The BCH Family of Storage Codes on Triangle-Free Graphs is of Unit Rate

Haihua Deng, Hexiang Huang, Guobiao Weng, and Qing Xiang*

ABSTRACT. Let Γ be a simple connected graph on n vertices, and let C be a code of length n whose coordinates are indexed by the vertices of Γ . We say that C is a *storage code* on Γ if for any codeword $c \in C$, one can recover the information on each coordinate of c by accessing its neighbors in Γ . The main problem here is to construct high-rate storage codes on triangle-free graphs. In this paper, we solve an open problem posed by Barg and Zémor in 2022, showing that the BCH family of storage codes is of unit rate. Furthermore, we generalize the construction of the BCH family and obtain more storage codes of unit rate on triangle-free graphs.

CONTENTS

| 1. Introduction | 1 |
|--|----|
| 2. The BCH family is of unit rate | 2 |
| 2.1. An upper bound | 2 |
| 2.2. Proof of Theorem 2 | 5 |
| 2.3. The ambient graphs of the BCH Family | 8 |
| 3. The generalized BCH family | 8 |
| 4. The case when $n = 2^r + 1$ | 10 |
| 5. Some cases of three-bit n by using a computer | 13 |
| Acknowledgments | 15 |
| References | 15 |

1. Introduction

A simple graph, also called a strict graph, is an unweighted, undirected graph containing no loops or multiple edges. A simple graph is said to be *connected* if there is a path between each pair of its vertices.

Let Γ be a simple connected graph on n vertices, and let C be a code of length n whose coordinates are indexed by the vertices of Γ . We say that C is a *storage code* on Γ if for any codeword $c \in C$, one can recover the information on each coordinate of c by accessing its neighbors in Γ . In 2014, Mazumdar [9, 10], Shanmugam and Dimakis [11] introduced storage codes on graphs independently. The concept of storage codes on graphs was introduced, in a different way, by the authors of [3] and [5]. Throughout this paper, we will only consider binary linear storage codes.

Key words and phrases. Cayley graphs, storage codes, the BCH family, the polynomial method.

^{*}Research partially supported by the National Natural Science Foundation of China Grant No. 12071206, 12131011, 12150710510, and the Sino-German Mobility Programme M-0157.

Suppose that Γ has *n* vertices, say v_1, v_2, \ldots, v_n . We define a storage code on Γ in the following way: let $A(\Gamma)$ be the adjacency matrix of Γ whose rows and columns are indexed by the vertices v_1, v_2, \ldots, v_n . Let $H := A(\Gamma) + I$ where *I* is the $n \times n$ identity matrix, and let *C* be the linear code over \mathbb{F}_2 with *H* as a parity-check matrix. Given a codeword $c = (c_{v_1}, c_{v_2}, \ldots, c_{v_n}) \in C$, we are able to recover any v_i^{th} entry of *c* by accessing its neighbors since the v_i^{th} row of *H* implies a linear equation, namely, $c_{v_i} = \sum_{v_j \in N(v_i)} c_{v_j}$, where $N(v_i)$ is the set of neighbors of v_i in Γ . The rate of a linear storage code *C*, denoted by R(C), is simply the ratio of its dimension to the dimension of the ambient space. If we have a family of storage codes $\{C_m\}$, where *m* is a parameter, assuming that $\lim_{m \to \infty} R(C_m)$ exists, then this limit is called the *rate* of the family.

Constructing a family of storage codes of unit rate is easy: let Γ_n be the complete graph on *n* vertices, and let C_n be the binary linear code defined by the equation $\sum_{i=1}^n x_{v_i} = 0$. Then C_n is a storage code on Γ with rate 1 - 1/n; hence the family $\{C_n\}$ is of unit rate.

In the above example, the graph used to obtain the storage code of rate close to one is very dense (in fact, as dense as possible), and contains a large number of cliques. It is therefore natural to consider the question of the largest attainable rate of storage codes on graphs that contain no cliques K_t ($t \ge 3$), i.e., triangle-free graphs.

Constructing storage codes of high rate on such graphs represents a challenge. A triangle-free graph with many edges does not necessarily give rise to a storage code of high rate. To see this, consider the complete bipartite graph $K_{t,t}$ which is triangle-free and quite dense, and a storage code C on it. Note that there are two independent vertex sets of $K_{t,t}$ and so for each vertex, we can recover the message on it from the ones in the other (vertex) independent set; hence $R(C) \leq 1/2$. In early studies [5], the authors had conjectured that for triangle-free graphs, R = 1/2 is the largest attainable rate value. Later on this conjecture was refuted in [3] by some sporadic examples.

Recently, the authors of [2] constructed four infinite families of storage codes on triangle-free graphs. They used the *Cayley graph method*: Let S be a subset of \mathbb{F}_2^r such that $0 \notin S$ and the sum of any three distinct vectors in S is nonzero. Then the resulting Cayley graph $\Gamma = \text{Cay}(\mathbb{F}_2^r, S)$ is triangle-free. Let $H := A(\Gamma) + I$ and C be the binary linear code defined by using H as its parity-check matrix. Then we obtain a storage code C on the triangle-free graph Γ . Using this method, a proper subset $S \subseteq \mathbb{F}_2^r$ will give rise to a triangle-free graph and a storage code on it. In their work, the Hamming family is of rate 3/4 and the BCH family shows the record of rate 0.8196. It remains unknown whether the BCH family can approach unit rate or not; this was left as an open problem in [2].

Subsequently, the authors of [1] and the authors of [7] presented the generalized Hamming family which could reach unit rate. In this paper, we solve the open problem about the BCH family, showing that the BCH family is indeed of unit rate. We also generalize the construction of the BCH family to obtain more storage codes on triangle-free graphs with rates approaching one.

2. The BCH family is of unit rate

2.1. An upper bound. In 2022, Barg and Zémor [2] presented a new family, the so-called BCH family, which can reach the rate of 0.8196. This data can be calculated by using a computer. It is left as an open problem whether the BCH family can reach unit rate or not. In order to understand why the BCH family exhibits a phenomenon of high rate, we use the *polynomial method* to investigate the intrinsic algebraic structure of the BCH family. As a consequence, we give an upper bound for the rank of the parity-check matrix of the BCH family, which shows that the BCH family is indeed of unit rate.

The BCH family of storage codes is constructed by using the Cayley graph method. We first recall the definition of Cayley graphs.

DEFINITION 1. Let G be a finite multiplicatively written group with identity element e, and let S be a subset of G such that $e \notin S$ and $S = S^{-1}$, where $S^{-1} = \{g^{-1} \mid g \in S\}$. The Cayley graph on G with connection set S, denoted by $\Gamma = \text{Cay}(G, S)$, is the graph with elements of G as vertices, two vertices $g_1, g_2 \in G$ are adjacent if and only if $g_1g_2^{-1} \in S$.

Now we are going to construct the Cayley graphs of the BCH family. Let $q = 2^m$ with $m \ge 1$ being an integer. The vertex set is given by $G = \mathbb{F}_q^2$ and the connection set is given by $S_m \setminus \{0\}$, where

$$S_m := \left\{ (a, a^3) \mid a \in \mathbb{F}_q \right\} \subseteq \mathbb{F}_q^2.$$

The graph is $\Gamma = \operatorname{Cay}(\mathbb{F}_q^2, S_m \setminus \{0\}).$

Let $H_m := A(\Gamma) + I$ and C_m be the binary linear code defined by using H_m as a parity-check matrix. Since each row of the parity-check matrix H_m for the storage code C_m can be regarded as a characteristic vector of a coset in $\{x + S_m \mid x \in \mathbb{F}_q^2\}$, we may call H_m the *coset matrix* of S_m in \mathbb{F}_q^2 . In order to better understand the structure of the matrix H_m , we will express the (x, y)-entry of H_m as the value of a polynomial evaluated at (x, y). More precisely, the coset matrix H_m over \mathbb{F}_2 can be formulated as

$$H_m = (a_{x,y})_{x,y \in \mathbb{F}_q^2},$$

where the (x, y)-entry is given by

$$a_{x,y} = \begin{cases} 1, & \text{if } x - y \in S_m \\ 0, & \text{otherwise.} \end{cases}$$

Next, we apply the *polynomial method* to investigate the rank of H_m . If we write $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{F}_q^2$, then $a_{x,y}$ can be expressed as the value of a polynomial g evaluated at (x, y):

$$a_{x,y} = \left((x_1 - y_1)^3 - (x_2 - y_2) \right)^{q-1} + 1$$

= $\left(x_1^3 + x_1^2 y_1 + x_1 y_1^2 + y_1^3 + x_2 + y_2 \right)^{q-1} + 1$
=: $g(x_1, x_2, y_1, y_2)$.

Let $W_m = (a_{x,y} + 1)_{x,y \in \mathbb{F}_q^2}$. Then $W_m = H_m + J$, where J is the all-one matrix, and so

$$\operatorname{rank}(H_m) - \operatorname{rank}(J) \le \operatorname{rank}(W_m) \le \operatorname{rank}(H_m) + \operatorname{rank}(J)$$

that is,

$$\operatorname{rank}(H_m) - 1 \le \operatorname{rank}(W_m) \le \operatorname{rank}(H_m) + 1.$$

Therefore, the matrix W_m has almost the same rank as that of H_m . We define the rate of a square matrix $A_{n\times n}$ to be the ratio of rank(A) to the size n; that is, $R(A) = \operatorname{rank}(A)/n$. That the BCH family is of unit rate is equivalent to saying that the rate of W_m converges to 0 as $m \to \infty$. Now the problem is reduced to computing the rank of W_m whose entry $a_{x,y} + 1$ is given by

$$h(x_1, x_2, y_1, y_2) = (x_1^3 + x_1^2 y_1 + x_1 y_1^2 + y_1^3 + x_2 + y_2)^{q-1}$$

The following proposition simplifies the question further by dropping some terms from the polynomial h.

PROPOSITION 1. Let $D_m = (f(x, y))_{x,y \in \mathbb{F}^2_{\sigma}}$, where

$$f(x_1, x_2, y_1, y_2) = (x_1^2 y_1 + x_1 y_1^2 + x_2 + y_2)^{q-1}$$

Then D_m has the same \mathbb{F}_2 -rank as that of W_m .

PROOF. Note that changing (x_1, x_2) to $(x_1, x_2 + x_1^3)$ is a permutation on \mathbb{F}_q^2 . Thus changing $h(x_1, x_2, y_1, y_2)$ to $f = h(x_1, x_2 + x_1^3, y_1, y_2 + y_1^3)$ is in fact a permutation on the rows and columns of W_m . The conclusion of the proposition now follows.

To find an upper bound on the rank of D_m , we first decompose D_m as the product of two matrices. Let

$$\Omega := \left\{ (l_1, l_2, l_3, l_4) \mid \sum_{i=1}^4 l_i = q - 1, 0 \le l_i \le q - 1, \forall i \right\}.$$

Then we can expand the polynomial f as follows:

$$f = (x_1^2 y_1 + x_1 y_1^2 + x_2 + y_2)^{q-1}$$

= $\sum_{(l_1, l_2, l_3, l_4) \in \Omega} {q-1 \choose l_1, l_2, l_3, l_4} x_1^{2l_1+l_2} x_2^{l_3} y_1^{l_1+2l_2} y_2^{l_4}$
= $\left[\cdots {\binom{q-1}{l_1, l_2, l_3, l_4}} x_1^{2l_1+l_2} x_2^{l_3} \cdots \right] \left[\begin{array}{c} \vdots \\ y_1^{l_1+2l_2} y_2^{l_4} \\ \vdots \end{array} \right],$

where the coordinates of the row/column vector are indexed by elements in Ω . Therefore we can write D_m as the product of two matrices

$$D_m = LR = \begin{bmatrix} \cdots & \binom{q-1}{l_1, l_2, l_3, l_4} x_1^{2l_1 + l_2} x_2^{l_3} & \cdots \end{bmatrix} \begin{bmatrix} \vdots \\ y_1^{l_1 + 2l_2} y_2^{l_4} \\ \vdots \end{bmatrix}$$

where the rows of L and columns of R are indexed by elements of \mathbb{F}_q^2 . Let N_m be the number of distinct nonzero monomials in L. That is,

$$N_m := \# \left\{ (2l_1 + l_2, l_3) \, \middle| \, \begin{pmatrix} q - 1 \\ l_1, l_2, l_3, l_4 \end{pmatrix} \equiv 1 \pmod{2} \right\}.$$

We then have an upper bound on $\operatorname{rank}(D_m)$:

(1)
$$\operatorname{rank}(D_m) \le \operatorname{rank}(L) \le N_m,$$

Using some counting techniques, we can give an explicit formula for N_m ; and hence obtain an upper bound on the rank of D_m . The obtained upper bound is good enough for us to show that the BCH family is of unit rate. We state the following theorem whose proof is postponed to the next subsection.

THEOREM 2. Let D_m be defined as above with $m \ge 1$ being an integer. Then

$$\operatorname{rank}(D_m) \le \frac{1+\sqrt{2}}{2}(2+\sqrt{2})^m,$$

and so

$$R(D_m) \le \frac{1+\sqrt{2}}{2} \left(\frac{2+\sqrt{2}}{4}\right)^m.$$

2.2. Proof of Theorem 2. Since the sequence of numbers N_m is defined by a property involving multinomial coefficients, we will use Lucas' theorem to analyse the behavior of N_m . Surprisingly, we can even compute the exact values of N_m .

Let n be a non-negative integer and p a prime. Suppose that the base p expansion of n is given by $n = n_k p^k + n_{k-1} p^{k-1} + \cdots + n_1 p + n_0$, where $0 \le n_i \le p-1$ for all i. We may use the abbreviation $n = \langle n_k n_{k-1} \cdots n_1 n_0 \rangle_p$ or $n = \langle n_k, n_{k-1}, \cdots, n_1, n_0 \rangle_p$. In the case where p = 2, we may drop the subscript p. We state Lucas' theorem as follows.

THEOREM 3 (Lucas' Theorem [4]). Let p be a prime, and express the non-negative integers n, l_1, l_2, \ldots, l_s in base p as

$$n = \langle n_k, n_{k-1}, \dots, n_1, n_0 \rangle_p; \quad l_i = \langle l_{i,k}, l_{i,k-1}, \dots, l_{i,1}, l_{i,0} \rangle_p,$$

where $n_j, l_{i,j} \in \{0, 1, \dots, p-1\}$ for $j = 0, 1, \dots, k$ and $i = 1, 2, \dots, s$. Then

$$\binom{n}{l_1, l_2, \dots, l_s} \equiv \prod_{j=0}^k \binom{n_j}{l_{1,j}, l_{2,j}, \dots, l_{s,j}} \pmod{p}$$

In the case where p = 2, we will drop (mod 2) to simplify notation. Before doing the actual computations, we will fix some notation as follows.

DEFINITION 2. Let a, b, c be non-negative integers. We write a + b < c, if the following conditions hold:

$$a_i + b_i \leq c_i \text{ for all } i = 0, \dots, k,$$

where $a = \langle a_k a_{k-1} \cdots a_1 a_0 \rangle_2, \ b = \langle b_k b_{k-1} \cdots b_1 b_0 \rangle_2, \ c = \langle c_k c_{k-1} \cdots c_1 c_0 \rangle_2.$

For $0 \leq s \leq q - 1$, define

$$B_s := \left\{ 2l_1 + l_2 \left| \begin{pmatrix} q - 1 \\ l_1, l_2, q - 1 - s, l_4 \end{pmatrix} \right| \equiv 1 \text{ for some } l_4 \right\}$$

and $b_s := |B_s|$. Note that the base 2 expansion of q-1 is $\langle \underbrace{11\cdots 1}_m \rangle$. By Theorem 3 we know

that $\binom{q-1}{l_1,l_2,q-1-s,l_4} \equiv 1 \pmod{2}$ if and only if the addition $l_1 + l_2 + (q-1-s) + l_4 = q-1$ involves no carries, which in turn is equivalent to $l_1 + l_2 < s$ and $l_4 = s - l_1 - l_2$. We now rewrite B_s as

$$B_s = \{2l_1 + l_2 \mid l_1 + l_2 \leqslant s\}.$$

Note that we have $N_m = \sum_{s=0}^{q-1} b_s$.

LEMMA 4. Let $s = \langle \alpha_1, \alpha_2, \dots, \alpha_n, \beta_1, \beta_2, \dots, \beta_k \rangle$. Then $B = B \times 2^k + B := \int r 2^k + t | r \in B \quad t \in B$

$$B_s = B_{s_1} \times 2^{\kappa} + B_{s_2} := \{ r2^{\kappa} + t \mid r \in B_{s_1}, t \in B_{s_2} \},\$$

where $s_1 = \langle \alpha_1, \alpha_2, \dots, \alpha_n \rangle$ and $s_2 = \langle \beta_1, \beta_2, \dots, \beta_k \rangle$.

PROOF. On the one hand, $B_s \subseteq B_{s_1} \times 2^k + B_{s_2}$. This can be seen as follows. Assume that $l_1 + l_2 < s$. By the division algorithm we write $l_1 = r_1 2^k + t_1, 0 \le t_1 < 2^k$, where the quotient and remainder, r_1, t_1 , are uniquely determined. Similarly for l_2 we obtain the quotient and the remainder, r_2, t_2 , respectively. As $l_1 + l_2 < s$, we have $r_1 + r_2 < s_1, t_1 + t_2 < s_2$ and thus $2l_1 + l_2 = 2(r_1 \times 2^k + t_1) + (r_2 \times 2^k + t_2) = (2r_1 + r_2) \times 2^k + (2t_1 + t_2) \in B_{s_1} \times 2^k + B_{s_2}$.

On the other hand, $B_s \supseteq B_{s_1} \times 2^k + B_{s_2}$: Assume $r_1 + r_2 < s_1, t_1 + t_2 < s_2$. Let $l_1 = r_1 \times 2^k + t_1, l_2 = r_2 \times 2^k + t_2$. Then $l_1 + l_2 < s$. So $(2r_1 + r_2) \times 2^k + (2t_1 + t_2) = 2(r_1 \times 2^k + t_1) + (r_2 \times 2^k + t_2) = 2l_1 + l_2 \in B_s$.

PROPOSITION 5. Let *i* be a positive integer. Then

$$b_{2^{i-1}-1} = 2^i - 1.$$

PROOF. Note that $B_{2^{i-1}-1} = \{2l_1 + l_2 \mid l_1 + l_2 \leq 2^{i-1} - 1\}$. Fixing $l_1 = 0$, we can take $l_2 = 0, 1, 2, \dots, 2^{i-1} - 1$, then $2l_1 + l_2 = 0, 1, 2, \dots, 2^{i-1} - 1$.

Let $l_1+l_2 = 2^{i-1}-1$. Then we have $l_1+l_2 < 2^{i-1}-1$ and $2l_1+l_2 = l_1+2^{i-1}-1$. As l_1 varies from 0 to $2^{i-1}-1$, $2l_1+l_2$ varies from $2^{i-1}-1$ to 2^i-2 . So $B_{2^{i-1}-1} = \{0, 1, 2, \dots, 2^i-2\}$. The claim now follows.

LEMMA 6. Let $s = \langle \alpha_1, \alpha_2, \dots, \alpha_n, 0, \beta_1, \beta_2, \dots, \beta_k \rangle$. Then

$$b_s = b_{s_1} b_{s_2},$$

where $s_1 = \langle \alpha_1, \alpha_2, \dots, \alpha_n \rangle$ and $s_2 = \langle \beta_1, \beta_2, \dots, \beta_k \rangle$.

PROOF. By Lemma 4 we have

$$B_s = B_{s_1} \times 2^{k+1} + B_{s_2}.$$

Note that for any $t \in B_{s_2}$, $t \le 2s_2 < 2^{k+1}$. Assume there are two pairs $(r_1, t_1), (r_2, t_2) \in B_{s_1} \times B_{s_2}$, such that $r_1 \times 2^{k+1} + t_1 = r_2 \times 2^{k+1} + t_2$. Then $(r_1 - r_2) \times 2^{k+1} + (t_1 - t_2) = 0$ and thus $r_1 - r_2 = t_1 - t_2 = 0$, i.e., $(r_1, t_1) = (r_2, t_2)$. Hence $\#B_s = \#(B_{s_1} \times 2^{k+1} + B_{s_2}) = \#(B_{s_1} \times B_{s_2})$.

EXAMPLE 2.1. By direct calculations, we have

| $b_0 = b_{2^0 - 1} = 2^1 - 1 = 1,$ | by Proposition 5 |
|--|-------------------------|
| $b_1 = b_{2^1 - 1} = 2^2 - 1 = 3,$ | by Proposition 5 |
| $b_2 = b_{\langle 10\rangle} = b_1 b_0 = 3,$ | by Lemma <mark>6</mark> |
| $b_3 = b_{2^2 - 1} = 2^3 - 1 = 7.$ | by Proposition 5 |

Thus $N_1 = b_0 + b_1 = 4$, $N_2 = b_0 + b_1 + b_2 + b_3 = 14$. We will use the initial values N_1, N_2 to determine the general formula of N_m in Theorem 8.

PROPOSITION 7. The sequence of numbers N_m satisfies:

$$N_m = \sum_{j=1}^{m+1} (2^j - 1) N_{m-j}, \quad m \ge 1,$$

where $N_0 = N_{-1} = 1$.

PROOF. Define

$$E^{(m)} := \left\{ t \mid 0 \le t \le 2^m - 1 \right\}$$

= $\left\{ \langle t_{m-1}, \dots, t_1, t_0 \rangle \mid t_i \in \{0, 1\}, \forall i \right\},$
$$E_j^{(m)} := \left\{ \langle \underbrace{1, 1, \dots, 1}_{j-1}, 0, t_{m-j-1}, \dots, t_1, t_0 \rangle \mid t_i \in \{0, 1\}, \forall i \right\}$$

= $\{2^{j-1} - 1\} \times 2^{m-j+1} + E^{(m-j)}, j = 1, \dots, m-1;$
$$E_m^{(m)} := \left\{ \langle \underbrace{11 \cdots 1}_{m-1} 0 \rangle \right\}, \quad E_{m+1}^{(m)} := \left\{ \langle \underbrace{11 \cdots 1}_{m} \rangle \right\}.$$

It is clear that $E^{(m)}$ is the disjoint union of $E_j^{(m)}$, $j = 1, \ldots, m+1$, namely $E^{(m)} = \bigcup_{j=1}^{m+1} E_j^{(m)}$. Note that $N_m = \sum_{t=0}^{2^m-1} b_s = \sum_{s \in E^{(m)}} b_s$. Then by Proposition 5 and Lemma

6 we have

$$N_{m} = \sum_{s \in E^{(m)}} b_{s} = \sum_{j=1}^{m-1} \sum_{s \in E_{j}^{(m)}} b_{s} + 2^{m} - 1 + 2^{m+1} - 1$$

$$= \sum_{j=1}^{m-1} \sum_{t \in E^{(m-j)}} b_{2^{j-1}-1} b_{t} + 2^{m} - 1 + 2^{m+1} - 1$$

$$= \sum_{j=1}^{m-1} b_{2^{j-1}-1} \sum_{t \in E^{(m-j)}} b_{t} + 2^{m} - 1 + 2^{m+1} - 1$$

$$= \sum_{j=1}^{m-1} b_{2^{j-1}-1} N_{m-j} + 2^{m} - 1 + 2^{m+1} - 1$$

$$= \sum_{j=1}^{m-1} (2^{j} - 1) N_{m-j} + 2^{m} - 1 + 2^{m+1} - 1$$

$$= \sum_{j=1}^{m+1} (2^{j} - 1) N_{m-j}.$$

| | п | |
|--|---|--|
| | | |
| | | |
| | | |

THEOREM 8. We have

(2)
$$N_m = \frac{1+\sqrt{2}}{2}(2+\sqrt{2})^m + \frac{1-\sqrt{2}}{2}(2-\sqrt{2})^m, \quad m \ge 0.$$

PROOF. By Proposition 7, we have

(3)
$$N_m = \sum_{j=1}^{m+1} (2^j - 1) N_{m-j}, \quad m \ge 1.$$

Replacing m by m-1, we get

(4)
$$N_{m-1} = \sum_{j=1}^{m} (2^{j} - 1) N_{m-1-j}$$
$$= \sum_{j=2}^{m+1} (2^{j-1} - 1) N_{m-j}, \quad m \ge 2,$$

Using (3) and (4), we obtain

$$N_m - 2N_{m-1} = N_{m-1} + \sum_{j=2}^{m+1} N_{m-j}, \quad m \ge 2.$$

It follows that

(5)
$$N_m = 3N_{m-1} + \sum_{j=2}^{m+1} N_{m-j}, \quad m \ge 2.$$

Again, replacing m by m-1, we get

(6)
$$N_{m-1} = 3N_{m-2} + \sum_{j=3}^{m+1} N_{m-j}, \quad m \ge 3,$$

Using (5) and (6), we obtain

$$N_m - N_{m-1} = 3N_{m-1} - 3N_{m-2} + N_{m-2}, \quad m \ge 3.$$

Consequently,

$$N_m = 4N_{m-1} - 2N_{m-2}, \quad m \ge 3.$$

Taking m = 2, we find that the initial values $N_0 = 1, N_1 = 4, N_2 = 14$ satisfy this linear recurrence relation. So the above recurrence holds whenever $m \ge 2$. Solving the linear recurrence we obtain

$$N_m = \frac{1+\sqrt{2}}{2}(2+\sqrt{2})^m + \frac{1-\sqrt{2}}{2}(2-\sqrt{2})^m, \quad m \ge 2.$$

We are now ready to give the proof of Theorem 2. Proof of Theorem 2. We have the following upper bound:

$$\operatorname{rank}(D_m) \le N_m = \frac{1+\sqrt{2}}{2}(2+\sqrt{2})^m + \frac{1-\sqrt{2}}{2}(2-\sqrt{2})^m$$
$$\le \frac{1+\sqrt{2}}{2}(2+\sqrt{2})^m.$$

Therefore,

$$R(D_m) = \frac{\operatorname{rank}(D_m)}{q^2} \le \frac{1+\sqrt{2}}{2} \left(\frac{2+\sqrt{2}}{4}\right)^m.$$

From the above upper bound on $R(D_m)$, we immediately see that the BCH family is of unit rate.

2.3. The ambient graphs of the BCH Family. For a given positive integer m, the graph of the BCH family is $\Gamma(V, E) = \operatorname{Cay}(\mathbb{F}_q^2, S_m \setminus \{0\})$, where $q = 2^m$. The number of vertices is $N = |V| = q^2 = 2^{2m}$. Note that Γ is a regular graph and each vertex has degree $|S_m| - 1 = q - 1 = 2^m - 1$, so the number of edges is

$$|E| = \frac{N(2^m - 1)}{2} = \frac{N(\sqrt{N} - 1)}{2} = O(N^{3/2}).$$

It is clear that Γ is simple. We claim that Γ is connected when m > 2. The proof is given in the next section.

We show that Γ is triangle-free: Let $a, b, c \in \mathbb{F}_q$ be distinct nonzero elements such that a+b+c=0. We claim that $a^3+b^3+c^3 \neq 0$. If not, then $c^3 = (a+b)^3 = a^3+a^2b+ab^2+b^3 = a^3+b^3$ and we obtain a = b, a contradiction. Hence the sum of any three distinct nonzero vectors in S_m is nonzero and thus Γ is triangle-free.

3. The generalized BCH family

Recall that in the BCH family, we investigate the coset matrix of S_m in \mathbb{F}_q^2 , where $q = 2^m$ and $S_m = \{(a, a^3) \mid a \in \mathbb{F}_q\}$. Now define

$$S_{n,m} := \{ (a, a^n) \mid a \in \mathbb{F}_q \} \subseteq \mathbb{F}_q^2,$$

where n is a fixed odd integer and $1 < n \leq q - 1$. Then we obtain the generalized BCH family F_n on the graph $\Gamma_{n,m} = \operatorname{Cay}(\mathbb{F}_q^2, S_{n,m} \setminus \{0\}).$

REMARK 3.1. In the above generalization, we require n to be odd. In fact, the matrix $H_{n,m}$ has the same rank as $H_{n/2,m}$ when n is even, where $H_{n,m}$ denotes the coset matrix of $S_{n,m}$ in \mathbb{F}_q^2 .

To prove that $\Gamma_{n,m}$ is connected, we need to show any vector in \mathbb{F}_q^2 is a sum of vectors in S_m . That is, viewing \mathbb{F}_q^2 as a 2*m*-dimensional \mathbb{F}_2 -vector space, we need to show that S_m contains a basis of \mathbb{F}_q^2 .

We now show that when m is large enough, the graph $\Gamma_{n,m}$ is connected. The following proof can be found in most coding theory textbooks. For more details, we refer the readers to [8].

THEOREM 9. Let n > 1 be an odd integer. If $2^{\frac{m}{2}} + 1 > n$, then S_m contains a \mathbb{F}_2 -basis for \mathbb{F}_a^2 ; and the graph $\Gamma_{n,m}$ is connected.

PROOF. Let $\mathbb{F}_q^* = \langle \alpha \rangle$. We claim that $\{(\alpha^k, \alpha^{nk}) \mid k = 0, 1, \dots, 2m - 1\}$ are linearly independent over \mathbb{F}_2 . Assume that $\sum_{k=0}^{2m-1} c_k \alpha^k = \sum_{k=0}^{2m-1} c_k \alpha^{nk} = 0$, where $c_k \in \mathbb{F}_2$. Let $g(x) = \sum_{k=0}^{2m-1} c_k x^k$. Then $g(\alpha) = g(\alpha^n) = 0$.

Let $p_1(x), p_2(x)$ be the minimal polynomials of α, α^n in $\mathbb{F}_2[x]$ respectively. We know that p_1, p_2 are irreducible polynomials. As α is a primitive element of \mathbb{F}_q , we have deg $p_1 = m$. Note that α and α^n are not conjugate to each other as n is odd and $1 < n < 2^{\frac{m}{2}} + 1 < q - 1$, so p_1, p_2 are coprime to each other.

Suppose that deg $p_2 = d$. We show that d = m: We know that d|m and $\alpha^n = \alpha^{n2^d}$, so $(2^m - 1)|n(2^d - 1)$ and thus $n \ge \frac{2^m - 1}{2^d - 1}$. Combining with the assumption that $2^{\frac{m}{2}} + 1 > n$, we have $2^{\frac{m}{2}} + 1 > \frac{2^m - 1}{2^d - 1}$, so $d > \frac{m}{2}$ and consequently d = m since d|m.

The polynomial g(x) should be a multiple of $p_1(x)p_2(x)$ since g(x) has the roots α, α^n . As the degree of $p_1(x)p_2(x)$ is 2m and g(x) cannot be of degree 2m, we deduce that g(x) is the zero polynomial. Hence the claim follows.

COROLLARY 10. Let r be a positive integer and $n = 2^r + 1$. Then the graph $\Gamma_{n,m}$ is connected if m > 2r.

For what values of n, will $\Gamma_{n,m}$ be triangle-free? The next lemma answers this question.

LEMMA 11. The graph $\Gamma_{n,m}$ is not triangle-free if and only if the equation $(x+1)^n+1 = x^n + 1$ has solution other than x = 0, 1.

PROOF. First note that $\Gamma_{n,m}$ is triangle-free if and only if the sum of any three nonzero distinct vectors in $S_{n,m}$ is nonzero. Assume that there are three distinct elements $a, b, c \in \mathbb{F}_q^*$ such that $a + b + c = a^n + b^n + c^n = 0$. Then $c^n = (a + b)^n = a^n + b^n$. Let $x = ab^{-1}$. We obtain $(x + 1)^n = x^n + 1$, where $x \neq 0, 1$. Hence the necessary condition holds.

To verify the sufficient condition, we assume that there exists an $x \neq 0, 1$ satisfying $(x+1)^n + 1 = x^n + 1$. Then a = x, b = 1, c = x + 1 are three distinct nonzero elements. The vectors $(a, a^n), (b, b^n), (c, c^n)$ will cause a triangle in the graph as the sum of them is zero.

PROPOSITION 12. Let $n = 2^r + 1$, where $r \ge 1$ is an integer. Then the graph $\Gamma_{n,m}$ is triangle-free if and only if gcd(r,m) = 1.

PROOF. We only need to consider the equation

$$(x+1)^{2^r+1} = x^{2^r+1} + 1,$$

 $x(x^{2^r-1}+1) = 0.$

The equation $x(x^{2^r-1}+1) = 0$ only has solutions x = 0, 1 if and only if $gcd(x(x^{2^r-1}+1), x^q+x) = x(x+1)$. That is

$$\begin{aligned} x(x+1) &= \gcd(x(x^{2^{r}-1}+1), x^{q}+x) \\ &= x \gcd(x^{2^{r}-1}+1, x^{2^{m}-1}+1) \\ &= x(x^{2^{\gcd(r,m)}-1}+1), \end{aligned}$$

which means that gcd(r, m) = 1.

PROPOSITION 13. Let $n = 2^r - 1$, where $r \ge 2$ is an integer. Then the graph $\Gamma_{n,m}$ is triangle-free if and only if gcd(r-1,m) = 1.

PROOF. We only need to consider the equation

$$(x+1)^{2^{r}-1} = x^{2^{r}-1} + 1,$$
$$\sum_{k=1}^{2^{r}-2} x^{k} = x \frac{1+x^{2^{r}-2}}{1+x} = 0.$$

The equation only has solutions x = 0, 1 if and only if $gcd(x^{2^{r-2}}+1, x^{q-1}+1) = x+1$, which implies that $gcd(2^{r}-2, 2^{m}-1) = gcd(2^{r-1}-1, 2^{m}-1) = 2^{gcd(r-1,m)}-1 = 1$ and thus gcd(r-1, m) = 1. The converse is also true.

4. The case when $n = 2^r + 1$

In this section, we are going to show that the generalized BCH family F_n is of unit rate provided that $n = 2^r + 1$, where r is a positive integer.

Let H_m be the coset matrix of S_m in \mathbb{F}_q^2 . Then we can formulate H_m as

$$H_m = (g(x,y))_{x,y \in \mathbb{F}_q^2},$$

where the (x, y)-entry is given by

$$g(x,y) = \left((x_1 + y_1)^{2^r + 1} + x_2 + y_2 \right)^{q-1} + 1.$$

By the same argument as in Section 2.1, we deduce that H_m has almost the same rank as that of the matrix $D_m := (f(x, y))_{x,y \in \mathbb{F}^2_r}$, where the (x, y)-entry is given by

$$f(x,y) = \left(x_1^{2^r}y_1 + x_1y_1^{2^r} + x_2 + y_2\right)^{q-1}.$$

Similarly, we have an upper bound: $\operatorname{rank}(D_m) \leq N_m$, where

$$N_m := \# \left\{ (2^r l_1 + l_2, l_3) \middle| \begin{pmatrix} q - 1 \\ l_1, l_2, l_3, l_4 \end{pmatrix} \equiv 1 \right\}$$
$$= \# \left\{ (2^r l_1 + l_2, l_3) \middle| l_1 + l_2 \leqslant q - 1 - l_3 \right\}.$$

We may redefine B_s in Section 2.2 by

$$B_s := \{2^r l_1 + l_2 \mid l_1 + l_2 \leqslant s\}$$

where $0 \le s \le q - 1$. Then

$$N_m = \sum_{s=0}^{2^m - 1} \# \{ 2^r l_1 + l_2 \mid l_1 + l_2 \leqslant q - 1 - s \}$$
$$= \sum_{s=0}^{2^m - 1} |B_{q-1-s}| = \sum_{s=0}^{2^m - 1} |B_s|.$$

For this more general definition of B_s , Lemma 4 still holds. We omit the proof since it is completely the same as before.

LEMMA 14. Let r be a positive integer and B_s defined above. Then

(1)
$$\sum_{s=0}^{2^{k}-1} |\{(l_{1}, l_{2}) \mid l_{1} + l_{2} \leq s\}| = 4^{k};$$

(2) $N_{k} = \sum_{s=0}^{2^{k}-1} |B_{s}| = 4^{k}, \text{ for } k = 0, 1, \dots, r;$
(3) $N_{r+1} = \sum_{s=0}^{2^{r+1}-1} |B_{s}| \leq 15 \times 4^{r-1}.$

PROOF. 1) We can classify all s between 0 and $2^k - 1$ by its weight, namely the number of 1s in its base 2 expansion. If the weight of s is i, then $\#\{(l_1, l_2) \mid l_1 + l_2 \leq s\} = 3^i$ since each pair (l_1, l_2) implies a distribution of each 1s to l_1, l_2 or l_3 , where $l_3 = s - l_1 - l_2$. We have

$$\sum_{s=0}^{2^{k}-1} \#\{(l_{1}, l_{2}) \mid l_{1}+l_{2} \leqslant s\} = \sum_{i=0}^{k} \binom{k}{i} 3^{i} = 4^{k}.$$

2) Assume $k \leq r$. For each $0 \leq s \leq 2^k - 1$, we have a map from $\{(l_1, l_2) \mid l_1 + l_2 \leq s\}$ to B_s , sending (l_1, l_2) to $2^r l_1 + l_2$. We want to show this is bijective map. It is clearly surjective, so we only need to show it is injective.

If there are two pairs $(l_1, l_2), (l'_1, l'_2)$ such that $2^r l_1 + l_2 = 2^r l'_1 + l'_2$, then $2^r (l_1 - l'_1) + l_2 - l'_2 = 0$ and thus $l_1 - l'_1 = l_2 - l'_2 = 0$ as $0 \le l_1, l_2, l'_1, l'_2 \le s \le 2^r - 1$. Hence the map is injective and thus bijective. The result follows by (1).

3) Notice that $B_1 = \{2^r l_1 + l_2 \mid l_1 + l_2 \leq 1\} = \{0, 1, 2^r\}$. We first calculate

$$\sum_{s=0}^{2^{r}-1} |B_{2s+1}| = \sum_{s=0}^{2^{r}-1} \#(B_s \times 2 + B_1)$$

$$= \sum_{s=0}^{2^{r}-1} \#(B_s \times 2 + \{0, 1, 2^r\})$$

$$= \sum_{s=0}^{2^{r}-1} \#(B_s \times 2 + 1) + \sum_{s=0}^{2^{r}-1} \#(B_s \times 2 + 2^r)$$

$$+ \sum_{s=0}^{2^{r}-1} \#(B_s \times 2) - \sum_{s=0}^{2^{r}-1} \#[(B_s \times 2) \cap (B_s \times 2 + 2^r)]$$

$$= 3N_r - \sum_{s=0}^{2^{r}-1} \#[(B_s \times 2) \cap (B_s \times 2 + 2^r)]$$

$$= 3N_r - \sum_{s=0}^{2^{r}-1} \#[B_s \cap (B_s + 2^{r-1})] := 3N_r - N'.$$

To determine the second term N', let $(l_1, l_2), (l'_1, l'_2)$ with $l_1 + l_2 \leq s, l'_1 + l'_2 \leq s$, where $0 \leq s \leq 2^r - 1$. Assume $2^r l_1 + l_2 = 2^r l'_1 + l'_2 + 2^{r-1}$ in the intersection. Then we obtain $2^r (l_1 - l'_1) + (l_2 - l'_2) = 2^{r-1}$. This equality holds if $l_1 - l'_1 = 0, l_2 - l'_2 = 2^{r-1}$. Hence $2^r l'_1 + l'_2 + 2^{r-1}$ is in the intersection if $2^{r-1} \leq s \leq 2^r - 1$ and $l'_1 + l'_2 \leq s - 2^{r-1}$. Then

$$N' \ge \sum_{s=2^{r-1}}^{2^{r-1}} |B_{s-2^{r-1}}| = \sum_{s=0}^{2^{r-1}-1} |B_s| = 4^{r-1}.$$

The value of N_{r+1} is given by

$$N_{r+1} = \sum_{s=0}^{2^{r+1}-1} |B_s| = \sum_{s=0}^{2^r-1} (|B_{2s}| + |B_{2s+1}|)$$
$$= \sum_{s=0}^{2^r-1} \#(B_s \times 2) + \sum_{s=0}^{2^r-1} |B_{2s+1}|$$
$$= N_r + 3N_r - N' \le 4^{r+1} - 4^{r-1} = 15 \times 4^{r-1}.$$

| | _ | _ | |
|--|---|---|--|

THEOREM 15. We have

s

$$N_m \le \left(\frac{15}{16}\right)^{\frac{m}{r+1}} 4^m.$$

PROOF. Assume m = t(r+1) + a, where $0 \le a < r+1$. Then for any $0 \le s \le 2^m - 1$, we have

$$= s_1 \times 2^{m-(r+1)} + s_2 \times 2^{m-2(r+1)} + \dots + s_t \times 2^{m-t(r+1)} + s_{t+1}$$

where $0 \le s_1, s_2, \ldots, s_t \le 2^{r+1} - 1$ and $0 \le s_{t+1} \le 2^a - 1$, which implies an expansion

$$B_s = \sum_{i=1}^{t} B_{s_i} \times 2^{m-i(r+1)} + B_{s_{t+1}},$$

and thus

$$|B_s| \le \prod_{i=1}^t |B_{s_i}| \cdot |B_{s_{t+1}}|.$$

Applying the above inequality, we obtain

$$N_{m} = \sum_{s=0}^{2^{m}-1} |B_{s}|$$

$$\leq \sum_{s_{1},\dots,s_{t}=0}^{2^{r+1}-1} \sum_{s_{t+1}=0}^{2^{a}-1} \left(\prod_{i=1}^{t} |B_{s_{i}}| \cdot |B_{s_{t+1}}| \right)$$

$$= \prod_{i=1}^{t} \left(\sum_{s_{i}=0}^{2^{r+1}-1} |B_{s_{i}}| \right) \cdot \sum_{s_{t+1}=0}^{2^{a}-1} |B_{s_{t+1}}|$$

$$= N_{r+1}^{t} N_{a} \leq (15 \times 4^{r-1})^{t} 4^{a}$$

$$= \left(\frac{15}{16} \right)^{t} 4^{m} \leq \left(\frac{15}{16} \right)^{\frac{m}{r+1}} 4^{m}.$$

The proof is now complete.

COROLLARY 16. The rate $R(D_m)$ converges to 0; so the generalized BCH family F_n (with $n = 2^r + 1$) is of unit rate.

PROOF. We denote the rate of D_m by R_m . Then $R_m = \operatorname{rank}(D_m)/4^m \le N_m/4^m$ and thus

$$R_m \le \left(\frac{15}{16}\right)^{\frac{m}{r+1}}$$

Hence $\{R_m\}$ converges to 0 as m goes to infinity.

5. Some cases of three-bit n by using a computer

DEFINITION 3. Let A, B be two matrices, say $A = (a_{i_1,j_1})_{m_1 \times n_1}, B = (b_{i_2,j_2})_{m_2 \times n_2}$. Then the tensor product of two matrices is $A \otimes B := (a_{i_1,j_1}B)_{m_1 \times n_1}$, namely each block entry is the product of the matrix B and an entry of A.

Note that the entry in the $((i_1 - 1)m_1 + i_2)^{th}$ row and the $((j_1 - 1)n_1 + j_2)^{th}$ column of $A \otimes B$ is $a_{i_1,j_1}b_{i_2,j_2}$.

DEFINITION 4. Let A, B be two matrices of the same size, say $A = (a_{i_1,j_1})_{m \times n}, B = (b_{i_2,j_2})_{m \times n}$. Then the Hadamard product of two matrices is $A \circ B := (a_{i,j}b_{i,j})_{m \times n}$.

THEOREM 17 ([6]). Let A, B be two matrices. Then

$$\operatorname{rank}(A \otimes B) = \operatorname{rank}(A)\operatorname{rank}(B).$$

COROLLARY 18. Let A, B be two $m \times n$ matrices. Then

 $\operatorname{rank}(A \circ B) \le \operatorname{rank}(A)\operatorname{rank}(B).$

PROOF. By Theorem 17, it suffices to show that $A \circ B$ is a submatrix of $A \otimes B$. Let $R = \{(i-1)m + i \mid i = 1, 2, ..., m\}, L = \{(j-1)m + j \mid j = 1, 2, ..., n\}$ and $C = A \otimes B(L \times R)$. Then we have

$$C(i,j) = A \otimes B((i-1)m+i, (j-1)n+j)$$

= $a_{i,j}b_{i,j} = A \circ B(i,j).$

The proof is now complete.

As used previously, $(h(x_1, x_2, y_1, y_2))_{(x_1, x_2), (y_1, y_2) \in \mathbb{F}_q^2}$ denotes the matrix in which each $((x_1, x_2), (y_1, y_2))$ -entry is $h(x_1, x_2, y_1, y_2)$. When there is no ambiguity regarding the finite field \mathbb{F}_q , we can omit the subscript and simply write the matrix as $(h(x_1, x_2, y_1, y_2))$ or (h). Below we always assume that the finite field has characteristic 2. We have the following lemma.

LEMMA 19. Let i be a non-negative integer. Then

$$\operatorname{rank}((h(x_1, x_2, y_1, y_2))) = \operatorname{rank}((h(x_1, x_2, y_1, y_2)^{2^{\epsilon}})).$$

PROOF. Note that $h(x_1, x_2, y_1, y_2)^{2^i} = h(x_1^{2^i}, x_2^{2^i}, y_1^{2^i}, y_2^{2^i})$. Furthermore, this expression represents a permutation of both the rows and columns. Thus the result follows. \Box

PROPOSITION 20. Let $d(x_1, x_2, y_1, y_2) \in \mathbb{F}_q[x_1, x_2, y_1, y_2]$ and t a positive integer. Then for any integer m > t, we have

$$\operatorname{rank}((d^{2^m-1})) \le c \cdot \left(\operatorname{rank}((d^{2^t-1}))\right)^{\frac{m}{t}}.$$

where $c = \max\{\operatorname{rank}((d^{2^{i}-1})) \mid 0 \le i < t\}$ only depends on t.

PROOF. Suppose that $m = k \cdot t + r$, where $0 \le r < t$. Then we can write

$$2^{m} - 1 = \sum_{j=0}^{k-1} (2^{t} - 1)2^{j+r} + 2^{r} - 1.$$

Hence,

$$\operatorname{rank}((d^{2^{m}-1})) \leq \operatorname{rank}((d^{2^{r}-1})) \prod_{j=0}^{k-1} \operatorname{rank}((d^{(2^{t}-1)\times 2^{j+r}})) \leq c \cdot \prod_{j=0}^{k-1} \operatorname{rank}((d^{2^{t}-1})) = c \cdot \left(\operatorname{rank}((d^{2^{t}-1}))\right)^{k} \leq c \cdot \left(\operatorname{rank}((d^{2^{t}-1}))\right)^{\frac{m}{t}},$$

where $c = \max\{ \operatorname{rank}((d^{2^{i}-1})) \mid 0 \le i < t \}.$

REMARK 5.1. The above theorem tells us that the rank of $A_m = (d^{2^t-1})_{\mathbb{F}^2_{2m} \times \mathbb{F}^2_{2m}}$ will give an upper bound for the rank of (d^{2^m-1}) . However, the matrix A_m is changing as m increases. We next show that rank (A_m) would not change when m is sufficiently large.

DEFINITION 5 (Rank of a polynomial). Assume that a polynomial $h \in \mathbb{F}_2[x_1, x_2, y_1, y_2]$, say

$$h = \sum_{i_1, i_2, j_1, j_2} a_{i_1, i_2, j_1, j_2} x_1^{i_1} x_2^{i_2} y_1^{j_1} y_2^{j_2},$$

where $a_{i_1,i_2,j_1,j_2} \in \mathbb{F}_2$. Then the coefficient matrix of h, the rows indexed by (i_1, i_2) and the columns indexed by (j_1, j_2) , is $M_h = (a_{i_1,i_2,j_1,j_2})$. The rank of the polynomial h is the rank of its coefficient matrix rank (M_h) , and it will be denoted by rank(h).

LEMMA 21. Let $h \in \mathbb{F}_2[x_1, x_2, y_1, y_2]$. Assume that $d = \max\{\deg_{x_1} h, \deg_{x_2} h, \deg_{y_1} h, \deg_{y_2} h\}$, where $\deg_{x_1} h$ is the degree of h in variable x_1 . If q > d, then

$$\operatorname{rank}((h)_{\mathbb{F}^2_q \times \mathbb{F}^2_q}) = \operatorname{rank}(h)$$

PROOF. Suppose that

$$h = \sum_{i_1, i_2, j_1, j_2 = 0}^{q-1} a_{i_1, i_2, j_1, j_2} x_1^{i_1} x_2^{i_2} y_1^{j_1} y_2^{j_2}.$$

Then we have

$$(h)_{\mathbb{F}_q^2 \times \mathbb{F}_q^2} = LM_hR$$

$$= \begin{bmatrix} \cdots & x_1^{i_1} x_2^{i_2} & \cdots \end{bmatrix} \begin{bmatrix} \vdots & \vdots & \vdots \\ \cdots & a_{i_1, i_2, j_1, j_2} & \cdots \\ \vdots & \vdots & \end{bmatrix} \begin{bmatrix} \vdots \\ y_1^{j_1} y_2^{j_2} \\ \vdots \end{bmatrix},$$

where the rows of L and the columns of R are indexed by elements of \mathbb{F}_q^2 . Note that the matrices L, R are invertible, so $\operatorname{rank}((h)) = \operatorname{rank}(M_h) = \operatorname{rank}(h)$.

We now consider the generalized BCH family F_n . In the following, we denote $d = (x_1 + y_1)^n + x_2 + y_2$, $f = d^{q-1}$.

THEOREM 22. If there exists a positive integer t such that

$$\operatorname{rank}(d^{2^t-1}) < 4^t,$$

then the generalized BCH family F_n is of unit rate.

PROOF. When m satisfies $2^m > n(2^t - 1)$, by Proposition 20 and Lemma 21 we have

$$\operatorname{rank}((f)_{\mathbb{F}_{2m}^{2} \times \mathbb{F}_{2m}^{2}}) = \operatorname{rank}((d^{2^{m}-1})_{\mathbb{F}_{2m}^{2} \times \mathbb{F}_{2m}^{2}})$$
$$\leq c \cdot \left(\operatorname{rank}((d^{2^{t}-1})_{\mathbb{F}_{2m}^{2} \times \mathbb{F}_{2m}^{2}})\right)^{\frac{m}{t}}$$
$$= c \cdot \left(\operatorname{rank}(d^{2^{t}-1})\right)^{\frac{m}{t}}$$
$$\leq c \cdot (4^{t}-1)^{\frac{m}{t}}$$
$$\frac{\operatorname{rank}((f))}{4^{m}} \leq \frac{c \cdot (4^{t}-1)^{\frac{m}{t}}}{4^{m}} = c \cdot \left(\frac{4^{t}-1}{4^{t}}\right)^{\frac{m}{t}}$$

Hence the evaluation matrix $(f)_{\mathbb{F}_q^2 \times \mathbb{F}_q^2}$ is of low rank and thus the parity-check matrix of the generalized BCH family F_n is also of low rank. We are done.

We can use a computer to search for the smallest t such that the rank of the polynomial $d^{2^{t}-1}$ is strictly smaller than 4^{t} . For instance, using Magma, we know that $\operatorname{rank}(d^{2^{6}-1}) = 3256 < 4096 = 4^{6}$ for F_{7} , $\operatorname{rank}(d^{2^{7}-1}) = 15018 < 16384 = 4^{7}$ for F_{11} , and $\operatorname{rank}(d^{2^{7}-1}) = 14442 < 16384 = 4^{7}$ for F_{13} . Therefore, we obtain the following result.

COROLLARY 23. The generalized BCH families F_7 , F_{11} and F_{13} are all of unit rate. \Box

Acknowledgments

We would like to thank Sihuang Hu for bringing the open problem of Barg and Zémor to our attention. We also thank Zhen Jia for his help with computer programming.

References

- A. Barg, M. Schwartz, and L. Yohananov, "Storage codes on triangle-free graphs with asyptotically unit rate," arXiv:2212.12117v1.
- [2] A. Barg and G. Zémor, "High-rate storage codes on triangle-free graphs," IEEE Trans. Inf. Theory, vol. 68, no. 12, pp. 7787–7797, Dec. 2022.
- [3] P. Cameron, A. Dang, and S. Riis, "Guessing games on triangle-free graphs," *Electron. J. Combina-torics*, vol. 23, no. 1, p. P1.48, Mar. 2016.
- [4] C. Chao and M. Zhang, "On Multinomial Coefficients Modulo a Prime," European Journal of Combinatorics, vol. 9, no. 1, pp. 23-26, Jan. 1988.
- [5] D. Christofides and K. Markström, "The guessing number of undirected graphs," *Electron. J. Combinatorics*, vol. 18, no. 1, p. P192, Sep, 2011.
- [6] R. A. Horn and C. R. Johnson, *Matrix analysis*. In Statistical Inference for Engineers and Data Scientists, 1985.
- [7] H. Huang and Q. Xiang, "Construction of storage codes of rates approaching one on triangle-free graphs," arXiv:2301.01668v2.
- [8] S. Ling and C. Xing, Coding Theory: A First Course. Cambridge University Press, 2004.
- [9] A. Mazumdar, "Storage capacity of repairable networks," *IEEE Trans. Inf. Theory*, vol. 61, no. 11, pp. 5810–5821, Nov. 2015.
- [10] A. Mazumdar, A. Mcgregor, and S. Vorotnikova, "Storage capacity as an information-theoretic analogue of vertex cover," 2017 IEEE Int. Symp. Inf. Theory (ISIT), pp. 2810–2814, Jun. 2017.
- [11] K. Shanmugam and A. G. Dimakis, "Bounding multiple unicasts through index coding and locally repairable codes," 2014 IEEE Int. Symp. Inf. Theory, pp. 296–300, Jun. 2014.

Department of Mathematics, Southern University of Science and Technology, Shenzhen 518055, China

Email address: 12131225@mail.sustech.edu.cn

Department of Mathematics, Southern University of Science and Technology, Shenzhen 518055, China

Email address: hhxiang1999@foxmail.com

School of Mathematical Sciences, Dalian University of Technology, Dalian 116024, China

Email address: weng@dlut.edu.cn

Department of Mathematics, Southern University of Science and Technology, Shenzhen 518055, China

Email address: xiangq@sustech.edu.cn

16