arXiv:1902.10766v1 [math.FA] 27 Feb 2019

BOUNDEDNESS OF WEIGHTED ITERATED HARDY-TYPE OPERATORS INVOLVING SUPREMA FROM WEIGHTED LEBESGUE SPACES INTO WEIGHTED CESÀRO FUNCTION SPACES

R.CH. MUSTAFAYEV AND N. BİLGİÇLİ

Abstract. In this paper the boundedness of the weighted iterated Hardy-type operators $T_{u,b}$ and $T^*_{u,b}$ involving suprema from weighted Lebesgue space $L_p(v)$ into weighted Cesàro function spaces $\mathrm{Ces}_q(w,a)$ are characterized. These results allow us to obtain the characterization of the boundedness of the supremal operator R_u from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions. For the convenience of the reader, we formulate the statement on the boundedness of the weighted Hardy operator $P_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions. Under additional condition on u and b, we are able to characterize the boundedness of weighted iterated Hardy-type operator $T_{u,b}$ involving suprema from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions. At the end of the paper, as an application of obtained results, we calculate the norm of the fractional maximal function M_γ from $\Lambda^p(v)$ into $\Gamma^q(w)$.

1. Introduction

Many Banach spaces which play an important role in functional analysis and its applications are obtained in a special way: the norms of these spaces are generated by positive sublinear operators and by L_p-norms.

In connection with Hardy and Copson operators

$$(Pf)(x) := \frac{1}{x}\int_0^x f(t)\,dt \qquad \text{and} \qquad (Qf)(x) := \int_x^\infty \frac{f(t)}{t}\,dt, \qquad (x>0),$$

the classical Cesàro function space

$$\mathrm{Ces}(p) := \left\{f : \|f\|_{\mathrm{Ces}(p)} := \left(\int_0^\infty \left(\frac{1}{x}\int_0^x |f(t)|\,dt\right)^p dx\right)^{\frac{1}{p}} < \infty\right\},$$

and the classical Copson function space

$$\mathrm{Cop}(p) := \left\{f : \|f\|_{\mathrm{Cop}(p)} := \left(\int_0^\infty \left(\int_x^\infty \frac{|f(t)|}{t}\,dt\right)^p dx\right)^{\frac{1}{p}} < \infty\right\},$$

where $1 < p \le \infty$, with the usual modifications if $p = \infty$, are of interest.

The classical Cesàro function spaces $\mathrm{Ces}(p)$ have been introduced in 1970 by Shiue [48]. These spaces have been defined analogously to the Cesàro sequence spaces that appeared two years earlier in [40] when the Dutch Mathematical Society posted a problem to find a representation of their dual spaces. In 1971 Leibowitz proved that $\mathrm{ces}_1 = \{0\}$ and for $1 < q < p \le \infty$, ℓ_p and ces_q sequence spaces are proper subspaces of ces_p [32]. The problem posted [40] was resolved by Jagers [28] in 1974 who gave an explicit isometric description of the dual of Cesàro sequence space. In [51], Sy, Zhang and Lee gave a description of dual spaces of $\mathrm{Ces}(p)$ spaces based on Jagers' result. In 1996 different, isomorphic description due to Bennett appeared in [4]. In [4, Theorem 21.1] Bennett observes that the classical Cesàro function space and the classical Copson function space coincide for $p > 1$. He also derives estimates for the norms of the corresponding inclusion operators. The same result, with different estimates, is due to Boas [7], who in fact obtained the integral analogue of the Askey-Boas Theorem [6, Lemma 6.18] and [1]. These results generalized in [27] using the blocking technique. In [2] they investigated dual spaces for $\mathrm{Ces}(p)$ for $1 < p < \infty$. Their description can be viewed as being analogous to one given for sequence spaces in [4]. For a long time, Cesàro function spaces have not attracted a lot of attention contrary to their sequence counterparts. In fact there is quite rich literature concerning different topics studied in Cesàro sequence spaces as for instance in [11–15]. However, recently in a series of papers, Astashkin and Maligranda started to study the structure of Cesàro

2010 *Mathematics Subject Classification.* 46E30, 26D10, 42B25, 42B35.

Key words and phrases. weighted iterated Hardy operators involving suprema, Cesàro function spaces, fractional maximal functions, classical Lorentz spaces.

function spaces. Among others, in [2] they investigated dual spaces for $\mathrm{Ces}(p)$ for $1 < p < \infty$. Their description can be viewed as being analogous to one given for sequence spaces in [4] (For more detailed information about history of classical Cesàro spaces see recent survey paper [3]).

Throughout the paper we assume that $I := (a, b) \subseteq (0, \infty)$. By $\mathfrak{M}(I)$ we denote the set of all measurable functions on I. The symbol $\mathfrak{M}^+(I)$ stands for the collection of all $f \in \mathfrak{M}(I)$ which are non-negative on I, while $\mathfrak{M}^{+,\downarrow}(I)$ is used to denote the subset of those functions which are non-increasing on I, respectively. A weight is a function $v \in \mathfrak{M}^+(0, \infty)$ such that $0 < V(x) < \infty$ for all $x \in (0, \infty)$, where

$$V(x) := \int_0^x v(t)\, dt.$$

The family of all weight functions (also called just weights) on $(0, \infty)$ is given by $\mathcal{W}(0, \infty)$.

For $p \in (0, \infty]$ and $w \in \mathfrak{M}^+(I)$, we define the functional $\|\cdot\|_{p,w,I}$ on $\mathfrak{M}(I)$ by

$$\|f\|_{p,w,I} := \begin{cases} \left(\int_I |f(x)|^p w(x)\, dx\right)^{1/p} & \text{if} \quad p < \infty \\ \operatorname{ess\,sup}_I |f(x)| w(x) & \text{if} \quad p = \infty. \end{cases}$$

If, in addition, $w \in \mathcal{W}(I)$, then the weighted Lebesgue space $L^p(w, I)$ is given by

$$L^p(w, I) = \{f \in \mathfrak{M}(I) : \ \|f\|_{p,w,I} < \infty\}$$

and it is equipped with the quasi-norm $\|\cdot\|_{p,w,I}$.

When $I = (0, \infty)$, we write $L^p(w)$ instead of $L^p(w, (0, \infty))$.

We adopt the following usual conventions.

Convention 1.1. We adopt the following conventions:

- Throughout the paper we put $0 \cdot \infty = 0$, $\infty/\infty = 0$ and $0/0 = 0$.
- If $p \in [1, +\infty]$, we define p' by $1/p + 1/p' = 1$.
- If $0 < q < p < \infty$, we define r by $1/r = 1/q - 1/p$.
- If $I = (a, b) \subseteq \mathbb{R}$ and g is monotone function on I, then by $g(a)$ and $g(b)$ we mean the limits $\lim_{x \to a+} g(x)$ and $\lim_{x \to b-} g(x)$, respectively.

Throughout the paper, we always denote by c and C a positive constant, which is independent of main parameters but it may vary from line to line. However a constant with subscript or superscript such as c_1 does not change in different occurrences. By $a \lesssim b$, $(b \gtrsim a)$ we mean that $a \leq \lambda b$, where $\lambda > 0$ depends on inessential parameters. If $a \lesssim b$ and $b \lesssim a$, we write $a \approx b$ and say that a and b are equivalent.

Unless a special remark is made, the differential element dx is omitted when the integrals under consideration are the Lebesgue integrals.

The weighted Cesàro and Copson function spaces are defined as follows:

Definition 1.2. Let $0 < p \leq \infty$, $u \in \mathfrak{M}^+(0, \infty)$ and $v \in \mathcal{W}(0, \infty)$. The weighted Cesàro and Copson spaces are defined by

$$\mathrm{Ces}_p(u, v) := \left\{ f \in \mathfrak{M}^+(0, \infty) : \|f\|_{\mathrm{Ces}_p(u,v)} := \left\| \|f\|_{1,v,(0,\cdot)} \right\|_{p,u,(0,\infty)} < \infty \right\},$$

and

$$\mathrm{Cop}_p(u, v) := \left\{ f \in \mathfrak{M}^+(0, \infty) : \|f\|_{\mathrm{Cop}_p(u,v)} := \left\| \|f\|_{1,v,(\cdot,\infty)} \right\|_{p,u,(0,\infty)} < \infty \right\},$$

respectively.

When $v \equiv 1$ on $(0, \infty)$, we simply write $\mathrm{Ces}_p(u)$ and $\mathrm{Cop}_p(u)$ instead of $\mathrm{Ces}_p(u, v)$ and $\mathrm{Cop}_p(u, v)$, respectively.

Recall that $\mathrm{Ces}_p(u, v)$ and $\mathrm{Cop}_p(u, v)$ are contained in the scale of weighted Cesàro and Copson function spaces $\mathrm{Ces}_{p,q}(u, v)$ and $\mathrm{Cop}_{p,q}(u, v)$ defined in [22]. Obviously, $\mathrm{Ces}(p) = \mathrm{Ces}_p(x^{-1})$ and $\mathrm{Cop}(p) = \mathrm{Cop}_p(x^{-1})$. In [29], Kamińska and Kubiak computed the dual norm of the Cesàro function space $\mathrm{Ces}_p(u)$, generated by $1 < p < \infty$ and an arbitrary positive weight u. A description presented in [29] resembles the approach of Jagers [28] for sequence spaces.

Let $u \in \mathcal{W}(0,\infty) \cap C(0,\infty)$, $b \in \mathcal{W}(0,\infty)$ and $B(t) := \int_0^t b(s)\,ds$. Assume that b is a weight such that $b(t) > 0$ for a.e. $t \in (0,\infty)$. The weighted iterated Hardy-type operators involving suprema $T_{u,b}$ and $T^*_{u,b}$ are defined at $g \in \mathfrak{M}^+(0,\infty)$ by

$$(T_{u,b}g)(t) := \sup_{t\le\tau} \frac{u(\tau)}{B(\tau)} \int_0^\tau g(y)b(y)\,dy, \qquad t \in (0,\infty),$$

$$(T^*_{u,b}g)(t) := \sup_{t\le\tau} \frac{u(\tau)}{B(\tau)} \int_\tau^\infty g(y)b(y)\,dy, \qquad t \in (0,\infty).$$

Such operators have been found indispensable in the search for optimal pairs of rearrangement-invariant norms for which a Sobolev-type inequality holds (cf. [30]). They constitute a very useful tool for characterization of the associate norm of an operator-induced norm, which naturally appears as an optimal domain norm in a Sobolev embedding (cf. [38], [39]). Supremum operators are also very useful in limiting interpolation theory as can be seen from their appearance for example in [18], [17], [16], [45]. Recall that $T_{u,b}$ successfully controls non-increasing rearrangements of wide range of maximal functions (see, for instance, [34] and references therein).

It was shown in [23] that for every $h \in \mathfrak{M}^+(0,\infty)$ and $t \in (0,\infty)$

$$(T_{u,b}h)(t) = (T_{\bar{u},b}h)(t),$$

where

$$\bar{u}(t) := B(t) \sup_{t\le\tau} \frac{u(\tau)}{B(\tau)}, \qquad t \in (0,\infty).$$

Moreover, if the condition

$$\sup_{0<t<\infty} \frac{u(t)}{B(t)} \int_0^t \frac{b(\tau)}{u(\tau)}\,d\tau < \infty. \tag{1.1}$$

holds, then for all $f \in \mathfrak{M}^{+,\downarrow}(0,\infty)$,

$$(T_{u,b}f)(t) \approx (R_u f)(t) + (P_{\bar{u},b}f)(t), \qquad t \in (0,\infty), \tag{1.2}$$

where the supremal operator R_u and the weighted Hardy operator $P_{u,b}$ are defined for $h \in \mathfrak{M}^+(0,\infty)$ and $t \in (0,\infty)$ by

$$(R_u h)(t) := \sup_{t\le\tau} u(\tau)h(\tau),$$

$$(P_{u,b}h)(t) := \frac{u(t)}{B(t)} \int_0^t h(\tau)b(\tau)\,d\tau,$$

respectively.

Recall that the boundedness of R_u from $L^p(v)$ into $L^q(w)$ on the cone of monotone non-increasing functions, that is, the validity of the inequality

$$\|R_u f\|_{L^q(w)} \le C\,\|f\|_{L^p(v)}, \qquad f \in \mathfrak{M}^{+,\downarrow}(0,\infty) \tag{1.3}$$

was completely characterized in [23] in the case $0 < p \le q < \infty$. In the case $0 < q < p < \infty$, [23] provides solution when u is equivalent to a non-decreasing function on $(0,\infty)$. The complete solution of inequality (1.3) using a certain reduction method was presented in [21]. Another solution of (1.3) was obtained in [31].

Note that inequality

$$\|P_{u,b}(f)\|_{q,w,(0,\infty)} \le c\|f\|_{p,v,(0,\infty)}, \qquad f \in \mathfrak{M}^{+,\downarrow}(0,\infty) \tag{1.4}$$

was considered by many authors and there exist several characterizations of this inequality (see, papers [5, 8, 9, 19, 20, 26]).

The complete characterizations of inequality

$$\|T_{u,b}f\|_{q,w,(0,\infty)} \le C\|f\|_{p,v,(0,\infty)}, \qquad f \in \mathfrak{M}^{+,\downarrow}(0,\infty) \tag{1.5}$$

for $0 < q \le \infty$, $0 < p \le \infty$ were given in [21] and [34]. Inequality (1.5) was characterized in [23, Theorem 3.5] under condition (1.1). Note that the case when $0 < p \le 1 < q < \infty$ was not considered in [23]. It is also worth to mention that in the case when $1 < p < \infty$, $0 < q < p < \infty$, $q \ne 1$ [23, Theorem 3.5] contains only discrete condition. In [25] the new reduction theorem was obtained when $0 < p \le 1$, and this technique allowed to characterize inequality (1.5)

when $b \equiv 1$, and in the case when $0 < q < p \le 1$, [25] contains only discrete condition. Using the results in [41–44], another characterization of (1.5) was obtained in [50] and [46].

In this paper we investigate the boundedness of $T_{u,b}$ and $T^*_{u,b}$ from the weighted Lebesgue spaces $L_p(v)$ into the weighted Cesàro spaces $\mathrm{Ces}_q(w,a)$, when $1 < p, q < \infty$ (see, Theorems 3.1 and 3.3). These results allow us to obtain the characterization of the boundedness of R_u from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions (see, Theorem 4.1). For the convenience of the reader, we formulate the statement on the boundedness of $P_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions (see, Theorem 5.1). In view of (1.2), we are able to characterize the boundedness of $T_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions (see, Theorem 6.1). At the end of the paper, as an application of obtained results, we calculate the norm of the fractional maximal function M_γ from $\Lambda^p(v)$ into $\Gamma^q(w)$.

The paper is organized as follows. We start with formulations of "an integration by parts" formula in Section 2. The boundedness results for $T_{u,b}$ and $T^*_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ are presented in Section 3. The characterizations of the boundedness of R_u, $P_{u,b}$ and $T_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions are given in Sections 4, 5 and 6, respectively. Finally, the obtained in previous sections results are applied to calculate the norm of the operator $M_\gamma : \Lambda_p(v) \to \Gamma_q(w)$ in Section 7.

2. "An integration by parts" formula

We recall the following "an integration by parts" formula. For the convenience of the reader we give the proof here (cf. [49, Lemma, p. 176]).

Theorem 2.1. *Let $\alpha > 0$. Let g be a non-negative function on $(0,\infty)$ such that $0 < \int_0^t g < \infty$, $t > 0$ and let f be a non-negative non-increasing right-continuous function on $(0,\infty)$. Then*

$$A_1 := \int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t)[f(t) - \lim_{t\to+\infty} f(t)]\,dt < \infty \quad \Longleftrightarrow \quad A_2 := \int_{(0,\infty)} \Big(\int_0^t g\Big)^{\alpha+1} d[-f(t)] < \infty.$$

Moreover, $A_1 \approx A_2$.

Proof. Assume at first that $\lim_{t\to+\infty} f(t) = 0$. Let

$$A_1 = \int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t) f(t)\,dt < \infty.$$

Then

$$\int_0^x \Big(\int_0^t g\Big)^\alpha g(t) f(t)\,dt \to 0, \quad \text{as} \quad x \to 0+.$$

Since

$$\int_0^x \Big(\int_0^t g\Big)^\alpha g(t) f(t)\,dt \ge f(x) \int_0^x \Big(\int_0^t g\Big)^\alpha g(t)\,dt \approx f(x)\Big(\int_0^x g\Big)^{\alpha+1}, \quad x > 0,$$

we have that

$$f(x)\Big(\int_0^x g\Big)^{\alpha+1} \to 0, \quad \text{as} \quad x \to 0+.$$

Integrating by parts, we get that

$$\begin{aligned} A_2 &= \int_{(0,\infty)} \Big(\int_0^t g\Big)^{\alpha+1} d[-f(t)] = -f(t)\Big(\int_0^t g\Big)^{\alpha+1}\Big|_0^\infty + \int_{(0,\infty)} f(t)\,d\Big(\int_0^t g\Big)^{\alpha+1} \\ &= \lim_{t\to0+} f(t)\Big(\int_0^t g\Big)^{\alpha+1} - \lim_{t\to+\infty} f(t)\Big(\int_0^t g\Big)^{\alpha+1} + (\alpha+1)\int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t) f(t)\,dt \\ &\le (\alpha+1)\int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t) f(t)\,dt = (\alpha+1) A_1. \end{aligned}$$

Thus

$$A_2 \lesssim A_1.$$

Now assume that

$$A_2 := \int_{(0,\infty)} \Big(\int_0^t g \Big)^{\alpha+1} d[-f(t)] < \infty.$$

Then

$$\int_{[x,\infty)} \Big(\int_0^t g \Big)^{\alpha+1} d[-f(t)] \to 0, \quad \text{as} \quad x \to +\infty.$$

Since

$$\begin{aligned} \int_{[x,\infty)} \Big(\int_0^t g \Big)^{\alpha+1} d[-f(t)] &\geq \Big(\int_0^x g \Big)^{\alpha+1} \int_{[x,\infty)} d[-f(t)] \\ &= \Big(\int_0^x g \Big)^{\alpha+1} [f(x) - \lim_{x\to+\infty} f(x)] = f(x) \Big(\int_0^x g \Big)^{\alpha+1}, \quad x > 0, \end{aligned}$$

we obtain that

$$f(x) \Big(\int_0^x g \Big)^{\alpha+1} \to 0, \quad \text{as} \quad x \to +\infty.$$

Thus, integrating by parts, we get that

$$\begin{aligned} A_1 &= \int_0^\infty \Big(\int_0^t g \Big)^{\alpha} g(t) f(t)\, dt \approx \int_0^\infty f(t)\, d \Big(\int_0^t g \Big)^{\alpha+1} \\ &= f(t) \Big(\int_0^t g \Big)^{\alpha+1} \Big|_0^\infty + \int_0^\infty \Big(\int_0^t g \Big)^{\alpha+1} d[-f(t)] \\ &= \lim_{t\to\infty} f(t) \Big(\int_0^t g \Big)^{\alpha+1} - \lim_{t\to 0+} f(t) \Big(\int_0^t g \Big)^{\alpha+1} + \int_0^\infty \Big(\int_0^t g \Big)^{\alpha+1} d[-f(t)] \\ &\leq \int_0^\infty \Big(\int_0^t g \Big)^{\alpha+1} d[-f(t)] = A_2. \end{aligned}$$

Hence

$$A_1 \lesssim A_2.$$

We have shown that if $\lim_{x\to+\infty} f(x) = 0$, then

$$A_1 < \infty \quad \Longleftrightarrow \quad A_2 < \infty,$$

and

$$A_1 \approx A_2.$$

Now assume that $\lim_{x\to+\infty} f(x) > 0$. Then, applying previous statement to the function $f(x) - \lim_{x\to+\infty} f(x)$, we arrive at

$$\int_0^\infty \Big(\int_0^t g \Big)^{\alpha} g(t) [f(t) - \lim_{x\to+\infty} f(x)]\, dt \approx \int_{(0,\infty)} \Big(\int_0^t g \Big)^{\alpha+1} d[-f(t)].$$

The proof is completed. □

Remark 2.2. Note that if $f \in \mathfrak{M}^{+,\downarrow}(0,\infty)$ is such that $\lim_{x\to+\infty} f(x) > 0$, then

$$\int_0^\infty \Big(\int_0^t g \Big)^{\alpha} g(t) f(t)\, dt < \infty \quad \Longrightarrow \quad \int_0^\infty g(x)\, dx < \infty.$$

Indeed: for each $x \in (0,\infty)$

$$\begin{aligned} \infty > \int_0^\infty \Big(\int_0^t g \Big)^{\alpha} g(t) f(t)\, dt &\geq \int_0^x \Big(\int_0^t g \Big)^{\alpha} g(t) f(t)\, dt \\ &\geq f(x) \int_0^x \Big(\int_0^t g \Big)^{\alpha} g(t)\, dt \approx f(x) \Big(\int_0^x g \Big)^{\alpha+1} \end{aligned}$$

holds. Thus

$$\lim_{x\to+\infty} f(x) \cdot \Big(\int_0^x g \Big)^{\alpha+1} \leq f(x) \Big(\int_0^x g \Big)^{\alpha+1} \leq \int_0^\infty \Big(\int_0^t g \Big)^{\alpha} g(t) f(t)\, dt < \infty.$$

Hence

$$\lim_{x\to+\infty} f(x)\cdot\Big(\int_0^\infty g\Big)^{\alpha+1} < \infty.$$

Therefore

$$\int_0^\infty g < \infty.$$

Corollary 2.3. *Let $\alpha > 0$. Let g be a non-negative function on $(0,\infty)$ such that $0 < \int_0^t g < \infty$, $t > 0$ and let f be a non-negative non-increasing right-continuous function on $(0,\infty)$. Then*

$$\int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t)f(t)\,dt \approx \int_{(0,\infty)} \Big(\int_0^t g\Big)^{\alpha+1} d[-f(t)] + \lim_{x\to+\infty} f(x)\cdot\Big(\int_0^\infty g\Big)^{\alpha+1}.$$

Proof. If $\lim_{x\to+\infty} f(x) = 0$, then the statement follows by Theorem 2.1. If $\lim_{x\to+\infty} f(x) > 0$, then by Remark 2.2, we know that

$$\int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t)f(t)\,dt < \infty \quad\Longrightarrow\quad \int_0^\infty g(x)\,dx < \infty.$$

Therefore, by Theorem 2.1, we get that

$$\begin{aligned}\int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t)f(t)\,dt &= \int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t)[f(t) - \lim_{x\to+\infty} f(x)]\,dt + \lim_{x\to+\infty} f(x)\cdot\int_0^\infty \Big(\int_0^t g\Big)^\alpha g(t)\,dt\\ &\approx \int_{(0,\infty)} \Big(\int_0^t g\Big)^{\alpha+1} d[-f(t)] + \lim_{x\to+\infty} f(x)\cdot\Big(\int_0^\infty g\Big)^{\alpha+1}.\end{aligned}$$

The proof is completed. □

Theorem 2.4. *Let $\alpha > 0$. Let g be a non-negative function on $(0,\infty)$ such that $0 < \int_t^\infty g < \infty$, $t > 0$ and let f be a non-negative non-decreasing left-continuous function on $(0,\infty)$. Then*

$$B_1 := \int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)[f(t) - f(0+)]\,dt < \infty \quad\Longleftrightarrow\quad B_2 := \int_{(0,\infty)} \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] < \infty.$$

Moreover, $B_1 \approx B_2$.

Proof. Assume at first that $f(0+) = 0$. Let

$$B_1 := \int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt < \infty.$$

Then

$$\int_x^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt \to 0, \quad \text{as} \quad x\to\infty.$$

Since

$$\int_x^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt \ge f(x)\int_x^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)\,dt \approx f(x)\Big(\int_x^\infty g\Big)^{\alpha+1}, \quad x > 0,$$

we have that

$$f(x)\Big(\int_x^\infty g\Big)^{\alpha+1} \to 0, \quad \text{as} \quad x\to\infty.$$

Hence, integrating by parts, we get that

$$\begin{aligned}B_2 &= \int_{(0,\infty)} \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] = f(t)\Big(\int_t^\infty g\Big)^{\alpha+1}\Big|_0^\infty - \int_{(0,\infty)} f(t)\,d\Big(\int_t^\infty g\Big)^{\alpha+1}\\ &= \lim_{t\to\infty} f(t)\Big(\int_t^\infty g\Big)^{\alpha+1} - \lim_{t\to0+} f(t)\Big(\int_t^\infty g\Big)^{\alpha+1} + (\alpha+1)\int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt\\ &\le (\alpha+1)\int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt = (\alpha+1)B_1.\end{aligned}$$

Now assume that

$$B_2 := \int_{(0,\infty)} \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] < \infty.$$

Then

$$\int_{(0,x]}\Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] \to 0, \quad \text{as} \quad x \to 0+.$$

Since

$$\begin{aligned}\int_{(0,x]}\Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] &\geq \Big(\int_x^\infty g\Big)^{\alpha+1}\int_{(0,x]} d[f(t)] \\ &= \Big(\int_x^\infty g\Big)^{\alpha+1}[f(x)-f(0+)] = f(x)\Big(\int_x^\infty g\Big)^{\alpha+1}, \quad x>0,\end{aligned}$$

we obtain that

$$f(x)\Big(\int_x^\infty g\Big)^{\alpha+1} \to 0, \quad \text{as} \quad x \to 0+.$$

Thus, integrating by parts, we get that

$$\begin{aligned}B_1 &= \int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha} g(t)f(t)\,dt \approx \int_0^\infty f(t)\,d\Big[-\Big(\int_t^\infty g\Big)^{\alpha+1}\Big] \\ &= -f(t)\Big(\int_t^\infty g\Big)^{\alpha+1}\Big|_0^\infty + \int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] \\ &= \lim_{t\to 0+} f(t)\Big(\int_t^\infty g\Big)^{\alpha+1} - \lim_{t\to\infty} f(t)\Big(\int_t^\infty g\Big)^{\alpha+1} + \int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] \\ &\leq \int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] = B_2.\end{aligned}$$

We have shown that if $f(0+)=0$, then

$$B_1 < \infty \quad \Longleftrightarrow \quad B_2 < \infty,$$

and

$$B_1 \approx B_2.$$

Now assume that $f(0+)>0$. Then, applying previous statement to the function $f(x)-f(0+)$, we arrive at

$$\int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha} g(t)[f(t)-f(0+)]\,dt \approx \int_{(0,\infty)}\Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)].$$

The proof is completed. □

Remark 2.5. Note that if f is a non-negative non-decreasing function on $(0,\infty)$ such that $f(0+)>0$, then

$$\int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha} g(t)f(t)\,dt < \infty \quad \Longrightarrow \quad \int_0^\infty g(x)\,dx < \infty.$$

Indeed: for each $x \in (0,\infty)$

$$\begin{aligned}\infty > \int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha} g(t)f(t)\,dt &\geq \int_x^\infty \Big(\int_t^\infty g\Big)^{\alpha} g(t)f(t)\,dt \\ &\geq f(x)\int_x^\infty \Big(\int_t^\infty g\Big)^{\alpha} g(t)\,dt \approx f(x)\Big(\int_x^\infty g\Big)^{\alpha+1}\end{aligned}$$

holds. Thus

$$f(0+)\Big(\int_x^\infty g\Big)^{\alpha+1} \leq f(x)\Big(\int_x^\infty g\Big)^{\alpha+1} \leq \int_0^\infty \Big(\int_t^\infty g\Big)^{\alpha} g(t)f(t)\,dt < \infty.$$

Hence

$$f(0+)\Big(\int_0^\infty g\Big)^{\alpha+1} < \infty.$$

Therefore

$$\int_0^\infty g < \infty.$$

Corollary 2.6. *Let $\alpha > 0$. Let g be a non-negative function on $(0,\infty)$ such that $0 < \int_t^\infty g < \infty$, $t > 0$ and let f be a non-negative non-decreasing left-continuous function on $(0,\infty)$. Then*

$$\int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt \approx \int_{(0,\infty)} \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] + f(0+)\Big(\int_0^\infty g\Big)^{\alpha+1}.$$

Proof. If $f(0+) = 0$, then the statement follows by Theorem 2.4. If $f(0+) > 0$, then by Remark 2.5, we know that

$$\int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt < \infty \quad \Longrightarrow \quad \int_0^\infty g(x)\,dx < \infty.$$

Therefore, by Theorem 2.4, we get that

$$\begin{aligned}\int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)f(t)\,dt &= \int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)[f(t)-f(0+)]\,dt + f(0+)\int_0^\infty \Big(\int_t^\infty g\Big)^\alpha g(t)\,dt \\ &\approx \int_{(0,\infty)} \Big(\int_t^\infty g\Big)^{\alpha+1} d[f(t)] + f(0+)\Big(\int_0^\infty g\Big)^{\alpha+1}.\end{aligned}$$

The proof is completed. □

3. The boundedness of $T_{u,b}$ and $T^*_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$

In this section we give solutions of the following two inequalities

$$\Big(\int_0^\infty \Big(\int_0^x \Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\int_0^\tau h(y)b(y)\,dy\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}} \le C\Big(\int_0^\infty h(s)^p v(s)\,ds\Big)^{\frac{1}{p}}, \quad h\in\mathfrak{M}^+(0,\infty) \tag{3.1}$$

and

$$\Big(\int_0^\infty \Big(\int_0^x \Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\int_\tau^\infty h(y)b(y)\,dy\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}} \le C\Big(\int_0^\infty h(s)^p v(s)\,ds\Big)^{\frac{1}{p}}, \quad h\in\mathfrak{M}^+(0,\infty), \tag{3.2}$$

where $1 < p \le q < \infty$ and $a, u, v, w \in \mathcal{W}(0,\infty)$. Using the duality argument, we reduce the problem to the boundedness for the dual of integral Volterra operator with a kernel satisfying Oinarovs condition and weighted Stieltjes operator.

Note that the characterization of inequalities

$$\Big(\int_0^\infty \Big(\int_x^\infty \Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\int_0^\tau h(y)b(y)\,dy\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}} \le C\Big(\int_0^\infty h(s)^p v(s)\,ds\Big)^{\frac{1}{p}}, \quad h\in\mathfrak{M}^+(0,\infty) \tag{3.3}$$

and

$$\Big(\int_0^\infty \Big(\int_x^\infty \Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\int_\tau^\infty h(y)b(y)\,dy\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}} \le C\Big(\int_0^\infty h(s)^p v(s)\,ds\Big)^{\frac{1}{p}}, \quad h\in\mathfrak{M}^+(0,\infty) \tag{3.4}$$

can be reduced to the solutions of (3.1) and (3.2).

Recall that, if F is a non-negative non-decreasing function on $(0,\infty)$, then

$$\operatorname*{ess\,sup}_{t\in(0,\infty)} F(t)G(t) = \operatorname*{ess\,sup}_{t\in(0,\infty)} F(t) \operatorname*{ess\,sup}_{\tau\in(t,\infty)} G(\tau), \tag{3.5}$$

likewise, when F is a non-negative non-increasing function on $(0,\infty)$, then

$$\operatorname*{ess\,sup}_{t\in(0,\infty)} F(t)G(t) = \operatorname*{ess\,sup}_{t\in(0,\infty)} F(t) \operatorname*{ess\,sup}_{\tau\in(0,t)} G(\tau) \tag{3.6}$$

(see, for instance, [24, p. 85]).

We need the following notations:

$$A(t) := \int_0^t a(s)ds, \qquad U(t) := \int_0^t u(s)ds, \qquad W(t) := \int_0^t w(s)ds.$$

Theorem 3.1. *Let $1 < p, q < \infty$. Assume that $u \in \mathcal{W}(0,\infty) \cap C(0,\infty)$ and $a, v, w \in \mathcal{W}(0,\infty)$. Moreover, assume that*

$$0 < \int_0^x v(t)^{1-p'}\,dt < \infty \qquad \text{for all} \quad x > 0.$$

(i) *If $p \le q$, then*

$$\begin{aligned}
&\sup_{h\ge 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} u(\tau) \int_0^\tau h(y)\,dy\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\quad \approx \sup_{t\in(0,\infty)} \left(\int_0^t v(x)^{1-p'} \left(\int_x^t \left(\sup_{s\le\tau} u(\tau)\right) a(s)\,ds\right)^{p'} dx\right)^{\frac{1}{p'}} \left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \sup_{t\in(0,\infty)} \left(\int_0^t v(x)^{1-p'}\,dx\right)^{\frac{1}{p'}} \left(\int_t^\infty \left(\int_t^y \left(\sup_{s\le\tau} u(\tau)\right) a(s)\,ds\right)^q w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_{[x,\infty)} d\left(-\sup_{t\le\tau} u(\tau)^{p'} \left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)^{\frac{1}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\sup_{t\le\tau} u(\tau)^{p'} \left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)^{\frac{1}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \lim_{t\to\infty} \left(\sup_{t\le\tau} u(\tau) \left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right);
\end{aligned}$$

(ii) *If $q < p$, then*

$$\begin{aligned}
&\sup_{h\ge 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} u(\tau) \int_0^\tau h(y)\,dy\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\quad \approx \left(\int_0^\infty \left(\int_0^t v(x)^{1-p'}\,dx\right)^{\frac{r}{q'}} v(t)^{1-p'} \left(\int_t^\infty \left(\int_t^z \left(\sup_{s\le y} u(y)\right) a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{r}{q}} dt\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_0^t v(x)^{1-p'} \left(\int_x^t \left(\sup_{s\le y} u(y)\right) a(s)\,ds\right)^{p'} dx\right)^{\frac{r}{p'}} \left(\int_z^\infty w(s)\,ds\right)^{\frac{r}{p}} w(t)\,dt\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_{[x,\infty)} d\left(-\left(\sup_{t\le\tau} u(\tau)^{p'} \left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}} A(x)^q w(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\left(\sup_{t\le\tau} u(\tau)^{p'} \left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}} w(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \lim_{t\to\infty} \left(\sup_{t\le\tau} u(\tau) \left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right).
\end{aligned}$$

Proof. Assume that $1 < p \le q < \infty$. By duality, using Fubini's Theorem, and interchanging the suprema, we get that

$$\begin{aligned}
&\sup_{h\ge 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} u(\tau) \int_0^\tau h(y)\,dy\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\qquad = \sup_{h\ge 0} \frac{1}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \sup_{g\ge 0} \frac{\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} u(\tau) \int_0^\tau h(y)\,dy\right) a(t)\,dt\right) g(x)\,dx}{\left(\int_0^\infty g(x)^{q'} w(x)^{1-q'}\,dx\right)^{\frac{1}{q'}}}
\end{aligned}$$

$$= \sup_{g\geq 0} \frac{1}{\left(\int_0^\infty g(x)^{q'} w(x)^{1-q'}\,dx\right)^{\frac{1}{q'}}} \sup_{h\geq 0} \frac{\int_0^\infty \left(\sup_{t\leq\tau} u(\tau)\int_0^\tau h(y)\,dy\right)\left(\int_t^\infty g(x)\,dx\right)a(t)\,dt}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}}.$$

Applying [23, Theorems 4.4], on using (3.5), we arrive at

$$\sup_{h\geq 0} \frac{\int_0^\infty \left(\sup_{t\leq\tau} u(\tau)\int_0^\tau h(y)\,dy\right)\left(\int_t^\infty g(x)\,dx\right)a(t)\,dt}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \approx D + E,$$

where

$$D := \left(\int_0^\infty \left(\int_t^\infty \left(\sup_{s\leq\tau} u(\tau)\right)\left(\int_s^\infty g(x)\,dx\right)a(s)\,ds\right)^{\frac{p'}{p}} \left(\sup_{t\leq\tau} u(\tau)\right)\left(\int_0^t v(s)^{1-p'}\,ds\right)\left(\int_t^\infty g(x)\,dx\right)a(t)\,dt\right)^{\frac{1}{p'}},$$

$$E := \left(\int_0^\infty \left(\int_0^t \left(\int_s^\infty g(x)\,dx\right)a(s)\,ds\right)^{\frac{p'}{p}} \left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\left(\int_t^\infty g(x)\,dx\right)a(t)\,dt\right)^{\frac{1}{p'}}.$$

Integrating by parts (applying Corollary 2.6), on using Fubini's Theorem, we arrive at

$$\begin{aligned} D &\approx \left(\int_0^\infty \left(\int_t^\infty \left(\sup_{s\leq\tau} u(\tau)\right)\left(\int_s^\infty g(x)\,dx\right)a(s)\,ds\right)^{p'} v(t)^{1-p'}\,dt\right)^{\frac{1}{p'}} \\ &= \left(\int_0^\infty \left(\int_t^\infty g(x)\int_t^x \left(\sup_{s\leq\tau} u(\tau)\right)a(s)\,ds\,dx\right)^{p'} v(t)^{1-p'}\,dt\right)^{\frac{1}{p'}}. \end{aligned}$$

Similarly, integrating by parts (applying Corollary 2.3), on using Fubini's Theorem, we get at

$$\begin{aligned} E &\approx \left(\int_0^\infty \left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right) d\left(\int_0^t \left(\int_s^\infty g(x)\,dx\right)a(s)\,ds\right)^{p'}\right)^{\frac{1}{p'}} \\ &\approx \left(\int_0^\infty \left(\int_0^t \left(\int_s^\infty g(x)\,dx\right)a(s)\,ds\right)^{p'} d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}} \\ &\quad + \left(\int_0^\infty \left(\int_s^\infty g(x)\,dx\right)a(s)\,ds\right)\lim_{t\to\infty}\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right) \\ &\approx \left(\int_0^\infty \left(\int_0^t g(x)A(x)\,dx\right)^{p'} d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}} \\ &\quad + \left(\int_0^\infty \left(\int_t^\infty g(x)\,dx\right)^{p'} A(t)^{p'}\, d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}} \\ &\quad + \left(\int_0^\infty g(x)A(x)\,dx\right)\lim_{t\to\infty}\left(\sup_{t\leq\tau} u(\tau)\left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right) := E_1 + E_2 + E_3. \end{aligned}$$

(i) Let $p \leq q$. By [35, Theorem 1.1], we obtain that

$$\begin{aligned} &\sup_{g\geq 0} \frac{D}{\left(\int_0^\infty g^{q'} w^{1-q'}\right)^{\frac{1}{q'}}} \\ &= \sup_{g\geq 0} \frac{1}{\left(\int_0^\infty g^{q'} w^{1-q'}\right)^{\frac{1}{q'}}} \left(\int_0^\infty \left(\int_t^\infty g(x)\int_t^x \left(\sup_{s\leq y} u(y)\right)a(s)\,ds\,dx\right)^{p'} v(t)^{1-p'}\,dt\right)^{\frac{1}{p'}} \\ &\approx \sup_{t\in(0,\infty)} \left(\int_0^t v(x)^{1-p'}\left(\int_x^t \left(\sup_{s\leq y} u(y)\right)a(s)\,ds\right)^{p'} dx\right)^{\frac{1}{p'}} \left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}} \\ &\quad + \sup_{t\in(0,\infty)} \left(\int_0^t v(x)^{1-p'}\,dx\right)^{\frac{1}{p'}} \left(\int_t^\infty \left(\int_t^z \left(\sup_{s\leq y} u(y)\right)a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{1}{q}}. \end{aligned} \tag{3.7}$$

By [33, Theorem 1, p. 40 and Theorem 3, p. 44], respectively, we have that

$$\begin{aligned}
&\sup_{g\geq 0}\frac{E_1}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&=\sup_{g\geq 0}\frac{\left(\int_0^\infty\left(\int_0^t g(x)A(x)\,dx\right)^{p'}d\left(-\left(\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}}}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&\approx\sup_{x\in(0,\infty)}\left(\int_{[x,\infty)}d\left(-\left(\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}}
\end{aligned}$$

and

$$\begin{aligned}
&\sup_{g\geq 0}\frac{E_2}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&=\sup_{g\geq 0}\frac{\left(\int_0^\infty\left(\int_t^\infty g(x)\,dx\right)^{p'}A(t)^{p'}\,d\left(-\left(\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}}}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&\approx\sup_{x\in(0,\infty)}\left(\int_{(0,x]}A(t)^{p'}\,d\left(-\left(\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}}\left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}}.
\end{aligned}$$

By duality, we have that

$$\begin{aligned}
&\sup_{g\geq 0}\frac{E_3}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&=\sup_{g\geq 0}\frac{\int_0^\infty g(x)A(x)\,dx}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\cdot\lim_{t\to\infty}\left(\sup_{t\leq\tau}u(\tau)\left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right)\\
&=\left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\leq\tau}u(\tau)\left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right).
\end{aligned}$$

Thus, we get that

$$\begin{aligned}
&\sup_{g\geq 0}\frac{E}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&\approx\sup_{x\in(0,\infty)}\left(\int_{[x,\infty)}d\left(-\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\\
&\quad+\sup_{x\in(0,\infty)}\left(\int_{(0,x]}A(t)^{p'}\,d\left(-\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}}\\
&\quad+\left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\leq\tau}u(\tau)\left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right).
\end{aligned}\tag{3.8}$$

Combining (3.7) and (3.8), we arrive at

$$\begin{aligned}
&\sup_{h\geq 0}\frac{\left(\int_0^\infty\left(\int_0^x\left(\sup_{t\leq\tau}u(\tau)\int_0^\tau h(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}}\\
&\quad\approx \sup_{t\in(0,\infty)}\left(\int_0^t v(x)^{1-p'}\left(\int_x^t\left(\sup_{s\leq\tau}u(\tau)\right)a(s)\,ds\right)^{p'}dx\right)^{\frac{1}{p'}}\left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\sup_{t\in(0,\infty)}\left(\int_0^t v(x)^{1-p'}\,dx\right)^{\frac{1}{p'}}\left(\int_t^\infty\left(\int_t^y\left(\sup_{s\leq\tau}u(\tau)\right)a(s)\,ds\right)^q w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\sup_{x\in(0,\infty)}\left(\int_{[x,\infty)}d\left(-\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\sup_{x\in(0,\infty)}\left(\int_{(0,x]}A(t)^{p'}\,d\left(-\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\leq\tau}u(\tau)\left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right).
\end{aligned}$$

(ii) Let now $q<p$. By [35, Theorem 1.2], we obtain that

$$\begin{aligned}
&\sup_{g\geq 0}\frac{D}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&\quad=\sup_{g\geq 0}\frac{\left(\int_0^\infty\left(\int_t^\infty g(x)\int_t^x\left(\sup_{s\leq y}u(y)\right)a(s)\,ds\,dx\right)^{p'}v(t)^{1-p'}\,dt\right)^{\frac{1}{p'}}}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&\quad\approx\left(\int_0^\infty\left(\int_0^t v(x)^{1-p'}\right)^{\frac{r}{q'}}v(t)^{1-p'}\left(\int_t^\infty\left(\int_t^z\left(\sup_{s\leq y}u(y)\right)a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{r}{q}}dt\right)^{\frac{1}{r}}\\
&\qquad+\left(\int_0^\infty\left(\int_0^t v(x)^{1-p'}\left(\int_x^t\left(\sup_{s\leq y}u(y)\right)a(s)\,ds\right)^{p'}dx\right)^{\frac{r}{p'}}\left(\int_t^\infty w(s)\,ds\right)^{\frac{r}{p}}w(t)\,dt\right)^{\frac{1}{r}}.
\end{aligned}\tag{3.9}$$

By [33, Theorem 2, p. 48], we have that

$$\begin{aligned}
&\sup_{g\geq 0}\frac{E_1}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&\quad=\sup_{g\geq 0}\frac{\left(\int_0^\infty\left(\int_0^t g(x)A(x)\,dx\right)^{p'}d\left(-\left(\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}}}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}\\
&\quad\approx\left(\int_0^\infty\left(\int_{[x,\infty)}d\left(-\left(\sup_{t\leq\tau}u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}}A(x)^q w(x)\,dx\right)^{\frac{1}{r}}
\end{aligned}$$

and

$$\sup_{g\geq 0}\frac{E_2}{\left(\int_0^\infty g^{q'}w^{1-q'}\right)^{\frac{1}{q'}}}$$

$$= \sup_{g\geq 0} \frac{\left(\int_0^\infty \left(\int_t^\infty g(x)\,dx\right)^{p'} A(t)^{p'}\, d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{1}{p'}}}{\left(\int_0^\infty g^{q'} w^{1-q'}\right)^{\frac{1}{q'}}}$$

$$\approx \left(\int_0^\infty \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}} w(x)\,dx\right)^{\frac{1}{r}}.$$

Consequently, we arrive at

$$\begin{aligned}
&\sup_{g\geq 0} \frac{E}{\left(\int_0^\infty g^{q'} w^{1-q'}\right)^{\frac{1}{q'}}} \\
&\approx \left(\int_0^\infty \left(\int_{[x,\infty)} d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}} A(x)^q w(x)\,dx\right)^{\frac{1}{r}} \\
&\quad + \left(\int_0^\infty \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}} w(x)\,dx\right)^{\frac{1}{r}} \\
&\quad + \left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \lim_{t\to\infty}\left(\sup_{t\leq\tau} u(\tau)\left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right).
\end{aligned} \tag{3.10}$$

Combining (3.9) and (3.10), we arrive at

$$\begin{aligned}
&\sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} u(\tau)\int_0^\tau h(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\approx \left(\int_0^\infty \left(\int_0^t v(x)^{1-p'}\right)^{\frac{r}{q'}} v(t)^{1-p'} \left(\int_t^\infty \left(\int_t^z \left(\sup_{s\leq y} u(y)\right)a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{r}{q}} dt\right)^{\frac{1}{r}} \\
&\quad + \left(\int_0^\infty \left(\int_0^t v(x)^{1-p'}\left(\int_x^t \left(\sup_{s\leq y} u(y)\right)a(s)\,ds\right)^{p'} dx\right)^{\frac{r}{p'}} \left(\int_t^\infty w(s)\,ds\right)^{\frac{r}{p}} w(t)\,dt\right)^{\frac{1}{r}} \\
&\quad + \left(\int_0^\infty \left(\int_{[x,\infty)} d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}} A(x)^q w(x)\,dx\right)^{\frac{1}{r}} \\
&\quad + \left(\int_0^\infty \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\left(\sup_{t\leq\tau} u(\tau)^{p'}\left(\int_0^\tau v(s)^{1-p'}\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}} w(x)\,dx\right)^{\frac{1}{r}} \\
&\quad + \left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \lim_{t\to\infty}\left(\sup_{t\leq\tau} u(\tau)\left(\int_0^\tau v(s)^{1-p'}\,ds\right)^{\frac{1}{p'}}\right).
\end{aligned}$$

The proof is completed. □

Theorem 3.2. *Let $1 < p, q < \infty$ and $b \in \mathcal{W}(0,\infty)$ be such that $b(t) > 0$ for a.e. $t \in (0,\infty)$. Assume that $u \in \mathcal{W}(0,\infty)\cap C(0,\infty)$ and $a, v, w \in \mathcal{W}(0,\infty)$. Moreover, assume that*

$$0 < \int_0^x v(t)^{1-p'}\,dt < \infty \qquad \text{for all} \quad x > 0.$$

(i) *If $p \leq q$, then*

$$\begin{aligned}
&\sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)}\int_0^\tau h(y)b(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\qquad \approx \sup_{t\in(0,\infty)} \left(\int_0^t b(x)^{p'} v(x)^{1-p'} \left(\int_x^t \left(\sup_{s\leq\tau} \frac{u(\tau)}{B(\tau)}\right)a(s)\,ds\right)^{p'} dx\right)^{\frac{1}{p'}} \left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}}
\end{aligned}$$

$$+\sup_{t\in(0,\infty)}\Big(\int_0^t b(x)^{p'}v(x)^{1-p'}\,dx\Big)^{\frac{1}{p'}}\Big(\int_t^\infty\Big(\int_t^y\Big(\sup_{s\le\tau}\frac{u(\tau)}{B(\tau)}\Big)a(s)\,ds\Big)^q w(y)\,dy\Big)^{\frac{1}{q}}$$

$$+\sup_{x\in(0,\infty)}\Big(\int_{[x,\infty)} d\Big(-\sup_{t\le\tau}\Big(\frac{u(\tau)}{B(\tau)}\Big)^{p'}\Big(\int_0^\tau b(s)^{p'}v(s)^{1-p'}\,ds\Big)\Big)\Big)^{\frac{1}{p'}}\Big(\int_0^x A(y)^q w(y)\,dy\Big)^{\frac{1}{q}}$$

$$+\sup_{x\in(0,\infty)}\Big(\int_{(0,x]} A(t)^{p'}\,d\Big(-\sup_{t\le\tau}\Big(\frac{u(\tau)}{B(\tau)}\Big)^{p'}\Big(\int_0^\tau b(s)^{p'}v(s)^{1-p'}\,ds\Big)\Big)\Big)^{\frac{1}{p'}}\Big(\int_x^\infty w(y)\,dy\Big)^{\frac{1}{q}}$$

$$+\Big(\int_0^\infty A(y)^q w(y)\,dy\Big)^{\frac{1}{q}}\lim_{t\to\infty}\Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\Big(\int_0^\tau b(s)^{p'}v(s)^{1-p'}\,ds\Big)^{\frac{1}{p'}}\Big);$$

(ii) *If $q<p$, then*

$$\sup_{h\ge0}\frac{\Big(\int_0^\infty\Big(\int_0^x\Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\int_0^\tau h(y)b(y)\,dy\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}}}{\Big(\int_0^\infty h(s)^p v(s)\,ds\Big)^{\frac{1}{p}}}$$

$$\approx\Big(\int_0^\infty\Big(\int_0^t b(x)^{p'}v(x)^{1-p'}\,dx\Big)^{\frac{r}{q'}}b(t)^{p'}v(t)^{1-p'}\Big(\int_t^\infty\Big(\int_t^z\Big(\sup_{s\le y}\frac{u(y)}{B(y)}\Big)a(s)\,ds\Big)^q w(z)\,dz\Big)^{\frac{r}{q}}dt\Big)^{\frac{1}{r}}$$

$$+\Big(\int_0^\infty\Big(\int_0^t b(x)^{p'}v(x)^{1-p'}\Big(\int_x^t\Big(\sup_{s\le y}\frac{u(y)}{B(y)}\Big)a(s)\,ds\Big)^{p'}dx\Big)^{\frac{r}{p'}}\Big(\int_z^\infty w(s)\,ds\Big)^{\frac{r}{p}}w(t)\,dt\Big)^{\frac{1}{r}}$$

$$+\Big(\int_0^\infty\Big(\int_{[x,\infty)} d\Big(-\Big(\sup_{t\le\tau}\Big(\frac{u(\tau)}{B(\tau)}\Big)^{p'}\Big(\int_0^\tau b(s)^{p'}v(s)^{1-p'}\,ds\Big)\Big)\Big)\Big)^{\frac{r}{p'}}\Big(\int_0^x A(y)^q w(y)\,dy\Big)^{\frac{r}{p}}A(x)^q w(x)\,dx\Big)^{\frac{1}{r}}$$

$$+\Big(\int_0^\infty\Big(\int_{(0,x]} A(t)^{p'}\,d\Big(-\Big(\sup_{t\le\tau}\Big(\frac{u(\tau)}{B(\tau)}\Big)^{p'}\Big(\int_0^\tau b(s)^{p'}v(s)^{1-p'}\,ds\Big)\Big)\Big)\Big)^{\frac{r}{p'}}\Big(\int_x^\infty w(y)\,dy\Big)^{\frac{r}{p}}w(x)\,dx\Big)^{\frac{1}{r}}$$

$$+\Big(\int_0^\infty A(y)^q w(y)\,dy\Big)^{\frac{1}{q}}\lim_{t\to\infty}\Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\Big(\int_0^\tau b(s)^{p'}v(s)^{1-p'}\,ds\Big)^{\frac{1}{p'}}\Big).$$

Proof. The statement follows by Theorem 3.1 at once if we note that

$$\sup_{h\ge0}\frac{\Big(\int_0^\infty\Big(\int_0^x\Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\int_0^\tau h(y)b(y)\,dy\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}}}{\Big(\int_0^\infty h(s)^p v(s)\,ds\Big)^{\frac{1}{p}}}$$

$$=\sup_{h\ge0}\frac{\Big(\int_0^\infty\Big(\int_0^x\Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\int_0^\tau h(y)\,dy\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}}}{\Big(\int_0^\infty h(s)^p b(s)^{-p}v(s)\,ds\Big)^{\frac{1}{p}}}.$$

□

Theorem 3.3. *Let $1<p,q<\infty$ and $b\in\mathcal{W}(0,\infty)$ be such that $b(t)>0$ for a.e. $t\in(0,\infty)$. Assume that $u\in\mathcal{W}(0,\infty)\cap C(0,\infty)$ and $a,v,w\in\mathcal{W}(0,\infty)$. Moreover, assume that*

$$0<\int_x^\infty v(t)^{1-p'}\,dt<\infty\qquad\text{for all}\quad x>0.$$

Denote by

$$\psi(x):=\Big(\int_x^\infty b(t)^{p'}v^{1-p'}(t)\,dt\Big)^{-\frac{p'}{p'+1}}b(x)^{p'}v^{1-p'}(x)$$

and

$$\Psi(x) := \left(\int_x^\infty b(t)^{p'} v^{1-p'}(t)\,dt\right)^{\frac{1}{p'+1}}.$$

(i) *If $p \le q$, then*

$$\begin{aligned}
&\sup_{h\ge 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} \frac{u(\tau)}{B(\tau)} \int_\tau^\infty h(y)b(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\quad\approx \sup_{t\in(0,\infty)} \left(\int_0^t \Psi(x)^{-p'}\psi(x)\left(\int_x^t \left(\sup_{s\le\tau} \frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\right)a(s)\,ds\right)^{p'} dx\right)^{\frac{1}{p'}} \left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \sup_{t\in(0,\infty)} \left(\int_0^t \Psi(x)^{-p'}\psi(x)\,dx\right)^{\frac{1}{p'}} \left(\int_t^\infty \left(\int_t^y \left(\sup_{s\le\tau} \frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\right)a(s)\,ds\right)^q w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_{[x,\infty)} d\left(-\sup_{t\le\tau}\left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'}\left(\int_0^\tau \Psi(s)^{-p'}\psi(s)\,ds\right)\right)\right)^{\frac{1}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\sup_{t\le\tau}\left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'}\left(\int_0^\tau \Psi(s)^{-p'}\psi(s)\,ds\right)\right)\right)^{\frac{1}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}} \\
&\qquad + \left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \lim_{t\to\infty}\left(\sup_{t\le\tau} \frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\left(\int_0^\tau \Psi(s)^{-p'}\psi(s)\,ds\right)^{\frac{1}{p'}}\right) \\
&\qquad + \left(\int_0^\infty \psi(s)\,ds\right)^{-\frac{1}{p}} \left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} \frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}};
\end{aligned}$$

(ii) *If $q < p$, then*

$$\begin{aligned}
&\sup_{h\ge 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} \frac{u(\tau)}{B(\tau)} \int_\tau^\infty h(y)b(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\quad\approx \left(\int_0^\infty \left(\int_0^t \Psi(x)^{-p'}\psi(x)\,dx\right)^{\frac{r}{q'}} \Psi(t)^{-p'}\psi(t)\left(\int_t^\infty \left(\int_t^z \left(\sup_{s\le y} \frac{u(y)}{B(y)}\Psi(y)^2\right)a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{r}{q}} dt\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_0^t \Psi(x)^{-p'}\psi(x)\left(\int_x^t \left(\sup_{s\le y} \frac{u(y)}{B(y)}\Psi(y)^2\right)a(s)\,ds\right)^{p'} dx\right)^{\frac{r}{p'}} \left(\int_z^\infty w(s)\,ds\right)^{\frac{r}{p}} w(t)\,dt\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_{[x,\infty)} d\left(-\left(\sup_{t\le\tau}\left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'}\left(\int_0^\tau \Psi(s)^{-p'}\psi(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}} A(x)^q w(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\left(\sup_{t\le\tau}\left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'}\left(\int_0^\tau \Psi(s)^{-p'}\psi(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}} w(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \lim_{t\to\infty}\left(\sup_{t\le\tau} \frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\left(\int_0^\tau \Psi(s)^{-p'}\psi(s)\,ds\right)^{\frac{1}{p'}}\right) \\
&\qquad + \left(\int_0^\infty \psi(s)\,ds\right)^{-\frac{1}{p}} \left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} \frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}.
\end{aligned}$$

Proof. By [20, Corollary 3.5], we have that

$$\sup_{h\ge 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\le\tau} \frac{u(\tau)}{B(\tau)} \int_\tau^\infty h(y)b(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$

$$= \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)} \int_\tau^\infty h(y)\,dy\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p b(s)^{-p} v(s)\,ds\right)^{\frac{1}{p}}}$$

$$\approx \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)} \Psi(\tau)^2 \int_0^\tau h(y)\,dy\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p \Psi(s)^p \psi(s)^{1-p}\,ds\right)^{\frac{1}{p}}}$$

$$+ \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)} \Psi(\tau)^2\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty \psi(s)\,ds\right)^{\frac{1}{p}}}.$$

(i) Let $p \leq q$. By Theorem 3.1, (i), we get that

$$\sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)} \int_\tau^\infty h(y)b(y)\,dy\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$

$$\approx \sup_{t\in(0,\infty)} \left(\int_0^t \Psi(x)^{-p'} \psi(x) \left(\int_x^t \left(\sup_{s\leq\tau} \frac{u(\tau)}{B(\tau)} \Psi(\tau)^2\right) a(s)\,ds\right)^{p'} dx\right)^{\frac{1}{p'}} \left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}}$$

$$+ \sup_{t\in(0,\infty)} \left(\int_0^t \Psi(x)^{-p'} \psi(x)\,dx\right)^{\frac{1}{p'}} \left(\int_t^\infty \left(\int_t^y \left(\sup_{s\leq\tau} \frac{u(\tau)}{B(\tau)} \Psi(\tau)^2\right) a(s)\,ds\right)^q w(y)\,dy\right)^{\frac{1}{q}}$$

$$+ \sup_{x\in(0,\infty)} \left(\int_{[x,\infty)} d\left(-\sup_{t\leq\tau} \left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'} \left(\int_0^\tau \Psi(s)^{-p'} \psi(s)\,ds\right)\right)\right)^{\frac{1}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}}$$

$$+ \sup_{x\in(0,\infty)} \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\sup_{t\leq\tau} \left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'} \left(\int_0^\tau \Psi(s)^{-p'} \psi(s)\,ds\right)\right)\right)^{\frac{1}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}}$$

$$+ \left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}} \lim_{t\to\infty} \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)} \Psi(\tau)^2 \left(\int_0^\tau \Psi(s)^{-p'} \psi(s)\,ds\right)^{\frac{1}{p'}}\right)$$

$$+ \left(\int_0^\infty \psi(s)\,ds\right)^{-\frac{1}{p}} \left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)} \Psi(\tau)^2\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}};$$

(ii) Let $q < p$. By Theorem 3.1, (ii), we obtain that

$$\sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\sup_{t\leq\tau} \frac{u(\tau)}{B(\tau)} \int_\tau^\infty h(y)b(y)\,dy\right) a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$

$$\approx \left(\int_0^\infty \left(\int_0^t \Psi(x)^{-p'} \psi(x)\,dx\right)^{\frac{r}{q'}} \Psi(t)^{-p'} \psi(t) \left(\int_t^\infty \left(\int_t^z \left(\sup_{s\leq y} \frac{u(y)}{B(y)} \Psi(y)^2\right) a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{r}{q}} dt\right)^{\frac{1}{r}}$$

$$+ \left(\int_0^\infty \left(\int_0^t \Psi(x)^{-p'} \psi(x) \left(\int_x^t \left(\sup_{s\leq y} \frac{u(y)}{B(y)} \Psi(y)^2\right) a(s)\,ds\right)^{p'} dx\right)^{\frac{r}{p'}} \left(\int_z^\infty w(s)\,ds\right)^{\frac{r}{p}} w(t)\,dt\right)^{\frac{1}{r}}$$

$$+ \left(\int_0^\infty \left(\int_{[x,\infty)} d\left(-\left(\sup_{t\leq\tau} \left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'} \left(\int_0^\tau \Psi(s)^{-p'} \psi(s)\,ds\right)\right)\right)^{\frac{r}{p'}} \left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}} A(x)^q w(x)\,dx\right)^{\frac{1}{r}}$$

$$+ \left(\int_0^\infty \left(\int_{(0,x]} A(t)^{p'}\, d\left(-\left(\sup_{t\leq\tau} \left(\frac{u(\tau)}{B(\tau)}\right)^{p'} \Psi(\tau)^{2p'} \left(\int_0^\tau \Psi(s)^{-p'} \psi(s)\,ds\right)\right)\right)^{\frac{r}{p'}} \left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}} w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\Big(\int_0^\infty A(y)^q w(y)\,dy\Big)^{\frac{1}{q}} \lim_{t\to\infty}\Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\Big(\int_0^\tau \Psi(s)^{-p'}\psi(s)\,ds\Big)^{\frac{1}{p'}}\Big)$$
$$+\Big(\int_0^\infty \psi(s)\,ds\Big)^{-\frac{1}{p}}\Big(\int_0^\infty\Big(\int_0^x\Big(\sup_{t\le\tau}\frac{u(\tau)}{B(\tau)}\Psi(\tau)^2\Big)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}}.$$

The proof is completed. □

4. The boundedness of R_u from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions

In this section we characterize the boundedness of R_u from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions.

Theorem 4.1. *Let $1<p,q<\infty$. Assume that $u\in\mathcal{W}(0,\infty)\cap C(0,\infty)$ and $a,v,w\in\mathcal{W}(0,\infty)$.*
(i) *If $p\le q$, then*

$$\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\Big(\int_0^\infty\Big(\int_0^x(R_uf)(t)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}}}{\Big(\int_0^\infty f(s)^p v(s)\,ds\Big)^{\frac{1}{p}}}$$
$$\approx \sup_{t\in(0,\infty)}\Big(\int_0^t V(x)^{p'}v(x)\Big(\int_x^t\Big(\sup_{s\le\tau}u(\tau)V(\tau)^{-2}\Big)a(s)\,ds\Big)^{p'}dx\Big)^{\frac{1}{p'}}\Big(\int_t^\infty w(y)\,dy\Big)^{\frac{1}{q}}$$
$$+\sup_{t\in(0,\infty)}\Big(\int_0^t V(x)^{p'}v(x)\,dx\Big)^{\frac{1}{p'}}\Big(\int_t^\infty\Big(\int_t^y\Big(\sup_{s\le\tau}u(\tau)V(\tau)^{-2}\Big)a(s)\,ds\Big)^q w(y)\,dy\Big)^{\frac{1}{q}}$$
$$+\sup_{x\in(0,\infty)}\Big(\int_{[x,\infty)}d\Big(-\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)\Big)\Big)^{\frac{1}{p'}}\Big(\int_0^x A(y)^q w(y)\,dy\Big)^{\frac{1}{q}}$$
$$+\sup_{x\in(0,\infty)}\Big(\int_{(0,x]}A(t)^{p'}\,d\Big(-\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)\Big)\Big)^{\frac{1}{p'}}\Big(\int_x^\infty w(y)\,dy\Big)^{\frac{1}{q}}$$
$$+\Big(\int_0^\infty A(y)^q w(y)\,dy\Big)^{\frac{1}{q}}\lim_{t\to\infty}\Big(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)^{\frac{1}{p'}}\Big);$$

(ii) *If $q<p$, then*

$$\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\Big(\int_0^\infty\Big(\int_0^x(R_uf)(t)a(t)\,dt\Big)^q w(x)\,dx\Big)^{\frac{1}{q}}}{\Big(\int_0^\infty f(s)^p v(s)\,ds\Big)^{\frac{1}{p}}}$$
$$\approx\Big(\int_0^\infty\Big(\int_0^t V(x)^{p'}v(x)\,dx\Big)^{\frac{r}{q'}}V(t)^{p'}v(t)\Big(\int_t^\infty\Big(\int_t^z\Big(\sup_{s\le y}u(y)V(y)^{-2}\Big)a(s)\,ds\Big)^q w(z)\,dz\Big)^{\frac{r}{q}}dt\Big)^{\frac{1}{r}}$$
$$+\Big(\int_0^\infty\Big(\int_0^t V(x)^{p'}v(x)\Big(\int_x^t\Big(\sup_{s\le y}u(y)V(y)^{-2}\Big)a(s)\,ds\Big)^{p'}dx\Big)^{\frac{r}{p'}}\Big(\int_z^\infty w(s)\,ds\Big)^{\frac{r}{p}}w(t)\,dt\Big)^{\frac{1}{r}}$$
$$+\Big(\int_0^\infty\Big(\int_{[x,\infty)}d\Big(-\Big(\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)\Big)\Big)\Big)^{\frac{r}{p'}}\Big(\int_0^x A(y)^q w(y)\,dy\Big)^{\frac{r}{p}}x^q w(x)\,dx\Big)^{\frac{1}{r}}$$
$$+\Big(\int_0^\infty\Big(\int_{(0,x]}A(t)^{p'}\,d\Big(-\Big(\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)\Big)\Big)\Big)^{\frac{r}{p'}}\Big(\int_x^\infty w(y)\,dy\Big)^{\frac{r}{p}}w(x)\,dx\Big)^{\frac{1}{r}}$$
$$+\Big(\int_0^\infty A(y)^q w(y)\,dy\Big)^{\frac{1}{q}}\lim_{t\to\infty}\Big(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)^{\frac{1}{p'}}\Big).$$

Proof. By [26, Theorem 3.2] (cf. [20, Theorem 2.3]), we get that

$$\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x\left(\sup_{t\le\tau}u(\tau)f(\tau)\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$
$$\approx\sup_{h\ge0}\frac{\left(\int_0^\infty\left(\int_0^x\left(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\int_0^\tau h(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^{-p}v(s)^{1-p}\,ds\right)^{\frac{1}{p}}}.$$

By Theorem 3.1, we have that
(i) if $p\le q$, then

$$\sup_{h\ge0}\frac{\left(\int_0^\infty\left(\int_0^x\left(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\int_0^\tau h(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^{-p}v(s)^{1-p}\,ds\right)^{\frac{1}{p}}}$$
$$\approx\sup_{t\in(0,\infty)}\left(\int_0^t V(x)^{p'}v(x)\left(\int_x^t\left(\sup_{s\le\tau}u(\tau)V(\tau)^{-2}\right)a(s)\,ds\right)^{p'}dx\right)^{\frac{1}{p'}}\left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}}$$
$$+\sup_{t\in(0,\infty)}\left(\int_0^t V(x)^{p'}v(x)\,dx\right)^{\frac{1}{p'}}\left(\int_t^\infty\left(\int_t^y\left(\sup_{s\le\tau}u(\tau)V(\tau)^{-2}\right)a(s)\,ds\right)^q w(y)\,dy\right)^{\frac{1}{q}}$$
$$+\sup_{x\in(0,\infty)}\left(\int_{[x,\infty)}d\left(-\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}}$$
$$+\sup_{x\in(0,\infty)}\left(\int_{(0,x]}A(t)^{p'}\,d\left(-\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}}$$
$$+\left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)^{\frac{1}{p'}}\right).$$

(ii) if $q<p$, then

$$\sup_{h\ge0}\frac{\left(\int_0^\infty\left(\int_0^x\left(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\int_0^\tau h(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^{-p}v(s)^{1-p}\,ds\right)^{\frac{1}{p}}}$$
$$\approx\left(\int_0^\infty\left(\int_0^t V(x)^{p'}v(x)\,dx\right)^{\frac{r}{q'}}V(t)^{p'}v(t)\left(\int_t^\infty\left(\int_t^z\left(\sup_{s\le y}u(y)V(y)^{-2}\right)a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{r}{q}}dt\right)^{\frac{1}{r}}$$
$$+\left(\int_0^\infty\left(\int_0^t V(x)^{p'}v(x)\left(\int_x^t\left(\sup_{s\le y}u(y)V(y)^{-2}\right)a(s)\,ds\right)^{p'}dx\right)^{\frac{r}{p'}}\left(\int_z^\infty w(s)\,ds\right)^{\frac{r}{p}}w(t)\,dt\right)^{\frac{1}{r}}$$
$$+\left(\int_0^\infty\left(\int_{[x,\infty)}d\left(-\left(\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}}A(x)^q w(x)\,dx\right)^{\frac{1}{r}}$$
$$+\left(\int_0^\infty\left(\int_{(0,x]}A(t)^{p'}\,d\left(-\left(\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}}\left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}}w(x)\,dx\right)^{\frac{1}{r}}$$
$$+\left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)^{\frac{1}{p'}}\right).$$

□

5. The boundedness of $P_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions

In this section we characterize the boundedness of weighted Hardy operator $P_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions.

Theorem 5.1. *Let $1 < p, q < \infty$ and $b \in \mathcal{W}(0,\infty)$ be such that $b(t) > 0$ for a.e. $t \in (0,\infty)$. Assume that $u \in \mathcal{W}(0,\infty) \cap C(0,\infty)$ and $a, v, w \in \mathcal{W}(0,\infty)$.*

(i) *If $p \le q$, then*

$$\begin{aligned}
&\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)} \frac{\left(\int_0^\infty \left(\int_0^x (P_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\quad\approx \sup_{x\in(0,\infty)} \left(\int_0^x \left(\int_0^t a(y)u(y)\,dy\right)^q w(t)\,dt\right)^{\frac{1}{q}} \left(\int_x^\infty V(s)^{-p'} v(s)\,ds\right)^{\frac{1}{p'}} \\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_x^\infty w(t)\,dt\right)^{\frac{1}{q}} \left(\int_0^x \left(\int_0^s a(y)u(y)\,dy\right)^{p'} V(s)^{-p'} v(s)\,ds\right)^{\frac{1}{p'}} \\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_x^\infty \left(\int_x^t \frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^q w(t)\,dt\right)^{\frac{1}{q}} \left(\int_0^x \left(\frac{B(s)}{V(s)}\right)^{p'} v(s)\,ds\right)^{\frac{1}{p'}} \\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_x^\infty w(t)\,dt\right)^{\frac{1}{q}} \left(\int_0^x \left(\int_s^x \frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^{p'} \left(\frac{B(s)}{V(s)}\right)^{p'} v(s)\,ds\right)^{\frac{1}{p'}} \\
&\qquad + \left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}} \left(\int_0^\infty \left(\int_0^x a(t)u(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}};
\end{aligned}$$

(ii) *If $q < p$, then*

$$\begin{aligned}
&\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)} \frac{\left(\int_0^\infty \left(\int_0^x (P_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}} \\
&\quad\approx \left(\int_0^\infty \left(\int_0^x \left(\int_0^t a(y)u(y)\,dy\right)^q w(t)\,dt\right)^{\frac{r}{p}} \left(\int_x^\infty V(z)^{-p'} v(z)\,dz\right)^{\frac{r}{p'}} \left(\int_0^x a(y)u(y)\,dy\right)^q w(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_x^\infty w(t)\,dt\right)^{\frac{r}{p}} \left(\int_0^x \left(\int_0^z a(y)u(y)\,dy\right)^{p'} V(z)^{-p'} v(z)\,dz\right)^{\frac{r}{p'}} w(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_x^\infty w(t)\,dt\right)^{\frac{r}{p}} \left(\int_0^x \left(\int_z^x \frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^{p'} \left(\frac{B(z)}{V(z)}\right)^{p'} v(z)\,dz\right)^{\frac{r}{p'}} w(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty \left(\int_x^\infty w(t)\left(\int_x^t \frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^q dt\right)^{\frac{r}{q}} \left(\int_0^x \left(\frac{B(s)}{V(s)}\right)^{p'} v(s)\,ds\right)^{\frac{r}{q'}} \left(\frac{B(x)}{V(x)}\right)^{p'} v(x)\,dx\right)^{\frac{1}{r}} \\
&\qquad + \left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}} \left(\int_0^\infty \left(\int_0^x a(y)u(y)\,dy\right)^q w(x)\,dx\right)^{\frac{1}{q}}.
\end{aligned}$$

Proof. By [26, Theorem 3.1], using Fubini's Theorem, we get that

$$\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)} \frac{\left(\int_0^\infty \left(\int_0^x (P_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$

$$
\begin{aligned}
&= \sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)} \frac{\left(\int_0^\infty \left(\int_0^x \left(\frac{u(t)}{B(t)}\int_0^t f(\tau)b(\tau)\,d\tau\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}\\
&\approx \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\frac{u(t)}{B(t)}\int_0^t \left(\int_\tau^\infty h(y)\,dy\right)b(\tau)\,d\tau\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^p v(s)^{1-p}\,ds\right)^{\frac{1}{p}}} + \frac{\left(\int_0^\infty \left(\int_0^x a(t)u(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty v(s)\,ds\right)^{\frac{1}{p}}}\\
&\approx \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\int_t^\infty h(y)\,dy\right)a(t)u(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^p v(s)^{1-p}\,ds\right)^{\frac{1}{p}}}\\
&\quad + \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x \left(\frac{u(t)}{B(t)}\int_0^t h(y)B(y)\,dy\right)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^p v(s)^{1-p}\,ds\right)^{\frac{1}{p}}} + \frac{\left(\int_0^\infty \left(\int_0^x a(t)u(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty v(s)\,ds\right)^{\frac{1}{p}}}\\
&\approx \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_x^\infty h(y)\,dy\right)^q \left(\int_0^x a(t)u(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^p v(s)^{1-p}\,ds\right)^{\frac{1}{p}}} + \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x h(y)\left(\int_0^y a(t)u(t)\,dt\right)dy\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p V(s)^p v(s)^{1-p}\,ds\right)^{\frac{1}{p}}}\\
&\quad + \sup_{h\geq 0} \frac{\left(\int_0^\infty \left(\int_0^x h(y)\left(\int_y^x \frac{a(t)}{B(t)}u(t)\,dt\right)dy\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty h(s)^p B(s)^{-p} V(s)^p v(s)^{1-p}\,ds\right)^{\frac{1}{p}}} + \frac{\left(\int_0^\infty \left(\int_0^x a(t)u(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty v(s)\,ds\right)^{\frac{1}{p}}}.
\end{aligned}
$$

(i) Let $p \leq q$. Using the characterizations of weighted Hardy-type inequalities (see, for instance, [37, Section 1]), by [35, Theorem 1.1], we obtain that

$$
\begin{aligned}
&\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)} \frac{\left(\int_0^\infty \left(\int_0^x (P_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}\\
&\quad\approx \sup_{x\in(0,\infty)} \left(\int_0^x \left(\int_0^t a(y)u(y)\,dy\right)^q w(t)\,dt\right)^{\frac{1}{q}} \left(\int_x^\infty V(s)^{-p'} v(s)\,ds\right)^{\frac{1}{p'}}\\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_x^\infty w(t)\,dt\right)^{\frac{1}{q}} \left(\int_0^x \left(\int_0^s a(y)u(y)\,dy\right)^{p'} V(s)^{-p'} v(s)\,ds\right)^{\frac{1}{p'}}\\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_x^\infty \left(\int_x^t \frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^q w(t)\,dt\right)^{\frac{1}{q}} \left(\int_0^x \left(\frac{B(s)}{V(s)}\right)^{p'} v(s)\,ds\right)^{\frac{1}{p'}}\\
&\qquad + \sup_{x\in(0,\infty)} \left(\int_x^\infty w(t)\,dt\right)^{\frac{1}{q}} \left(\int_0^x \left(\int_s^x \frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^{p'} \left(\frac{B(s)}{V(s)}\right)^{p'} v(s)\,ds\right)^{\frac{1}{p'}}\\
&\qquad + \left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}} \left(\int_0^\infty \left(\int_0^x a(t)u(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}.
\end{aligned}
$$

(ii) Let now $q<p$. Using the characterizations of weighted Hardy-type inequalities (see, for instance, [37, Section 1]), by [35, Theorem 1.2], we obtain that

$$\begin{aligned}
&\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x(P_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}\\
&\quad\approx\left(\int_0^\infty\left(\int_0^x\left(\int_0^t a(y)u(y)\,dy\right)^q w(t)\,dt\right)^{\frac{r}{p}}\left(\int_x^\infty V(z)^{-p'}v(z)\,dz\right)^{\frac{r}{p'}}\left(\int_0^x a(y)u(y)\,dy\right)^q w(x)\,dx\right)^{\frac{1}{r}}\\
&\qquad+\left(\int_0^\infty\left(\int_x^\infty w(t)\,dt\right)^{\frac{r}{p}}\left(\int_0^x\left(\int_0^z a(y)u(y)\,dy\right)^{p'}V(z)^{-p'}v(z)\,dz\right)^{\frac{r}{p'}}w(x)\,dx\right)^{\frac{1}{r}}\\
&\qquad+\left(\int_0^\infty\left(\int_x^\infty w(t)\,dt\right)^{\frac{r}{p}}\left(\int_0^x\left(\int_z^x\frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^{p'}\left(\frac{B(z)}{V(z)}\right)^{p'}v(z)\,dz\right)^{\frac{r}{p'}}w(x)\,dx\right)^{\frac{1}{r}}\\
&\qquad+\left(\int_0^\infty\left(\int_x^\infty w(t)\left(\int_x^t\frac{a(\tau)}{B(\tau)}u(\tau)\,d\tau\right)^q dt\right)^{\frac{r}{q}}\left(\int_0^x\left(\frac{B(s)}{V(s)}\right)^{p'}v(s)\,ds\right)^{\frac{r}{q'}}\left(\frac{B(x)}{V(x)}\right)^{p'}v(x)\,dx\right)^{\frac{1}{r}}\\
&\qquad+\left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}}\left(\int_0^\infty\left(\int_0^x a(y)u(y)\,dy\right)^q w(x)\,dx\right)^{\frac{1}{q}}.
\end{aligned}$$

The proof is completed. □

6. The boundedness of $T_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions

In this section we combine the results from previous two sections to present the characterization of the boundedness of $T_{u,b}$ from $L^p(v)$ into $\mathrm{Ces}_q(w,a)$ on the cone of monotone non-increasing functions.

Theorem 6.1. *Let $1<p,q<\infty$ and $b\in\mathcal{W}(0,\infty)$ be such that $b(t)>0$ for a.e. $t\in(0,\infty)$. Assume that $u\in\mathcal{W}(0,\infty)\cap C(0,\infty)$ and $a,v,w\in\mathcal{W}(0,\infty)$. Moreover, assume that condition (1.1) holds.*

(i) *If $p\le q$, then*

$$\begin{aligned}
&\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x(T_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}\\
&\quad\approx\sup_{t\in(0,\infty)}\left(\int_0^t V(x)^{p'}v(x)\left(\int_x^t\left(\sup_{s\le\tau}u(\tau)V(\tau)^{-2}\right)a(s)\,ds\right)^{p'}dx\right)^{\frac{1}{p'}}\left(\int_t^\infty w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\sup_{t\in(0,\infty)}\left(\int_0^t V(x)^{p'}v(x)\,dx\right)^{\frac{1}{p'}}\left(\int_t^\infty\left(\int_t^y\left(\sup_{s\le\tau}u(\tau)V(\tau)^{-2}\right)a(s)\,ds\right)^q w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\sup_{x\in(0,\infty)}\left(\int_{[x,\infty)}d\left(-\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\sup_{x\in(0,\infty)}\left(\int_{(0,x]}A(t)^{p'}\,d\left(-\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)^{\frac{1}{p'}}\left(\int_x^\infty w(y)\,dy\right)^{\frac{1}{q}}\\
&\qquad+\left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)^{\frac{1}{p'}}\right)\\
&\qquad+\sup_{x\in(0,\infty)}\left(\int_0^x\left(\int_0^t a(y)\bar{u}(y)\,dy\right)^q w(t)\,dt\right)^{\frac{1}{q}}\left(\int_x^\infty V(s)^{-p'}v(s)\,ds\right)^{\frac{1}{p'}}
\end{aligned}$$

$$+\sup_{x\in(0,\infty)}\left(\int_x^\infty w(t)\,dt\right)^{\frac{1}{q}}\left(\int_0^x\left(\int_0^s a(y)\bar{u}(y)\,dy\right)^{p'}V(s)^{-p'}v(s)\,ds\right)^{\frac{1}{p'}}$$

$$+\sup_{x\in(0,\infty)}\left(\int_x^\infty\left(\int_x^t\frac{a(\tau)}{B(\tau)}\bar{u}(\tau)\,d\tau\right)^q w(t)\,dt\right)^{\frac{1}{q}}\left(\int_0^x\left(\frac{B(s)}{V(s)}\right)^{p'}v(s)\,ds\right)^{\frac{1}{p'}}$$

$$+\sup_{x\in(0,\infty)}\left(\int_x^\infty w(t)\,dt\right)^{\frac{1}{q}}\left(\int_0^x\left(\int_s^x\frac{a(\tau)}{B(\tau)}\bar{u}(\tau)\,d\tau\right)^{p'}\left(\frac{B(s)}{V(s)}\right)^{p'}v(s)\,ds\right)^{\frac{1}{p'}}$$

$$+\left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}}\left(\int_0^\infty\left(\int_0^x a(t)\bar{u}(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}};$$

(ii) *If $q<p$, then*

$$\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x(T_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$

$$\approx\left(\int_0^\infty\left(\int_0^t V(x)^{p'}v(x)\,dx\right)^{\frac{r}{q'}}V(t)^{p'}v(t)\left(\int_t^\infty\left(\int_t^z\left(\sup_{s\le y}u(y)V(y)^{-2}\right)a(s)\,ds\right)^q w(z)\,dz\right)^{\frac{r}{q}}dt\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_0^t V(x)^{p'}v(x)\left(\int_x^t\left(\sup_{s\le y}u(y)V(y)^{-2}\right)a(s)\,ds\right)^{p'}dx\right)^{\frac{r}{p'}}\left(\int_z^\infty w(s)\,ds\right)^{\frac{r}{p}}w(t)\,dt\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_{[x,\infty)}d\left(-\left(\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}}\left(\int_0^x A(y)^q w(y)\,dy\right)^{\frac{r}{p}}x^q w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_{(0,x]}A(t)^{p'}\,d\left(-\left(\sup_{t\le\tau}u(\tau)^{p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}}\left(\int_x^\infty w(y)\,dy\right)^{\frac{r}{p}}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty A(y)^q w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\le\tau}u(\tau)V(\tau)^{-2}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)^{\frac{1}{p'}}\right)$$

$$+\left(\int_0^\infty\left(\int_0^x\left(\int_0^t a(y)\bar{u}(y)\,dy\right)^q w(t)\,dt\right)^{\frac{r}{p}}\left(\int_x^\infty V(z)^{-p'}v(z)\,dz\right)^{\frac{r}{p'}}\left(\int_0^x a(y)u(y)\,dy\right)^q w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_x^\infty w(t)\,dt\right)^{\frac{r}{p}}\left(\int_0^x\left(\int_0^z a(y)\bar{u}(y)\,dy\right)^{p'}V(z)^{-p'}v(z)\,dz\right)^{\frac{r}{p'}}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_x^\infty w(t)\,dt\right)^{\frac{r}{p}}\left(\int_0^x\left(\int_z^x\frac{a(\tau)}{B(\tau)}\bar{u}(\tau)\,d\tau\right)^{p'}\left(\frac{B(z)}{V(z)}\right)^{p'}v(z)\,dz\right)^{\frac{r}{p'}}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_x^\infty w(t)\left(\int_x^t\frac{a(\tau)}{B(\tau)}\bar{u}(\tau)\,d\tau\right)^q dt\right)^{\frac{r}{q}}\left(\int_0^x\left(\frac{B(s)}{V(s)}\right)^{p'}v(s)\,ds\right)^{\frac{r}{q'}}\left(\frac{B(x)}{V(x)}\right)^{p'}v(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}}\left(\int_0^\infty\left(\int_0^x a(y)\bar{u}(y)\,dy\right)^q w(x)\,dx\right)^{\frac{1}{q}}.$$

Proof. By (1.2), we have that

$$\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x(T_{u,b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}\approx\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x(R_u f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$

$$+\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x(P_{\bar{u},b}f)(t)a(t)\,dt\right)^q w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}.$$

It remains to apply Theorems 4.1 and 5.1. □

7. The boundedness of M_γ from $\Lambda^p(v)$ into $\Gamma^q(w)$

Suppose that f is a measurable a.e. finite function on $\mathbb{R}^n$. Then its non-increasing rearrangement f^* is given by

$$f^*(t)=\inf\{\lambda>0:|\{x\in\mathbb{R}^n:|f(x)|>\lambda\}|\le t\},\quad t\in(0,\infty),$$

and let f^{**} denotes the Hardy-Littlewood maximal function of f^*, i.e.

$$f^{**}(t):=\frac{1}{t}\int_0^t f^*(\tau)\,d\tau,\quad t>0.$$

Quite many familiar function spaces can be defined by using the non-increasing rearrangement of a function. One of the most important classes of such spaces are the so-called classical Lorentz spaces.

Let $p\in(0,\infty)$ and $w\in\mathcal{W}(0,\infty)$. Then the classical Lorentz spaces $\Lambda^p(w)$ and $\Gamma^p(w)$ consist of all measurable functions f on $\mathbb{R}^n$ for which $\|f\|_{\Lambda^p(w)}:=\|f^*\|_{p,w,(0,\infty)}<\infty$ and $\|f\|_{\Gamma^p(w)}:=\|f^{**}\|_{p,w,(0,\infty)}<\infty$, respectively. For more information about the Lorentz Λ and Γ spaces see e.g. [9] and the references therein.

The fractional maximal operator, M_γ, $\gamma\in(0,n)$, is defined at a locally integrable function f on $\mathbb{R}^n$ by

$$(M_\gamma f)(x):=\sup_{Q\ni x}|Q|^{\gamma/n-1}\int_Q|f(y)|\,dy,\quad x\in\mathbb{R}^n.$$

It was shown in [10, Theorem 1.1] that

$$(M_\gamma f)^*(t)\lesssim\sup_{\tau>t}\tau^{\gamma/n-1}\int_0^\tau f^*(y)\,dy\lesssim(M_\gamma\tilde{f})^*(t)\tag{7.1}$$

for every locally integrable function f on $\mathbb{R}^n$ and $t\in(0,\infty)$, where $\tilde{f}(x):=f^*(\omega_n|x|^n)$ and ω_n is the volume of the unit ball in $\mathbb{R}^n$.

The characterization of the boundedness of M_γ between classical Lorentz spaces $\Lambda^p(v)$ and $\Lambda^q(w)$ was obtained in [10] for the particular case when $1<p\le q<\infty$ and in [36, Theorem 2.10] in the case of more general operators and for extended range of p and q (For the characteriation of the boundedness of more general fractional maximal functions between $\Lambda^p(v)$ and $\Lambda^q(w)$, see [34], and the references therein).

As an application of obtained results, we calculate the norm of the fractional maximal function M_γ from $\Lambda^p(v)$ into $\Gamma^q(w)$.

Theorem 7.1. *Let $1<p,q<\infty$ and $0<\gamma<n$. Assume that $v,w\in\mathcal{W}(0,\infty)$.*

(i) *If $p\le q$, then*

$$\begin{aligned}
&\|M_\gamma\|_{\Lambda^p(v)\to\Gamma^q(w)}\\
&\approx\sup_{t\in(0,\infty)}\Big(\int_0^t V(x)^{p'}v(x)\Big(\int_x^t\Big(\sup_{s\le\tau}\tau^{\frac{\gamma}{n}}V(\tau)^{-2}\Big)ds\Big)^{p'}dx\Big)^{\frac{1}{p'}}\Big(\int_t^\infty y^{-q}w(y)\,dy\Big)^{\frac{1}{q}}\\
&\quad+\sup_{t\in(0,\infty)}\Big(\int_0^t V(x)^{p'}v(x)\,dx\Big)^{\frac{1}{p'}}\Big(\int_t^\infty\Big(\int_t^y\Big(\sup_{s\le\tau}\tau^{\frac{\gamma}{n}}V(\tau)^{-2}\Big)ds\Big)^q y^{-q}w(y)\,dy\Big)^{\frac{1}{q}}\\
&\quad+\sup_{x\in(0,\infty)}\Big(\int_{[x,\infty)}d\Big(-\sup_{t\le\tau}\tau^{\frac{\gamma}{n}p'}V(\tau)^{-2p'}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)\Big)\Big)^{\frac{1}{p'}}\Big(\int_0^x w(y)\,dy\Big)^{\frac{1}{q}}\\
&\quad+\sup_{x\in(0,\infty)}\Big(\int_{(0,x]}t^{p'}\,d\Big(-\sup_{t\le\tau}\tau^{\frac{\gamma}{n}p'}V(\tau)^{-2p'}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)\Big)\Big)^{\frac{1}{p'}}\Big(\int_x^\infty y^{-q}w(y)\,dy\Big)^{\frac{1}{q}}\\
&\quad+\Big(\int_0^\infty w(y)\,dy\Big)^{\frac{1}{q}}\lim_{t\to\infty}\Big(\sup_{t\le\tau}\tau^{\frac{\gamma}{n}}V(\tau)^{-2}\Big(\int_0^\tau V(s)^{p'}v(s)\,ds\Big)^{\frac{1}{p'}}\Big)\\
&\quad+\sup_{x\in(0,\infty)}\Big(\int_0^x t^{\frac{\gamma}{n}q}w(t)\,dt\Big)^{\frac{1}{q}}\Big(\int_x^\infty V(s)^{-p'}v(s)\,ds\Big)^{\frac{1}{p'}}
\end{aligned}$$

$$+\sup_{x\in(0,\infty)}\left(\int_x^\infty y^{-q}w(y)\,dy\right)^{\frac{1}{q}}\left(\int_0^x t^{(\frac{\gamma}{n}+1)p'}V(t)^{-p'}v(t)\,dt\right)^{\frac{1}{p'}}$$

$$+\sup_{x\in(0,\infty)}\left(\int_x^\infty \left(t^{\frac{\gamma}{n}}-x^{\frac{\gamma}{n}}\right)^q t^{-q}w(t)\,dt\right)^{\frac{1}{q}}\left(\int_0^x s^{p'}V(s)^{-p'}v(s)\,ds\right)^{\frac{1}{p'}}$$

$$+\sup_{x\in(0,\infty)}\left(\int_x^\infty t^{-q}w(t)\,dt\right)^{\frac{1}{q}}\left(\int_0^x \left(x^{\frac{\gamma}{n}}-s^{\frac{\gamma}{n}}\right)^{p'} s^{p'}V(s)^{-p'}v(s)\,ds\right)^{\frac{1}{p'}}$$

$$+\left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}}\left(\int_0^\infty x^{\frac{\gamma}{n}q}w(x)\,dx\right)^{\frac{1}{q}};$$

(ii) *If $q<p$, then*

$$\|M_\gamma\|_{\Lambda^p(v)\to\Gamma^q(w)}$$

$$\approx \left(\int_0^\infty\left(\int_0^t V(x)^{p'}v(x)\,dx\right)^{\frac{r}{q'}}V(t)^{p'}v(t)\left(\int_t^\infty\left(\int_t^z\left(\sup_{s\le y}y^{\frac{\gamma}{n}}V(y)^{-2}\right)ds\right)^q z^{-q}w(z)\,dz\right)^{\frac{r}{q}}dt\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_0^t V(x)^{p'}v(x)\left(\int_x^t\left(\sup_{s\le y}y^{\frac{\gamma}{n}}V(y)^{-2}\right)ds\right)^{p'}dx\right)^{\frac{r}{p'}}\left(\int_z^\infty s^{-q}w(s)\,ds\right)^{\frac{r}{p}}t^{-q}w(t)\,dt\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_{[x,\infty)}d\left(-\left(\sup_{t\le\tau}\tau^{\frac{\gamma}{n}p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}}\left(\int_0^x w(y)\,dy\right)^{\frac{r}{p}}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_{(0,x]}t^{p'}\,d\left(-\left(\sup_{t\le\tau}\tau^{\frac{\gamma}{n}p'}V(\tau)^{-2p'}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)\right)\right)\right)^{\frac{r}{p'}}\left(\int_x^\infty y^{-q}w(y)\,dy\right)^{\frac{r}{p}}x^{-q}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty w(y)\,dy\right)^{\frac{1}{q}}\lim_{t\to\infty}\left(\sup_{t\le\tau}\tau^{\frac{\gamma}{n}p'}V(\tau)^{-2}\left(\int_0^\tau V(s)^{p'}v(s)\,ds\right)^{\frac{1}{p'}}\right)$$

$$+\left(\int_0^\infty\left(\int_0^x t^{\frac{\gamma}{n}q}w(t)\,dt\right)^{\frac{r}{p}}\left(\int_x^\infty V(z)^{-p'}v(z)\,dz\right)^{\frac{r}{p'}}x^{\frac{\gamma}{n}q}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_x^\infty t^{-q}w(t)\,dt\right)^{\frac{r}{p}}\left(\int_0^x z^{(\frac{\gamma}{n}+1)p'}V(z)^{-p'}v(z)\,dz\right)^{\frac{r}{p'}}x^{-q}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_x^\infty t^{-q}w(t)\,dt\right)^{\frac{r}{p}}\left(\int_0^x \left(x^{\frac{\gamma}{n}}-z^{\frac{\gamma}{n}}\right)^{p'}z^{p'}V(z)^{-p'}v(z)\,dz\right)^{\frac{r}{p'}}x^{-q}w(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty\left(\int_x^\infty \left(t^{\frac{\gamma}{n}}-x^{\frac{\gamma}{n}}\right)^q t^{-q}w(t)\,dt\right)^{\frac{r}{q}}\left(\int_0^x s^{p'}V(s)^{-p'}v(s)\,ds\right)^{\frac{r}{q'}}x^{p'}V(x)^{-p'}v(x)\,dx\right)^{\frac{1}{r}}$$

$$+\left(\int_0^\infty v(s)\,ds\right)^{-\frac{1}{p}}\left(\int_0^\infty x^{\frac{\gamma}{n}q}w(x)\,dx\right)^{\frac{1}{q}}.$$

Proof. From inequalities (7.1), we have that

$$\|M_\gamma\|_{\Lambda^p(v)\to\Gamma^q(w)}\approx\sup_{f\in\mathfrak{M}^{+,\downarrow}(0,\infty)}\frac{\left(\int_0^\infty\left(\int_0^x(T_{u,b}f)(t)\,dt\right)^q x^{-q}w(x)\,dx\right)^{\frac{1}{q}}}{\left(\int_0^\infty f(s)^p v(s)\,ds\right)^{\frac{1}{p}}}$$

with $u(\tau)=\tau^{\gamma/n}$ and $b\equiv 1$. Note that

$$\sup_{0<t<\infty}\frac{u(t)}{B(t)}\int_0^t\frac{b(\tau)}{u(\tau)}\,d\tau<\infty$$

in this case. So, it remains to apply Theorem 6.1. □

References

[1] R. Askey and R. P. Boas Jr., *Some integrability theorems for power series with positive coefficients*, Mathematical Essays Dedicated to A. J. Macintyre, Ohio Univ. Press, Athens, Ohio, 1970, pp. 23–32. MR0277956 (43 #3689)

[2] S. V. Astashkin and L. Maligranda, *Structure of Cesàro function spaces*, Indag. Math. (N.S.) **20** (2009), no. 3, 329–379, DOI 10.1016/S0019-3577(10)00002-9. MR2639977 (2011c:46056)

[3] ———, *Structure of Cesàro function spaces: a survey*, Banach Center Publ. **102** (2014), 13–40.

[4] G. Bennett, *Factorizing the classical inequalities*, Mem. Amer. Math. Soc. **120** (1996), no. 576, viii+130, DOI 10.1090/memo/0576. MR1317938 (96h:26020)

[5] G. Bennett and K.- G. Grosse-Erdmann, *Weighted Hardy inequalities for decreasing sequences and functions*, Math. Ann. **334** (2006), no. 3, 489–531, DOI 10.1007/s00208-005-0678-7. MR2207873 (2006m:26038)

[6] R. P. Boas Jr., *Integrability theorems for trigonometric transforms*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 38, Springer-Verlag New York Inc., New York, 1967. MR0219973 (36 #3043)

[7] ———, *Some integral inequalities related to Hardy's inequality*, J. Analyse Math. **23** (1970), 53–63. MR0274685 (43 #447)

[8] M. Carro, A. Gogatishvili, J. Martin, and L. Pick, *Weighted inequalities involving two Hardy operators with applications to embeddings of function spaces*, J. Operator Theory **59** (2008), no. 2, 309–332. MR2411048 (2009f:26024)

[9] M. Carro, L. Pick, J. Soria, and V. D. Stepanov, *On embeddings between classical Lorentz spaces*, Math. Inequal. Appl. **4** (2001), no. 3, 397–428, DOI 10.7153/mia-04-37. MR1841071 (2002d:46026)

[10] A. Cianchi, R. Kerman, B. Opic, and L. Pick, *A sharp rearrangement inequality for the fractional maximal operator*, Studia Math. **138** (2000), no. 3, 277–284. MR1758860 (2001h:42029)

[11] S. Chen, Y. Cui, H. Hudzik, and B. Sims, *Geometric properties related to fixed point theory in some Banach function lattices*, Handbook of metric fixed point theory, Kluwer Acad. Publ., Dordrecht, 2001, pp. 339–389. MR1904283 (2003f:46031)

[12] Y. Cui and R. Płuciennik, *Local uniform nonsquareness in Cesàro sequence spaces*, Comment. Math. Prace Mat. **37** (1997), 47–58. MR1608225 (99b:46025)

[13] Y. Cui and H. Hudzik, *Some geometric properties related to fixed point theory in Cesàro spaces*, Collect. Math. **50** (1999), no. 3, 277–288. MR1744077 (2001f:46033)

[14] ———, *Packing constant for Cesaro sequence spaces*, Proceedings of the Third World Congress of Nonlinear Analysts, Part 4 (Catania, 2000), 2001, pp. 2695–2702, DOI 10.1016/S0362-546X(01)00389-3. MR1972393 (2004c:46033)

[15] Y. Cui, H. Hudzik, and Y. Li, *On the Garcia-Falset coefficient in some Banach sequence spaces*, Function spaces (Poznań, 1998), Lecture Notes in Pure and Appl. Math., vol. 213, Dekker, New York, 2000, pp. 141–148. MR1772119 (2001h:46009)

[16] M. Cwikel and E. Pustylnik, *Weak type interpolation near "endpoint" spaces*, J. Funct. Anal. **171** (2000), no. 2, 235–277, DOI 10.1006/jfan.1999.3502. MR1745635 (2001b:46118)

[17] R. Ya. Doktorskii, *Reiterative relations of the real interpolation method*, Dokl. Akad. Nauk SSSR **321** (1991), no. 2, 241–245 (Russian); English transl., Soviet Math. Dokl. **44** (1992), no. 3, 665–669. MR1153547 (93b:46143)

[18] W. D. Evans and B. Opic, *Real interpolation with logarithmic functors and reiteration*, Canad. J. Math. **52** (2000), no. 5, 920–960, DOI 10.4153/CJM-2000-039-2. MR1782334 (2001i:46115)

[19] A. Gogatishvili, M. Johansson, C. A. Okpoti, and L.-E. Persson, *Characterisation of embeddings in Lorentz spaces*, Bull. Austral. Math. Soc. **76** (2007), no. 1, 69–92, DOI 10.1017/S0004972700039484. MR2343440 (2008j:46017)

[20] A. Gogatishvili and R.Ch. Mustafayev, *Weighted iterated Hardy-type inequalities*, Math. Inequal. Appl. **20** (2017), no. 3, 683–728. MR3653914

[21] ———, *Iterated Hardy-type inequalities involving suprema*, Math. Inequal. Appl. **20** (2017), no. 4, 901–927. MR3711402

[22] A. Gogatishvili, R. Mustafayev, and T. Ünver, *Embeddings between weighted Copson and Cesàro function spaces*, Czechoslovak Math. J. **67(142)** (2017), no. 4, 1105–1132. MR3736022

[23] A. Gogatishvili, B. Opic, and L. Pick, *Weighted inequalities for Hardy-type operators involving suprema*, Collect. Math. **57** (2006), no. 3, 227–255.

[24] A. Gogatishvili and L. Pick, *Embeddings and duality theorems for weak classical Lorentz spaces*, Canad. Math. Bull. **49** (2006), no. 1, 82–95, DOI 10.4153/CMB-2006-008-3. MR2198721

[25] ———, *A reduction theorem for supremum operators*, J. Comput. Appl. Math. **208** (2007), no. 1, 270–279. MR2347749 (2009a:26013)

[26] A. Gogatishvili and V. D. Stepanov, *Reduction theorems for weighted integral inequalities on the cone of monotone functions*, Uspekhi Mat. Nauk **68** (2013), no. 4(412), 3–68, DOI 10.1070/rm2013v068n04abeh004849 (Russian, with Russian summary); English transl., Russian Math. Surveys **68** (2013), no. 4, 597–664. MR3154814

[27] K.-G. Grosse-Erdmann, *The blocking technique, weighted mean operators and Hardy's inequality*, Lecture Notes in Mathematics, vol. 1679, Springer-Verlag, Berlin, 1998. MR1611898 (99d:26024)

[28] A. A. Jagers, *A note on Cesàro sequence spaces*, Nieuw Arch. Wisk. (3) **22** (1974), 113–124. MR0348444 (50 #942)

[29] A. Kamińska and D. Kubiak, *On the dual of Cesàro function space*, Nonlinear Anal. **75** (2012), no. 5, 2760–2773, DOI 10.1016/j.na.2011.11.019. MR2878472 (2012m:46034)

[30] R. Kerman and L. Pick, *Optimal Sobolev imbeddings*, Forum Math. **18** (2006), no. 4, 535–570, DOI 10.1515/FORUM.2006.028. MR2254384 (2007g:46052)

[31] M. Křepela, *Integral conditions for Hardy-type operators involving suprema*, Collect. Math. **68** (2017), no. 1, 21–50, DOI 10.1007/s13348-016-0170-6. MR3591463

[32] G.M. Leibowitz, *A note on the Cesàro sequence spaces*, Tamkang J. Math. **2** (1971), 151–157.

[33] V. G. Maz'ja, *Sobolev spaces*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1985. Translated from the Russian by T. O. Shaposhnikova. MR817985

[34] R. Ch. Mustafayev and N. Bilgiçli, *Generalized fractional maximal functions in Lorentz spaces* Λ, J. Math. Inequal. **12** (2018), no. 3, 827–851, DOI 10.7153/jmi-2018-12-62. MR3857365

[35] R. Oinarov, *Two-sided estimates for the norm of some classes of integral operators*, Trudy Mat. Inst. Steklov. **204** (1993), no. Issled. po Teor. Differ. Funktsii Mnogikh Peremen. i ee Prilozh. 16, 240–250 (Russian); English transl., Proc. Steklov Inst. Math. **3 (204)** (1994), 205–214. MR1320028

[36] B. Opic, *On boundedness of fractional maximal operators between classical Lorentz spaces*, Function spaces, differential operators and nonlinear analysis (Pudasjärvi, 1999), Acad. Sci. Czech Repub., Prague, 2000, pp. 187–196. MR1755309 (2001g:42043)

[37] B. Opic and A. Kufner, *Hardy-type inequalities*, Pitman Research Notes in Mathematics Series, vol. 219, Longman Scientific & Technical, Harlow, 1990. MR1069756 (92b:26028)

[38] L. Pick, *Supremum operators and optimal Sobolev inequalities*, Function spaces, differential operators and nonlinear analysis (Pudasjärvi, 1999), Acad. Sci. Czech Repub., Prague, 2000, pp. 207–219. MR1755311 (2000m:46075)

[39] ———, *Optimal Sobolev embeddings—old and new*, Function spaces, interpolation theory and related topics (Lund, 2000), de Gruyter, Berlin, 2002, pp. 403–411. MR1943297 (2003j:46054)

[40] *Programma van Jaarlijkse Prijsvragen (Annual Problem Section)*, Nieuw Arch. Wiskd. **16** (1968), 47–51.

[41] D. V. Prokhorov and V. D. Stepanov, *On weighted Hardy inequalities in mixed norms*, Proc. Steklov Inst. Math. **283** (2013), 149–164.

[42] ———, *Weighted estimates for a class of sublinear operators*, Dokl. Akad. Nauk **453** (2013), no. 5, 486–488 (Russian); English transl., Dokl. Math. **88** (2013), no. 3, 721–723. MR3203323

[43] ———, *Estimates for a class of sublinear integral operators*, Dokl. Akad. Nauk **456** (2014), no. 6, 645–649 (Russian); English transl., Dokl. Math. **89** (2014), no. 3, 372–377. MR3287911

[44] D. V. Prokhorov, *On the boundedness of a class of sublinear integral operators*, Dokl. Akad. Nauk **92** (2015), no. 2, 602–605 (Russian).

[45] E. Pustylnik, *Optimal interpolation in spaces of Lorentz-Zygmund type*, J. Anal. Math. **79** (1999), 113–157, DOI 10.1007/BF02788238. MR1749309 (2001a:46028)

[46] G. È. Shambilova, *Weighted inequalities for a class of quasilinear integral operators on the cone of monotone functions*, Sibirsk. Mat. Zh. **55** (2014), no. 4, 912–936 (Russian, with Russian summary); English transl., Sib. Math. J. **55** (2014), no. 4, 745–767. MR3242605

[47] J.-S. Shiue, *On the Cesàro sequence spaces*, Tamkang J. Math. **1** (1970), no. 1, 19–25.

[48] ———, *A note on Cesàro function space*, Tamkang J. Math. **1** (1970), no. 2, 91–95. MR0276751 (43 #2491)

[49] V. D. Stepanov, *The weighted Hardy's inequality for nonincreasing functions*, Trans. Amer. Math. Soc. **338** (1993), no. 1, 173–186, DOI 10.2307/2154450. MR1097171

[50] V. D. Stepanov and G. È. Shambilova, *Weight boundedness of a class of quasilinear operators on the cone of monotone functions*, Dokl. Math. **90** (2014), no. 2, 569–572.

[51] P. W. Sy, W. Y. Zhang, and P. Y. Lee, *The dual of Cesàro function spaces*, Glas. Mat. Ser. III **22(42)** (1987), no. 1, 103–112 (English, with Serbo-Croatian summary). MR940098 (89g:46059)

Rza Mustafayev, Department of Mathematics, Faculty of Science, Karamanoglu Mehmetbey University, Karaman, 70100, Turkey
E-mail address: rzamustafayev@gmail.com

Nevin Bilgiçli, Department of Mathematics, Faculty of Science and Arts, Kirikkale University, 71450 Yahsihan, Kirikkale, Turkey
E-mail address: nevinbilgicli@gmail.com