

Electronic structure of a multiply-charged nitrogen atom

Christian Buth^{1,2,3,4, a)} and Mau Hsiung Chen (陳茂雄)⁵

¹⁾Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

²⁾The PULSE Institute for Ultrafast Energy Science, SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA

³⁾Department of Physics and Astronomy, Louisiana State University, Baton Rouge, Louisiana 70803, USA

⁴⁾Argonne National Laboratory, Argonne, Illinois 60439, USA

⁵⁾Physics Division, Lawrence Livermore National Laboratory, Livermore, California 94550, USA

(Dated: 15 April 2012)

I. DESCRIPTION

Here we provide all atomic electronic structure data that was used to create a rate-equation model for a nitrogen atom. Please consider Sec. II of the paper for a description of the employed methods. For a multiply-ionized nitrogen atom, we gather the electron binding energies in Tables III, VIII, and XIII. The Auger decay widths are listed in Tables IV and IX with the corresponding transition energies provided in Tables V and X; the x-ray fluorescence decay widths are presented in Tables I, VI and XI with the respective transition energies collected in Tables II, VII and XII.

II. AVERAGING OVER FINE-STRUCTURE RESOLVED ENERGIES AND DECAY WIDTHS

Assuming that the population with electrons of the $2p_{1/2}$ and $2p_{3/2}$ orbitals in $1s^\ell 2s^m 2p_{1/2}^{n_{1/2}} 2p_{3/2}^{n_{3/2}}$ jj coupled configurations is probabilistic, we average over configurations as follows. For decay widths in jj coupling $\tilde{\Gamma}_{\ell m n_{1/2} n_{3/2}}$, we calculate the weighted sum over the accessible decay channels via

$$\Gamma_{\ell m n} = \sum_{(n_{1/2}, n_{3/2}) \in \mathbb{K}_n} f(n_{1/2}, n_{3/2}) \tilde{\Gamma}_{\ell m n_{1/2} n_{3/2}}, \quad (1)$$

where the total of the weights is unity, i.e., $\sum_{(n_{1/2}, n_{3/2}) \in \mathbb{K}_n} f(n_{1/2}, n_{3/2}) = 1$. The sum runs over all possible distributions of the $0 \leq n \leq 3$ electrons over the $2p_{1/2}$ and the $2p_{3/2}$ orbitals: $\mathbb{K}_n = \{(n_{1/2}, n_{3/2}) \mid n_{1/2} + n_{3/2} = n \wedge 0 \leq n_{1/2} \leq 2 \wedge 0 \leq n_{3/2} \leq 4\}$, using the probabilistic weights

$$f(n_{1/2}, n_{3/2}) = \frac{\binom{2}{n_{1/2}} \binom{4}{n_{3/2}}}{\binom{6}{n_{1/2} + n_{3/2}}}. \quad (2)$$

To compute the average for binding energies subject to multiplet splitting and transition energies between multiplets, we need to multiply the relevant energies in jj -coupling scheme with the weights (2). However, in contrast to decay widths (1), for energies, we need not only the probabilistic weight for a distribution of n electrons over $2p_{1/2}^{n_{1/2}} 2p_{3/2}^{n_{3/2}}$ but we also have to average over contributions from the cases where the electron(s)—that is(are) involved in a transition or whose binding energy is calculated—stem(s) from $2p_{1/2}^{n_{1/2}}$ or $2p_{3/2}^{n_{3/2}}$. This is because, for the determination of total decay widths, we merely add all possible decays from a specific configuration $2p_{1/2}^{n_{1/2}} 2p_{3/2}^{n_{3/2}}$ whereas energies cannot simply be added but need to be additionally weighted according to their probabilistic contribution to the average energy of the LS coupled configuration. In other words, to obtain LS coupled energies, first, we average over the binding energies or transition energies starting from a single initial jj coupled state by multiplying it with proper probabilities; and, second, we average over the various initial jj coupled states arising from fine-structure splitting, i.e., from distributing n electrons over the energetically distinct $2p_{1/2}$ and $2p_{3/2}$ orbitals. For this purpose, we introduce into Eq. (1) the additional probabilistic weights $g(n_{1/2}, n_{3/2}, j)$ with $j \in \{1/2, 3/2\}$ and $\sum_{j=1/2}^{3/2} g(n_{1/2}, n_{3/2}, j) = 1$ that determine how likely one picks an(two) electron(s)¹ from the configuration $2p_{1/2}^{n_{1/2}} 2p_{3/2}^{n_{3/2}}$ by writing

$$E_{\ell m n} = \sum_{(n_{1/2}, n_{3/2}) \in \mathbb{K}_n} f(n_{1/2}, n_{3/2}) \times \sum_{j=1/2}^{3/2} g(n_{1/2}, n_{3/2}, j) \tilde{E}_{\ell m n_{1/2} n_{3/2} j}. \quad (3)$$

The new weights $g(n_{1/2}, n_{3/2}, j)$ are determined as follows. There are three different types of Auger decays in LS coupling: $1s-2s2s$, $1s-2s2p$, and $1s-2p2p$ transitions. In jj coupling, $1s-2s2p$ splits into $1s-2s2p_{1/2}$ and $1s-2s2p_{3/2}$ due to spin-orbit interaction.² For the same reason, $1s-2p2p$ splits up into $1s-2p_{1/2}2p_{1/2}$, $1s-2p_{1/2}2p_{3/2}$, and $1s-2p_{3/2}2p_{3/2}$. For $1s-2s2s$ Auger de-

^{a)}Corresponding author; Electronic mail: christian.buth@web.de

cay, the weights are $g(n_{1/2}, n_{3/2}, j) = 1/2$ because the transition energies $\bar{E}_{\ell m n_{1/2} n_{3/2} j}$ are in this case independent of j and we set the energies for $j \in \{1/2, 3/2\}$ to the same value. This holds also true for $1s-2s2p$ Auger transitions in jj coupling for charge states of nitrogen with only one $2p$ electron or a $1s-2p2p$ transition for an N ion with only two $2p$ electrons. In other words, in these special cases, one needs to average only over the initial states [Eqs. (1) and (2)] to obtain the LS averaged energies, in the same way as is done to determine the LS averaged Auger widths. For two or three $2p$ electrons, the probabilistic weight of a $1s-2s2p_j$ transition is $g(n_{1/2}, n_{3/2}, j) = \binom{n_j}{1} / \binom{n_{1/2} + n_{3/2}}{1}$ with $j \in \{1/2, 3/2\}$. The weight of a $1s-2p_i2p_j$ transition for three $2p$ electrons is $g(n_{1/2}, n_{3/2}, j) = \binom{n_{1/2}}{p_j} \binom{n_{3/2}}{q_j} / \binom{n_{1/2} + n_{3/2}}{2}$ where $p_j + q_j = 2$ with $0 \leq p_j, q_j \leq 2$ and p_j and q_j are the number of $2p_{1/2}$ and $2p_{3/2}$ electrons, respectively, involved in the Auger decay.¹ For the weighting of binding energies and x-ray fluorescence energies, the averaging procedure is the same as for $2s2p$ Auger energies. If only $2s$ electrons are involved in a transition or the binding energy of an s electron shall be calculated, then $g(n_{1/2}, n_{3/2}, j) = 1/2$. This choice of the weights is also true, for the binding energy and a fluorescence decay of a $2p$ electron of nitrogen in a charge state with only a single $2p$ electron. However, the weights are $g(n_{1/2}, n_{3/2}, j) = \binom{n_j}{1} / \binom{n_{1/2} + n_{3/2}}{1}$, if nitrogen is in a charge state with two or three $2p$ electrons and either a $2p$ electron is involved in a transition or the binding energy of a $2p$ electron is calculated.

REFERENCES

- ¹When we average over the transition energies in jj coupling scheme of $2p2p$ Auger decay in the case of three $2p$ electrons, the j label on the energies and the weights in Eq. (3) denotes the j value of the remaining $2p$ electron after Auger decay (in all other cases, the j label denotes the j value of the involved $2p$ electron). In this case, the weights are $g(n_{1/2}, n_{3/2}, j) = \frac{n_j}{3}$, i.e., they are given by the probability to choose one electron to remain in a $2p_{1/2}$ or a $2p_{3/2}$ orbital when two $2p$ electrons are removed. See text after Eq. (3) for a detailed discussion.
- ²E. U. Condon and G. H. Shortley, *The theory of atomic spectra* (Cambridge University Press, Cambridge, 1935) ISBN 978-0-521-09209-8.

TABLES

Initial Configuration		Electron involved in the decay
m	n	$2p$
1	3	9.97×10^{-4}
1	2	3.83×10^{-4}
1	1	8.27×10^{-5}
0	3	2.81×10^{-3}
0	2	1.06×10^{-3}
0	1	2.30×10^{-4}

TABLE I. No- K -shell-hole x-ray fluorescence decay widths for various configurations of a multiply-charged nitrogen atom ($1s^2 2s^m 2p^n$) in millielectronvolts. All transitions fill the inner-valence hole. All numbers correspond to E1 transitions.

Initial Configuration		Electron involved in the decay
m	n	$2p$
1	3	12.20
1	2	10.73
1	1	8.58
0	3	14.42
0	2	12.59
0	1	10.07

TABLE II. No- K -shell-hole x-ray fluorescence transition energies for various configurations of a multiply-charged nitrogen atom ($1s^2 2s^m 2p^n$) in electronvolts. All transitions fill the inner-valence hole. All numbers correspond to E1 transitions.

Configuration		Electron binding energy		
m	n	$1s$	$2s$	$2p$
2	3	411.74	24.27	12.08
2	2	432.46	38.78	28.04
2	1	458.03	55.70	47.12
2	0	488.28	74.99	—
1	3	431.24	40.50	26.58
1	2	455.62	57.56	44.96
1	1	484.67	76.49	66.41
1	0	518.22	97.68	—
0	3	453.93	—	43.14
0	2	481.79	—	63.89
0	1	514.15	—	87.61
0	0	550.80	—	—

TABLE III. No- K -shell-hole binding energies for various configurations of a multiply-charged nitrogen atom ($1s^2 2s^m 2p^n$) in electronvolts.

Initial Configuration		Electrons involved in the decay			
m	n	$2s\ 2s$	$2s\ 2p$	$2p\ 2p$	Total width
2	3	25.32	37.42	35.12	97.86
2	2	31.10	32.17	15.92	79.19
2	1	38.40	20.30	—	58.71
2	0	46.63	—	—	46.63
1	3	—	23.37	45.42	68.79
1	2	—	19.70	19.99	39.69
1	1	—	12.37	—	12.37
0	3	—	—	57.01	57.01
0	2	—	—	24.44	24.44

TABLE IV. Single- K -shell-hole Auger decay widths for various configurations of a multiply-charged nitrogen atom ($1s^1 2s^m 2p^n$) in millielectronvolts.

Initial Configuration		Electrons involved in the decay		
m	n	$2s\ 2s$	$2s\ 2p$	$2p\ 2p$
2	3	346.47	360.89	371.62
2	2	336.13	348.72	357.30
2	1	325.84	335.91	—
2	0	315.61	—	—
1	3	—	347.10	359.70
1	2	—	334.17	344.24
1	1	—	320.57	—
0	3	—	—	346.90
0	2	—	—	330.29

TABLE V. Single- K -shell-hole Auger transition energies for various configurations of a multiply-charged nitrogen atom ($1s^1 2s^m 2p^n$) in electronvolts. All transitions fill the core hole.

Initial Configuration		Electron involved in the decay	
m	n	$2s$	$2p$
2	3	—	0.46
2	2	—	0.36
2	1	—	0.21
2	0	2.08×10^{-10}	—
1	3	—	0.53
1	2	—	0.42
1	1	—	0.24
1	0	1.25×10^{-10}	—
0	3	—	0.62
0	2	—	0.48
0	1	—	0.28

TABLE VI. Single- K -shell-hole x-ray fluorescence decay widths for various configurations of a multiply-charged nitrogen atom ($1s^1 2s^m 2p^n$) in millielectronvolts. All transitions fill the core hole. The numbers in the “ $2p$ ” column correspond to E1 transitions and the numbers in the “ $2s$ ” column correspond to M1 transitions.

Initial Configuration		Electron involved in the decay	
m	n	$2s$	$2p$
2	3	—	399.67
2	2	—	404.42
2	1	—	410.91
2	0	413.29	—
1	3	—	404.66
1	2	—	410.66
1	1	—	418.25
1	0	420.54	—
0	3	—	410.79
0	2	—	417.90
0	1	—	426.54

TABLE VII. Single- K -shell-hole x-ray fluorescence transition energies for various configurations of a multiply-charged nitrogen atom ($1s^1 2s^m 2p^n$) in electronvolts.

Configuration		Electron binding energy		
m	n	$1s$	$2s$	$2p$
2	3	501.73	43.77	32.80
2	2	528.94	61.93	53.61
2	1	560.69	82.34	77.38
2	0	596.76	104.93	—
1	3	525.76	63.68	50.96
1	2	556.33	83.73	74.01
1	1	591.28	105.97	99.97
1	0	630.51	130.26	—
0	3	552.68	—	71.00
0	2	586.48	—	96.25
0	1	624.58	—	124.26
0	0	667.05	—	—

TABLE VIII. Single- K -shell-hole binding energies for various configurations of a multiply-charged nitrogen atom ($1s^1 2s^m 2p^n$) in electronvolts.

Initial Configuration		Electrons involved in the decay			
m	n	$2s\ 2s$	$2s\ 2p$	$2p\ 2p$	Total width
2	3	67.73	115.57	121.62	304.93
2	2	82.40	94.20	50.84	227.45
2	1	99.09	56.58	—	155.67
2	0	115.42	—	—	115.42
1	3	—	68.54	146.19	214.73
1	2	—	55.46	59.79	115.25
1	1	—	33.16	—	33.16
0	3	—	—	173.74	173.74
0	2	—	—	70.48	70.48

TABLE IX. Double- K -shell-hole Auger decay widths for various configurations of a multiply-charged nitrogen atom ($1s^0 2s^m 2p^n$) in millielectronvolts. All transitions fill the core hole.

Initial Configuration		Electron involved in the decay	
m	n	$2s$	$2p$
2	3	—	1.80
2	2	—	1.37
2	1	—	0.77
2	0	8.21×10^{-10}	—
1	3	—	2.03
1	2	—	1.54
1	1	—	0.87
1	0	4.66×10^{-10}	—
0	3	—	2.33
0	2	—	1.76
0	1	—	0.99

TABLE XI. Double- K -shell-hole x-ray fluorescence decay widths for various configurations of a multiply-charged nitrogen atom ($1s^0 2s^m 2p^n$) in millielectronvolts. All transitions fill the core hole. The numbers in the “ $2p$ ” column correspond to E1 transitions and the numbers in the “ $2s$ ” column correspond to M1 transitions.

Initial Configuration		Electron involved in the decay	
m	n	$2s$	$2p$
2	3	—	468.93
2	2	—	475.34
2	1	—	483.31
2	0	491.84	—
1	3	—	474.81
1	2	—	482.32
1	1	—	491.32
1	0	500.25	—
0	3	—	481.68
0	2	—	490.22
0	1	—	500.32

TABLE XII. Double- K -shell-hole x-ray fluorescence transition energies for various configurations of a multiply-charged nitrogen atom ($1s^0 2s^m 2p^n$) in electronvolts.

Initial Configuration		Electrons involved in the decay		
m	n	$2s\ 2s$	$2s\ 2p$	$2p\ 2p$
2	3	394.27	406.99	415.32
2	2	383.28	393.00	397.96
2	1	372.38	378.39	—
2	0	361.58	—	—
1	3	—	391.08	400.80
1	2	—	376.36	382.36
1	1	—	361.06	—
0	3	—	—	385.43
0	2	—	—	365.97

TABLE X. Double- K -shell-hole Auger transition energies for various configurations of a multiply-charged nitrogen atom ($1s^0 2s^m 2p^n$) in electronvolts.

Configuration		Electron binding energy	
m	n	$2s$	$2p$
2	3	67.81	60.02
2	2	89.32	85.35
2	1	112.93	113.45
2	0	138.68	—
1	3	90.60	81.53
1	2	113.87	108.96
1	1	139.27	139.20
1	0	166.79	—
0	3	—	104.80
0	2	—	134.36
0	1	—	166.73

TABLE XIII. Double- K -shell-hole binding energies for various configurations of a multiply-charged nitrogen atom ($1s^0 2s^m 2p^n$) in electronvolts.