

```

<< Units`  

<< PhysicalConstants`  

<< PlotLegends`  

SetDirectory[NotebookDirectory[]]  

Get["NitrogenRateEquations.m"]  

Get["NitrogenMoleculeRateEqs.m"]  

HartreeToeV := 27.2113845  

eVToHartree := 1 / HartreeToeV  

JouleToHartree := 2.29371269*^17  

MilliJouleToHartree := 1*^-3 * JouleToHartree  

BohrToAngstroem := 0.5291772083  

AngstroemToBohr := 1 / BohrToAngstroem  

MicrometerToBohr := 1*^4 * AngstroemToBohr  

BohrToMicrometer := 1 / MicrometerToBohr  

t0ToSecond := 24.18884326505*^-18  

t0ToFemtoSecond := 24.18884326505*^-3  

crossSectionAUtoBarn := 1*^8 * BohrToAngstroem^2  

crossSectionAUtoKiloBarn := 1*^-3 * crossSectionAUtoBarn  

crossSectionAUtoMegaBarn := 1*^-6 * crossSectionAUtoBarn  

crossSectionAUtoCm2 := crossSectionAUtoBarn * 1*^-24  

crossSectionCm2toAU := 1 / crossSectionAUtoCm2  

neginfinity := -1000 / t0ToFemtoSecond  

posinfinity := 10 000 / t0ToFemtoSecond

```

Dominant path analysis for a nitrogen atom in LCLS radiation

Christian Buth, 20 April 2012

Argonne National Laboratory, Argonne, Illinois 60439, USA

Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

The PULSE Institute for Ultrafast Energy Science, SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA

LCLS parameters

Gaussian x-ray flux pulse shape (See Eq. (92) in Ref. [9]):

```
jxgaussian[J0Xj_, tx_, t_] :=  
  J0Xj * Exp[-4 * Log[2] * (t / tx)^2] * HeavisideTheta[t + 2 * tx] * HeavisideTheta[2 * tx - t]
```

FWHM of the crater profile:

```
rhoXX := 2.2 * MicrometerToBohr  
rhoYY := 1.2 * MicrometerToBohr
```

Number of LCLS photons:

```
numXRays[pulseenergy_, photonenergy_] := pulseenergy / photonenergy
```

LCLS photon flux; fluence in photons / shot / Bohr²:

```

fluence[nxr_] := nxr / (Pi / Log[16] * rhoXX * rhoYY)

iondominant[pulseenergy_, photonenergy_, tauX_] := Module[{nxr, J0Xpeak, sol, jxdom},
  nxr = numXRays[pulseenergy, photonenergy];
  J0Xpeak = 2 * Sqrt[Log[2] / Pi] / tauX * fluence[nxr];
  jxdom[t_] := jxgaussian[J0Xpeak, tauX, t];
  sol = papaSolveRateEqs[photonenergy, jxdom, neginfinity, posinfinity, 1 / 1000, True];
  sol
]

```

Optimal fraction of nominal pulse energy determined with the fragmentation matrix model:

```

optdamp280 := 0.31
optdamp4 := 0.26

```

Detailed analysis for a pulse with nominal energy: 0.15 mJ, photon energy: 1100 eV, duration: 4 fs

```

sol =
  iondominant[optdamp4 * 0.15 * MilliJouleToHartree, 1100 * eVToHartree, 4 / t0ToFemtoSecond];


```

■ Neutral

```


Plot[{solGround[sol, t1 / t0ToFemtoSecond]},
 {t1, -10, 10}, PlotRange -> {-0.1, 1.1}, PlotStyle -> {{Thick, Red}}]

```




■ Singly

```
Plot[{solSingly[sol, t1 / t0ToFemtoSecond],
      (Replace[P213[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P222[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      Replace[P123[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}], {t1, -10, 25},
      PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}}]
```




■ Doubly

```
Plot[{solDoubly[sol, t1 / t0ToFemtoSecond],
      (Replace[P212[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P203[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P221[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      (Replace[P113[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P122[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      Replace[P023[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}], {t1, -10, 25},
      PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```




■ Triply

```
Plot[{solTriply[sol, t1 / t0ToFemtoSecond],
  (Replace[P202[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P211[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P220[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P103[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P112[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P121[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P013[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P022[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -10, 25},
  PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```




■ Quadruply

```
Plot[{solQuadruply[sol, t1 / t0ToFemtoSecond],
  (Replace[P201[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P210[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P102[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P111[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P120[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P003[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P012[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P021[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -10, 50},
  PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```




■ Quintuply

```
Plot[{solQuintuply[sol, t1 / t0ToFemtoSecond],
  Replace[P200[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond},
  (Replace[P101[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P110[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P011[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P020[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P002[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -10, 25},
  PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green},
  {Thick, Blue}, {Thick, Magenta}, {Thick, Orange}, {Thick, Cyan}}]
```




■ Sextuply

```
Plot[{solSextuply[sol, t1 / t0ToFemtoSecond],
  Replace[P100[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond},
  (Replace[P010[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P001[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -10, 25},
  PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Green}, {Thick, Blue}}]
```



■ Heptuply

```
Plot[{solSeptuply[sol, t1 / t0ToFemtoSecond]}, {t1, -100, 400}, PlotRange -> All, PlotStyle -> {Thick, Red}]
```




Detailed analysis for a pulse with nominal energy: 0.26 mJ, photon energy: 1100 eV, duration: 280 fs

```
sol = iondominant[optdamp280 * 0.26 * MilliJouleToHartree,
1100 * eVToHartree, 280 / t0ToFemtoSecond];
```


■ Neutral

```
Plot[{solGround[sol, t1 / t0ToFemtoSecond]}, {t1, -300, 300}, PlotRange -> {-0.1, 1.1}, PlotStyle -> {{Thick, Red}}]
```




■ Singly

```
Plot[{solSingly[sol, t1 / t0ToFemtoSecond],
      (Replace[P213[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P222[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      Replace[P123[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}], {t1, -300, 600},
      PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}}]
```




■ Doubly

```
Plot[{solDoubly[sol, t1 / t0ToFemtoSecond],
      (Replace[P212[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P203[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P221[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      (Replace[P113[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P122[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      Replace[P023[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}], {t1, -300, 600},
      PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```




■ Triply

```
Plot[{solTriply[sol, t1 / t0ToFemtoSecond],
  (Replace[P202[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P211[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P220[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P103[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P112[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P121[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P013[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P022[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -300, 600},
PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```




■ Quadruply

```
Plot[{solQuadruply[sol, t1 / t0ToFemtoSecond],
  (Replace[P201[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P210[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P102[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P111[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P120[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P003[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P012[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P021[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -300, 600},
PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```




■ Quintuply

```
Plot[{solQuintuply[sol, t1 / t0ToFemtoSecond],
  Replace[P200[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond},
  (Replace[P101[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P110[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
  (Replace[P011[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P020[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P002[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -300, 600},
  PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green},
  {Thick, Blue}, {Thick, Magenta}, {Thick, Orange}, {Thick, Cyan}}]
```




■ Sextuply

```
Plot[{solSextuply[sol, t1 / t0ToFemtoSecond],
  Replace[P100[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond},
  (Replace[P010[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
  (Replace[P001[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -300, 600},
  PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Green}, {Thick, Blue}}]
```



■ Septuply

```
Plot[{solSeptuply[sol, t1 / t0ToFemtoSecond]},  
{t1, -300, 600}, PlotRange -> All, PlotStyle -> {Thick, Red}]
```




Dominant path analysis for a nitrogen molecule with the nominal pulse energy: 0.15 mJ, photon energy: 1100 eV, duration: 4 fs

```
sol =  
  iondominant[optdamp4 * 0.15 * MilliJouleToHartree, 1100 * eVToHartree, 4 / t0ToFemtoSecond];
```


■ Neutral

```
Plot[{solGround[sol, t1 / t0ToFemtoSecond]^2},  
{t1, -10, 10}, PlotRange -> {-0.1, 1.1}, PlotStyle -> {{Thick, Red}}]
```




■ Singly

```
Plot[{2 * solGround[sol, t1 / t0ToFemtoSecond] * solSingly[sol, t1 / t0ToFemtoSecond],
      2 * solGround[sol, t1 / t0ToFemtoSecond] *
      (Replace[P123[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})},
      {t1, -10, 25}, PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}}]
```




■ Doubly

```
Plot[{solDoubly[sol, t1 / t0ToFemtoSecond],
      (Replace[P123[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})^2,
      2 * solGround[sol, t1 / t0ToFemtoSecond] *
      (Replace[P023[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond})}, {t1, -10, 25},
      PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```




■ Presentation graphics

```
Plot[{solNeutral[sol, t1 / t0ToFemtoSecond], solSCH[sol, t1 / t0ToFemtoSecond],
solDCHtwosites[sol, t1 / t0ToFemtoSecond], solDCHsinglesite[sol, t1 / t0ToFemtoSecond],
solPAP[sol, t1 / t0ToFemtoSecond]}, {t1, -5, 20}, PlotRange -> {{-6, 22}, {-0.1, 1.2}},
PlotStyle -> {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}, {Thick, Magenta}},
AspectRatio -> 1 / 1.3, AxesLabel -> {" $t$  [fs]", " $\tilde{P}(t)$ "},
AxesStyle -> {{Arrowheads[Automatic], AbsoluteThickness[0.8]},
{Arrowheads[Automatic], AbsoluteThickness[0.8]}},
LabelStyle -> {FontFamily -> "Arial", FontSlant -> Plain, FontSize -> 18}]
```



```
LogPlot[{solNeutral[sol, t1 / t0ToFemtoSecond], solSCH[sol, t1 / t0ToFemtoSecond],
solDCHtwosites[sol, t1 / t0ToFemtoSecond], solDCHsinglesite[sol, t1 / t0ToFemtoSecond],
solPAP[sol, t1 / t0ToFemtoSecond]}, {t1, -10, 20}, PlotRange -> {{-6, 23}, {5^-5, 5}},
PlotStyle -> {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}, {Thick, Magenta}},
AspectRatio -> 1 / 1.3, AxesLabel -> {" $t$  [fs]", " $\tilde{P}(t)$ "},
AxesStyle -> {{Arrowheads[Automatic], AbsoluteThickness[0.8]},
{Arrowheads[Automatic], AbsoluteThickness[0.8]}},
LabelStyle -> {FontFamily -> "Helvetica", FontSlant -> Plain, FontSize -> 17}]
```




Dominant path analysis for a nitrogen molecule with the nominal pulse energy: 0.26 mJ, photon energy: 1100 eV, duration: 280 fs

```
sol = iondominant[optdamp280 * 0.26 * MilliJouleToHartree,
 1100 * eVToHartree, 280 / t0ToFemtoSecond];
```


■ Neutral

```
Plot[{solGround[sol, t1 / t0ToFemtoSecond]}, {t1, -300, 300}, PlotRange -> {-0.1, 1.1}, PlotStyle -> {{Thick, Red}}]
```




■ Singly

```
Plot[{solsSingly[sol, t1 / t0ToFemtoSecond],
(Replace[P213[t], sol[[1]]] /. {t -> t1 / t0ToFemtoSecond}) +
(Replace[P222[t], sol[[1]]] /. {t -> t1 / t0ToFemtoSecond}),
Replace[P123[t], sol[[1]]] /. {t -> t1 / t0ToFemtoSecond}], {t1, -300, 600},
PlotRange -> All, PlotStyle -> {{Thick, Black}, {Thick, Red}, {Thick, Green}}]
```




■ Doubly

```
Plot[{solDoubly[sol, t1 / t0ToFemtoSecond],
      (Replace[P212[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P203[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P221[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      (Replace[P113[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}) +
      (Replace[P122[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}),
      Replace[P023[t], sol[[1]]] /. {t → t1 / t0ToFemtoSecond}], {t1, -300, 600},
PlotRange → All, PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}}]
```



■ Presentation graphics


```
Plot[{solNeutral[sol, t1 / t0ToFemtoSecond],
      solSCH[sol, t1 / t0ToFemtoSecond], soldCHtwosites[sol, t1 / t0ToFemtoSecond],
      soldCHsinglesite[sol, t1 / t0ToFemtoSecond], solPAP[sol, t1 / t0ToFemtoSecond}],
     {t1, -300, 300}, PlotRange → {{-350, 350}, {-0.1, 1.2}},
     PlotStyle → {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}, {Thick, Magenta}},
     AspectRatio → 1 / 1.3, AxesLabel → {"t [fs]", "P̃(t)" },
     AxesStyle → {{Arrowheads[Automatic], AbsoluteThickness[0.8]},
                   {Arrowheads[Automatic], AbsoluteThickness[0.8]}},
     LabelStyle → {FontFamily → "Arial", FontSlant → Plain, FontSize → 18}]
```



```

LogPlot[{solNeutral[sol, t1 / t0ToFemtoSecond],
  solSCH[sol, t1 / t0ToFemtoSecond], solDCHtwosites[sol, t1 / t0ToFemtoSecond],
  solDCHsinglesite[sol, t1 / t0ToFemtoSecond], solPAP[sol, t1 / t0ToFemtoSecond]}, {
  t1, -300, 330}, PlotRange -> {{-300, 400}, {5^-7, 30}}, 
  PlotStyle -> {{Thick, Black}, {Thick, Red}, {Thick, Green}, {Thick, Blue}, {Thick, Magenta}}, 
  AspectRatio -> 1 / 1.3, AxesLabel -> {"t [fs]", "P(t)"}, 
  AxesStyle -> {{Arrowheads[Automatic], AbsoluteThickness[0.8]}, 
    {Arrowheads[Automatic], AbsoluteThickness[0.8]}}, 
  LabelStyle -> {FontFamily -> "Arial", FontSlant -> Plain, FontSize -> 17}]

```



References

- [1] Nina Rohringer, Robin Santra, Phys. Rev. A **76**, 033416 (2007), <http://dx.doi.org/10.1103/PhysRevA.76.033416>
- [2] Los Alamos National Laboratory Atomic Physics Codes, <http://aphysics2.lanl.gov/cgi-bin/ION/runlanl08a.pl>
- [3] LCLS Parameter Index for X-RAY-OPTICS / RADIATION-SOURCE, http://www-srsl.slac.stanford.edu/htbin/rdb-web/LCLS_params_DB_public
- [4] M. G. Makris, P. Lambropoulos, A. Mihelič, Phys. Rev. Lett. **102**, 033002 (2009), <http://dx.doi.org/10.1103/PhysRevLett.102.033002>
- [5] J. L. Campbell, Tibor Papp, *Widths of the atomic K-N7 levels*, At. Data Nucl. Data Tables **77**, 1–56 (2001), <http://dx.doi.org/10.1006/adnd.2000.0848>
- [6] Ryan Coffee, John Bozek, Christian Buth, Marcus Hertlein, Markus Gühr, Nora Berrah, Ali Belkacem, Linda Young, Louis DiMauro, Hamed Merdji, Janos Hajdu, Philip H. Bucksbaum, *Xray multiple ionization of impulsively aligned molecules*, Linac Coherent Light Source, Menlo Park, California, USA, 2009, research proposal for the first round of beam time at the Linac Coherent Light Source
- [7] Mau Hsiung Chen, Phys. Rev. A **44**, 239–242 (1991), <http://dx.doi.org/10.1103/PhysRevA.44.239>
- [8] Albert C. Thompson *et al.*, *X-ray data booklet*, Center for X-ray Optics and Advanced Light Source, Lawrence Berkeley National Laboratory, <http://xdb.lbl.gov>
- [9] Christian Buth, Robin Santra, Phys. Rev. A **78**, 043409 (2008), <http://dx.doi.org/10.1103/PhysRevA.78.043409>