

Daya Bay Reactor Neutrino Experiment

Supplemental Material

August 2018

This document provides supplemental information for the article titled “Search for a time-varying electron antineutrino signal at Daya Bay”. Section I provides the correlation matrices associated with the fit of the Standard-Model Extension presented in the article. All other sections provide information that is necessary to reproduce the fit.

CONTENTS

I. LV-CPTV COEFFICIENT CORRELATION MATRICES	2
II. RESIDUAL SURVIVAL PROBABILITY VS. SIDEREAL TIME	4
III. ANTINEUTRINO ENERGIES	6
IV. DIRECTIONAL FACTORS	6
V. PREDICTED REACTOR ANTINEUTRINO EVENT FRACTIONS	7
VI. BASELINES, TRANSITION AMPLITUDES, AND EXPERIMENTAL FACTORS	9

I. LV-CPTV COEFFICIENT CORRELATION MATRICES

This section provides the correlation matrices associated with the fits of the Standard-Model Extension described in Section IV and presented in Table II of the main text. Tables I through VI gives the matrices for each of the six flavor pairs.

TABLE I. Correlation matrix for the $e\bar{e}$ flavor pair.

a_R^X	1	0.77	0.16	-0.01	-0.02	-0.05	-0.00	-0.00
c_R^{TX}	0.77	1	0.73	0.01	-0.00	-0.12	-0.00	0.00
c_R^{XZ}	0.16	0.73	1	0.05	0.12	0.00	0.00	0.00
a_R^Y	-0.01	0.01	0.05	1	0.77	0.16	0.00	0.00
c_R^{TY}	-0.02	-0.00	0.12	0.77	1	0.73	0.00	0.00
c_R^{YZ}	-0.05	-0.12	0.00	0.16	0.73	1	-0.00	0.00
$c_R^{XX} - c_R^{YY}$	-0.00	-0.00	0.00	0.00	0.00	-0.00	1	-0.00
c_R^{XY}	-0.00	0.00	0.00	0.00	0.00	0.00	-0.00	1

TABLE II. Correlation matrix for the $\mu\bar{\mu}$ flavor pair.

a_R^X	1	0.76	0.15	-0.00	-0.02	-0.04	-0.00	0.00
c_R^{TX}	0.76	1	0.74	0.01	-0.00	-0.12	-0.00	0.00
c_R^{XZ}	0.15	0.74	1	0.04	0.12	0.00	0.00	0.00
a_R^Y	-0.00	0.01	0.04	1	0.76	0.15	0.00	0.00
c_R^{TY}	-0.02	-0.00	0.12	0.76	1	0.73	0.00	0.00
c_R^{YZ}	-0.04	-0.12	0.00	0.15	0.73	1	-0.00	0.00
$c_R^{XX} - c_R^{YY}$	-0.00	-0.00	0.00	0.00	0.00	-0.00	1	-0.00
c_R^{XY}	0.00	0.00	0.00	0.00	0.00	0.00	-0.00	1

TABLE III. Correlation matrix for the $\tau\bar{\tau}$ flavor pair.

a_R^X	1	0.78	0.18	-0.00	-0.02	-0.05	-0.00	-0.00
c_R^{TX}	0.78	1	0.73	0.02	-0.00	-0.12	-0.00	0.00
c_R^{XZ}	0.18	0.73	1	0.05	0.12	0.00	0.00	0.00
a_R^Y	-0.00	0.02	0.05	1	0.78	0.18	0.00	0.00
c_R^{TY}	-0.02	-0.00	0.12	0.78	1	0.73	0.00	0.00
c_R^{YZ}	-0.05	-0.12	0.00	0.18	0.73	1	-0.00	0.00
$c_R^{XX} - c_R^{YY}$	-0.00	-0.00	0.00	0.00	0.00	-0.00	1	-0.00
c_R^{XY}	-0.00	0.00	0.00	0.00	0.00	0.00	-0.00	1

TABLE IV. Correlation matrix for the $e\bar{\mu}$ flavor pair.

a_R^X	1	0.88	0.10	-0.00	0.02	0.04	0.00	-0.00
c_R^{TX}	0.88	1	0.50	-0.03	-0.00	-0.05	0.00	-0.00
c_R^{XZ}	0.10	0.50	1	-0.04	0.05	-0.00	0.00	0.00
a_R^Y	-0.00	-0.03	-0.04	1	0.88	0.10	-0.01	0.00
c_R^{TY}	0.02	-0.00	0.05	0.88	1	0.50	-0.00	0.00
c_R^{YZ}	0.04	-0.05	-0.00	0.10	0.50	1	-0.00	0.00
$c_R^{XX} - c_R^{YY}$	0.00	0.00	0.00	-0.01	-0.00	-0.00	1	-0.00
c_R^{XY}	-0.00	-0.00	0.00	0.00	0.00	0.00	-0.00	1

TABLE V. Correlation matrix for the $e\tau$ flavor pair.

a_R^X	1	0.90	0.14	-0.00	0.03	0.05	0.00	-0.00
c_R^{TX}	0.90	1	0.49	-0.03	-0.00	-0.03	0.00	-0.00
c_R^{XZ}	0.14	0.49	1	-0.05	0.03	-0.00	0.00	0.00
a_R^Y	-0.00	-0.03	-0.05	1	0.90	0.14	-0.01	0.00
c_R^{TY}	0.03	-0.00	0.03	0.90	1	0.49	-0.01	0.00
c_R^{YZ}	0.05	-0.03	-0.00	0.14	0.49	1	-0.00	-0.00
$c_R^{XX} - c_R^{YY}$	0.00	0.00	0.00	-0.01	-0.01	-0.00	1	-0.00
c_R^{XY}	-0.00	-0.00	0.00	0.00	0.00	-0.00	-0.00	1

TABLE VI. Correlation matrix for the $\mu\tau$ flavor pair.

a_R^X	1	0.77	0.16	-0.00	-0.02	-0.05	-0.00	0.00
c_R^{TX}	0.77	1	0.73	0.01	-0.00	-0.12	-0.00	0.00
c_R^{XZ}	0.16	0.73	1	0.05	0.12	0.00	0.00	0.00
a_R^Y	-0.00	0.01	0.05	1	0.77	0.16	0.00	0.00
c_R^{TY}	-0.02	-0.00	0.12	0.77	1	0.73	0.00	0.00
c_R^{YZ}	-0.05	-0.12	0.00	0.16	0.73	1	-0.00	0.00
$c_R^{XX} - c_R^{YY}$	-0.00	-0.00	0.00	0.00	0.00	-0.00	1	-0.00
c_R^{XY}	0.00	0.00	0.00	0.00	0.00	0.00	-0.00	1

II. RESIDUAL SURVIVAL PROBABILITY VS. SIDEREAL TIME

Tables VII, VIII, and IX provide the residual survival probability of reactor $\bar{\nu}_e$, R , in 24 sidereal hour bins and 5 energy bins, for each experimental hall. These are the same R shown in Fig. 5 and used in Eq. (5) of the main text. The energy bins are defined in Table X. The errors include both statistical and systematic components.

TABLE VII. Residual survival probability of $\bar{\nu}_e$ in **experimental hall 1** vs. sidereal hour and energy bin.

Time bin	E bin 1	E bin 2	E bin 3	E bin 4	E bin 5
1	-0.0033 ± 0.0160	-0.0060 ± 0.0122	-0.0010 ± 0.0123	0.0033 ± 0.0147	-0.0058 ± 0.0166
2	-0.0027 ± 0.0161	-0.0003 ± 0.0122	0.0049 ± 0.0122	0.0124 ± 0.0146	0.0133 ± 0.0165
3	0.0085 ± 0.0160	-0.0027 ± 0.0122	0.0077 ± 0.0122	-0.0176 ± 0.0149	0.0016 ± 0.0166
4	-0.0138 ± 0.0162	0.0237 ± 0.0121	-0.0118 ± 0.0124	-0.0174 ± 0.0149	0.0003 ± 0.0166
5	-0.0082 ± 0.0162	0.0110 ± 0.0122	-0.0024 ± 0.0123	-0.0097 ± 0.0149	0.0278 ± 0.0164
6	-0.0084 ± 0.0162	0.0092 ± 0.0122	0.0051 ± 0.0123	0.0124 ± 0.0148	0.0170 ± 0.0166
7	-0.0000 ± 0.0162	0.0002 ± 0.0123	-0.0109 ± 0.0125	-0.0131 ± 0.0150	-0.0068 ± 0.0168
8	0.0145 ± 0.0162	0.0062 ± 0.0123	-0.0069 ± 0.0125	-0.0006 ± 0.0150	0.0252 ± 0.0166
9	0.0212 ± 0.0161	0.0150 ± 0.0123	0.0074 ± 0.0124	0.0226 ± 0.0148	0.0122 ± 0.0167
10	0.0029 ± 0.0163	0.0011 ± 0.0124	0.0015 ± 0.0125	-0.0005 ± 0.0150	-0.0055 ± 0.0169
11	-0.0028 ± 0.0163	-0.0017 ± 0.0123	0.0043 ± 0.0124	-0.0221 ± 0.0151	0.0103 ± 0.0167
12	0.0041 ± 0.0161	-0.0244 ± 0.0124	0.0055 ± 0.0123	-0.0072 ± 0.0149	0.0044 ± 0.0167
13	0.0162 ± 0.0160	0.0035 ± 0.0122	-0.0099 ± 0.0124	0.0004 ± 0.0148	-0.0115 ± 0.0168
14	-0.0099 ± 0.0162	0.0043 ± 0.0122	-0.0128 ± 0.0124	0.0177 ± 0.0147	-0.0326 ± 0.0170
15	0.0136 ± 0.0160	0.0001 ± 0.0122	0.0039 ± 0.0123	0.0145 ± 0.0147	0.0182 ± 0.0165
16	0.0097 ± 0.0160	0.0038 ± 0.0122	0.0117 ± 0.0122	-0.0031 ± 0.0148	0.0120 ± 0.0165
17	0.0114 ± 0.0160	-0.0132 ± 0.0123	0.0064 ± 0.0123	-0.0373 ± 0.0151	-0.0046 ± 0.0167
18	-0.0241 ± 0.0163	-0.0002 ± 0.0122	-0.0029 ± 0.0123	0.0134 ± 0.0147	-0.0258 ± 0.0168
19	-0.0266 ± 0.0163	0.0084 ± 0.0122	-0.0038 ± 0.0123	0.0155 ± 0.0147	-0.0126 ± 0.0167
20	0.0079 ± 0.0160	-0.0001 ± 0.0122	0.0041 ± 0.0123	-0.0004 ± 0.0147	-0.0287 ± 0.0168
21	-0.0010 ± 0.0160	-0.0117 ± 0.0123	-0.0075 ± 0.0123	-0.0051 ± 0.0148	-0.0088 ± 0.0167
22	-0.0082 ± 0.0161	-0.0142 ± 0.0123	0.0013 ± 0.0123	0.0002 ± 0.0147	-0.0151 ± 0.0167
23	-0.0157 ± 0.0162	-0.0113 ± 0.0122	0.0111 ± 0.0122	-0.0029 ± 0.0147	0.0074 ± 0.0165
24	0.0215 ± 0.0158	0.0030 ± 0.0121	-0.0016 ± 0.0122	0.0292 ± 0.0145	0.0180 ± 0.0164

TABLE VIII. Residual survival probability of $\bar{\nu}_e$ in **experimental hall 2** vs. sidereal hour and energy bin.

Time bin	E bin 1	E bin 2	E bin 3	E bin 4	E bin 5
1	-0.0004 ± 0.0182	-0.0170 ± 0.0139	0.0054 ± 0.0139	0.0171 ± 0.0167	-0.0178 ± 0.0189
2	0.0225 ± 0.0180	0.0013 ± 0.0137	0.0000 ± 0.0139	0.0074 ± 0.0167	-0.0308 ± 0.0190
3	-0.0061 ± 0.0182	-0.0168 ± 0.0138	0.0103 ± 0.0138	0.0027 ± 0.0167	-0.0089 ± 0.0188
4	0.0097 ± 0.0181	-0.0012 ± 0.0137	-0.0081 ± 0.0139	-0.0058 ± 0.0168	0.0133 ± 0.0186
5	-0.0255 ± 0.0185	-0.0140 ± 0.0139	-0.0050 ± 0.0139	-0.0083 ± 0.0169	0.0119 ± 0.0186
6	-0.0028 ± 0.0182	-0.0047 ± 0.0138	0.0221 ± 0.0137	-0.0349 ± 0.0171	0.0152 ± 0.0186
7	-0.0106 ± 0.0184	-0.0093 ± 0.0139	-0.0160 ± 0.0140	0.0078 ± 0.0168	0.0231 ± 0.0186
8	-0.0320 ± 0.0186	-0.0011 ± 0.0138	-0.0114 ± 0.0140	0.0140 ± 0.0167	0.0207 ± 0.0186
9	-0.0169 ± 0.0184	-0.0152 ± 0.0139	-0.0174 ± 0.0140	0.0162 ± 0.0167	0.0085 ± 0.0187
10	-0.0264 ± 0.0185	0.0173 ± 0.0137	-0.0045 ± 0.0139	-0.0013 ± 0.0168	-0.0149 ± 0.0189
11	0.0234 ± 0.0180	-0.0099 ± 0.0138	0.0141 ± 0.0138	-0.0121 ± 0.0169	0.0264 ± 0.0185
12	0.0165 ± 0.0180	-0.0087 ± 0.0138	0.0098 ± 0.0138	0.0153 ± 0.0166	0.0279 ± 0.0184
13	0.0222 ± 0.0179	0.0100 ± 0.0136	0.0144 ± 0.0137	-0.0227 ± 0.0169	-0.0014 ± 0.0186
14	-0.0151 ± 0.0183	0.0102 ± 0.0136	-0.0052 ± 0.0139	0.0142 ± 0.0166	-0.0093 ± 0.0188
15	-0.0348 ± 0.0185	0.0227 ± 0.0136	-0.0101 ± 0.0139	0.0123 ± 0.0166	-0.0475 ± 0.0191
16	0.0318 ± 0.0179	-0.0071 ± 0.0138	-0.0119 ± 0.0139	0.0057 ± 0.0167	-0.0167 ± 0.0188
17	0.0052 ± 0.0182	0.0196 ± 0.0136	0.0146 ± 0.0138	-0.0215 ± 0.0170	0.0291 ± 0.0185
18	-0.0010 ± 0.0182	-0.0011 ± 0.0138	-0.0209 ± 0.0140	-0.0166 ± 0.0169	0.0217 ± 0.0185
19	-0.0121 ± 0.0184	0.0033 ± 0.0137	-0.0023 ± 0.0139	0.0118 ± 0.0167	-0.0063 ± 0.0188
20	-0.0029 ± 0.0182	-0.0045 ± 0.0138	0.0043 ± 0.0138	0.0092 ± 0.0167	0.0094 ± 0.0186
21	0.0404 ± 0.0178	0.0153 ± 0.0136	0.0203 ± 0.0137	-0.0117 ± 0.0168	-0.0041 ± 0.0187
22	0.0019 ± 0.0181	0.0366 ± 0.0134	0.0174 ± 0.0137	-0.0046 ± 0.0167	-0.0242 ± 0.0189
23	0.0147 ± 0.0180	-0.0042 ± 0.0137	-0.0171 ± 0.0139	0.0022 ± 0.0167	-0.0378 ± 0.0190
24	0.0035 ± 0.0182	-0.0180 ± 0.0139	0.0013 ± 0.0139	0.0104 ± 0.0167	0.0198 ± 0.0185

TABLE IX. Residual survival probability of $\bar{\nu}_e$ in **experimental hall 3** vs. sidereal hour and energy bin.

Time bin	E bin 1	E bin 2	E bin 3	E bin 4	E bin 5
1	-0.0342 ± 0.0333	0.0037 ± 0.0233	0.0278 ± 0.0230	-0.0254 ± 0.0287	-0.0449 ± 0.0325
2	-0.0181 ± 0.0329	0.0040 ± 0.0233	0.0268 ± 0.0229	0.0450 ± 0.0276	0.0007 ± 0.0317
3	0.0085 ± 0.0323	-0.0252 ± 0.0235	0.0586 ± 0.0225	-0.0202 ± 0.0285	0.0021 ± 0.0316
4	0.0174 ± 0.0322	0.0405 ± 0.0227	-0.0369 ± 0.0237	0.0022 ± 0.0282	-0.0263 ± 0.0321
5	-0.0439 ± 0.0334	-0.0271 ± 0.0237	-0.0140 ± 0.0235	0.0182 ± 0.0281	0.0696 ± 0.0306
6	-0.0007 ± 0.0327	-0.0044 ± 0.0234	-0.0149 ± 0.0235	0.0320 ± 0.0279	-0.0267 ± 0.0322
7	0.0275 ± 0.0322	-0.0304 ± 0.0238	-0.0085 ± 0.0235	0.0027 ± 0.0283	-0.0050 ± 0.0319
8	-0.0036 ± 0.0329	0.0187 ± 0.0233	-0.0176 ± 0.0238	0.0136 ± 0.0283	0.0203 ± 0.0317
9	0.0113 ± 0.0325	-0.0066 ± 0.0235	-0.0008 ± 0.0234	0.0180 ± 0.0282	0.0024 ± 0.0319
10	0.0252 ± 0.0323	0.0409 ± 0.0229	0.0236 ± 0.0231	-0.0293 ± 0.0289	0.0218 ± 0.0315
11	0.0098 ± 0.0326	0.0029 ± 0.0233	-0.0093 ± 0.0235	0.0377 ± 0.0278	0.0042 ± 0.0317
12	-0.0250 ± 0.0331	-0.0033 ± 0.0234	0.0225 ± 0.0231	0.0227 ± 0.0280	-0.0014 ± 0.0318
13	-0.0036 ± 0.0325	-0.0121 ± 0.0234	-0.0195 ± 0.0235	0.0027 ± 0.0282	-0.0175 ± 0.0319
14	0.0305 ± 0.0320	-0.0272 ± 0.0237	-0.0099 ± 0.0234	-0.0522 ± 0.0291	0.0715 ± 0.0305
15	-0.0511 ± 0.0335	-0.0172 ± 0.0235	-0.0486 ± 0.0239	-0.0517 ± 0.0291	0.0490 ± 0.0308
16	0.0500 ± 0.0316	-0.0047 ± 0.0233	0.0097 ± 0.0231	-0.0026 ± 0.0283	0.0117 ± 0.0314
17	-0.0266 ± 0.0332	0.0408 ± 0.0228	-0.0264 ± 0.0236	0.0060 ± 0.0282	0.0128 ± 0.0315
18	-0.0204 ± 0.0330	0.0189 ± 0.0232	0.0216 ± 0.0231	-0.0147 ± 0.0286	-0.0483 ± 0.0327
19	0.0567 ± 0.0318	-0.0100 ± 0.0235	0.0166 ± 0.0232	0.0154 ± 0.0282	-0.0150 ± 0.0321
20	0.0125 ± 0.0323	-0.0119 ± 0.0234	0.0158 ± 0.0230	0.0383 ± 0.0277	-0.0147 ± 0.0318
21	0.0031 ± 0.0325	0.0452 ± 0.0227	0.0380 ± 0.0228	0.0038 ± 0.0283	-0.0139 ± 0.0320
22	0.0136 ± 0.0322	-0.0228 ± 0.0235	-0.0363 ± 0.0236	-0.0295 ± 0.0286	-0.0069 ± 0.0316
23	-0.0094 ± 0.0326	0.0331 ± 0.0228	-0.0024 ± 0.0232	-0.0043 ± 0.0283	0.0076 ± 0.0315
24	0.0561 ± 0.0317	0.0001 ± 0.0233	0.0262 ± 0.0229	0.0249 ± 0.0279	0.0093 ± 0.0316

III. ANTINEUTRINO ENERGIES

Table X lists the mean $\bar{\nu}_e$ energy corresponding to the selected prompt energy bins introduced in Section IV and illustrated in Fig. 5 of the main text. The root mean square (RMS) of each mean $\bar{\nu}_e$ energy is also listed.

TABLE X. Prompt energy bins and corresponding mean $\bar{\nu}_e$ energies.

E Bin	$E_{\text{prompt}}[\text{MeV}]$	$\bar{E}_{\bar{\nu}_e}[\text{MeV}]$	$\text{RMS}_{\bar{E}_{\bar{\nu}_e}}[\text{MeV}]$
1	(0.7, 2.0)	2.55	0.53
2	(2.0, 3.0)	3.36	0.37
3	(3.0, 4.0)	4.30	0.35
4	(4.0, 5.0)	5.28	0.35
5	(5.0, 12.0)	6.57	0.63

IV. DIRECTIONAL FACTORS

Table XI lists the directional factors for each reactor-hall pair. They were calculated following Eq. (A9) of the main text.

TABLE XI. Directional factors for each reactor-hall pair.

Hall	Reactor	\hat{N}^X	\hat{N}^Y	\hat{N}^Z
EH1	D1	-0.461	-0.004	0.887
EH1	D2	-0.448	0.236	0.862
EH1	L1	0.221	-0.753	-0.620
EH1	L2	0.210	-0.769	-0.604
EH1	L3	0.265	-0.668	-0.696
EH1	L4	0.261	-0.673	-0.692
EH2	D1	-0.390	0.263	0.883
EH2	D2	-0.382	0.323	0.866
EH2	L1	-0.329	-0.698	0.636
EH2	L2	-0.370	-0.560	0.741
EH2	L3	0.008	-0.989	-0.149
EH2	L4	-0.046	-0.998	-0.034
EH3	D1	-0.379	-0.303	0.874
EH3	D2	-0.384	-0.260	0.886
EH3	L1	-0.236	-0.820	0.521
EH3	L2	-0.254	-0.786	0.564
EH3	L3	-0.131	-0.954	0.269
EH3	L4	-0.151	-0.936	0.317

V. PREDICTED REACTOR ANTINEUTRINO EVENT FRACTIONS

Tables [XII](#), [XIII](#), and [XIV](#) provide the predicted fractions of inverse beta decays, f , in each experimental hall (EH) and each sidereal time bin, originating from each nuclear reactor as defined by Eq. (4) in Section IV of the main text. See Fig. 1 of the main text for the relative location of each reactor.

TABLE XII. Predicted fractions of inverse beta decays in **experimental hall 1** and time bin, originating from each nuclear reactor.

Time bin	D1	D2	L1	L2	L3	L4
1	0.389	0.405	0.063	0.082	0.030	0.031
2	0.389	0.405	0.063	0.082	0.030	0.031
3	0.387	0.406	0.063	0.083	0.030	0.031
4	0.387	0.406	0.063	0.083	0.030	0.031
5	0.387	0.406	0.063	0.083	0.030	0.031
6	0.386	0.406	0.064	0.083	0.030	0.031
7	0.386	0.406	0.064	0.083	0.030	0.031
8	0.386	0.406	0.064	0.083	0.030	0.030
9	0.385	0.407	0.064	0.083	0.030	0.031
10	0.384	0.408	0.064	0.083	0.030	0.031
11	0.385	0.408	0.063	0.082	0.030	0.031
12	0.386	0.408	0.063	0.082	0.030	0.031
13	0.387	0.408	0.063	0.082	0.030	0.031
14	0.388	0.407	0.062	0.081	0.030	0.031
15	0.389	0.406	0.062	0.081	0.030	0.031
16	0.389	0.406	0.062	0.082	0.030	0.031
17	0.389	0.406	0.062	0.082	0.030	0.031
18	0.388	0.407	0.062	0.082	0.030	0.031
19	0.388	0.407	0.062	0.082	0.030	0.031
20	0.387	0.407	0.063	0.082	0.030	0.031
21	0.387	0.406	0.063	0.082	0.030	0.031
22	0.387	0.407	0.063	0.082	0.030	0.031
23	0.387	0.407	0.063	0.082	0.030	0.031
24	0.388	0.407	0.063	0.082	0.030	0.031

TABLE XIII. Predicted fractions of inverse beta decays in **experimental hall 2** and time bin, originating from each nuclear reactor.

Time bin	D1	D2	L1	L2	L3	L4
1	0.033	0.033	0.257	0.254	0.197	0.227
2	0.033	0.033	0.258	0.254	0.196	0.227
3	0.033	0.033	0.258	0.254	0.196	0.226
4	0.032	0.033	0.257	0.257	0.196	0.226
5	0.032	0.033	0.257	0.256	0.196	0.226
6	0.032	0.033	0.259	0.255	0.196	0.226
7	0.032	0.033	0.260	0.255	0.195	0.225
8	0.032	0.033	0.261	0.254	0.196	0.224
9	0.032	0.033	0.261	0.254	0.196	0.225
10	0.032	0.033	0.260	0.254	0.195	0.227
11	0.032	0.033	0.259	0.253	0.195	0.228
12	0.032	0.033	0.259	0.253	0.196	0.227
13	0.032	0.033	0.258	0.253	0.196	0.227
14	0.033	0.033	0.257	0.253	0.197	0.228
15	0.032	0.033	0.256	0.253	0.197	0.228
16	0.032	0.033	0.256	0.254	0.196	0.228
17	0.033	0.033	0.256	0.254	0.196	0.228
18	0.032	0.033	0.256	0.254	0.196	0.228
19	0.032	0.033	0.256	0.254	0.196	0.229
20	0.032	0.033	0.256	0.254	0.196	0.229
21	0.032	0.033	0.256	0.253	0.196	0.229
22	0.032	0.033	0.257	0.253	0.196	0.229
23	0.032	0.033	0.257	0.253	0.196	0.228
24	0.033	0.033	0.257	0.254	0.196	0.228

TABLE XIV. Predicted fractions of inverse beta decays in **experimental hall 3** and time bin, originating from each nuclear reactor.

Time bin	D1	D2	L1	L2	L3	L4
1	0.118	0.130	0.183	0.197	0.193	0.180
2	0.118	0.129	0.183	0.197	0.193	0.180
3	0.117	0.129	0.184	0.197	0.193	0.180
4	0.116	0.129	0.183	0.199	0.193	0.179
5	0.117	0.129	0.183	0.199	0.193	0.180
6	0.116	0.129	0.185	0.198	0.193	0.179
7	0.116	0.129	0.186	0.198	0.193	0.179
8	0.116	0.129	0.187	0.197	0.193	0.178
9	0.115	0.129	0.187	0.197	0.193	0.179
10	0.115	0.129	0.186	0.197	0.192	0.181
11	0.115	0.130	0.185	0.196	0.192	0.181
12	0.116	0.130	0.184	0.197	0.192	0.181
13	0.117	0.130	0.184	0.196	0.193	0.181
14	0.117	0.131	0.183	0.195	0.193	0.181
15	0.117	0.130	0.182	0.196	0.193	0.182
16	0.117	0.130	0.182	0.196	0.193	0.182
17	0.117	0.130	0.182	0.196	0.193	0.182
18	0.117	0.130	0.182	0.196	0.193	0.182
19	0.117	0.130	0.182	0.196	0.192	0.182
20	0.117	0.130	0.183	0.196	0.192	0.182
21	0.117	0.130	0.182	0.197	0.193	0.182
22	0.116	0.130	0.182	0.197	0.193	0.182
23	0.117	0.130	0.182	0.197	0.193	0.182
24	0.117	0.130	0.182	0.197	0.193	0.181

VI. BASELINES, TRANSITION AMPLITUDES, AND EXPERIMENTAL FACTORS

Table XV lists the baselines L , and transition amplitudes $S_{ee}^{(0)}$ with experimental factors $\mathcal{M}_{ee}^{(1)}$, for each flavor pair and each reactor-hall pair (see Fig. 1 of the main text for the layout). The calculation of the latter two quantities followed Eqs. (A3) and (A5) of the main text. Neutrino oscillation parameters are from PDG 2015 ($\sin^2 2\theta_{12} = 0.846 \pm 0.021$, $\sin^2 2\theta_{23} = 0.999_{-0.018}^{+0.001}$, $\sin^2 2\theta_{13} = 0.085 \pm 0.005$, $\Delta m_{21}^2 = (7.53 \pm 0.18) \times 10^{-5} \text{eV}^2$, $\Delta m_{32}^2 = (2.42 \pm 0.06) \times 10^{-3} \text{eV}^2$) assuming normal neutrino mass ordering. We took the CP violating phase to be zero. The average antineutrino energy is 4.2 MeV.

TABLE XV. The baselines, transition amplitudes, and experimental factors for each flavor pair and each reactor-hall pair.

Hall	Reactor	$L[\text{m}]$	$ S_{ee}^{(0)} ^2$	$\text{Im}(S_{ee}^{(0)*}(\mathcal{M}_{ee}^{(1)})_{cd})$					
				ee	$\mu\mu$	$\tau\tau$	$e\mu$	$e\tau$	$\mu\tau$
EH1	D1	360.2	0.994	0.0146	-0.0054	-0.0092	0.0182	0.0279	-0.0138
EH1	D2	370.1	0.994	0.0150	-0.0055	-0.0095	0.0187	0.0286	-0.0141
EH1	L1	903.4	0.967	0.0325	-0.0122	-0.0203	0.0408	0.0614	-0.0271
EH1	L2	817.0	0.972	0.0302	-0.0113	-0.0189	0.0378	0.0571	-0.0259
EH1	L3	1353.9	0.938	0.0407	-0.0157	-0.0250	0.0513	0.0756	-0.0268
EH1	L4	1265.6	0.944	0.0397	-0.0152	-0.0244	0.0499	0.0739	-0.0278
EH2	D1	1335.0	0.940	0.0405	-0.0156	-0.0249	0.0510	0.0753	-0.0271
EH2	D2	1360.5	0.938	0.0408	-0.0158	-0.0250	0.0513	0.0757	-0.0267
EH2	L1	470.3	0.990	0.0188	-0.0069	-0.0118	0.0234	0.0358	-0.0175
EH2	L2	492.5	0.989	0.0196	-0.0073	-0.0123	0.0245	0.0373	-0.0182
EH2	L3	558.1	0.986	0.0219	-0.0081	-0.0138	0.0274	0.0417	-0.0201
EH2	L4	500.1	0.989	0.0199	-0.0074	-0.0125	0.0248	0.0378	-0.0184
EH3	D1	1921.4	0.915	0.0418	-0.0170	-0.0249	0.0509	0.0737	-0.0123
EH3	D2	1895.9	0.916	0.0420	-0.0170	-0.0250	0.0513	0.0742	-0.0132
EH3	L1	1536.9	0.929	0.0422	-0.0165	-0.0256	0.0528	0.0773	-0.0236
EH3	L2	1537.2	0.929	0.0422	-0.0165	-0.0256	0.0528	0.0773	-0.0236
EH3	L3	1555.6	0.928	0.0422	-0.0166	-0.0257	0.0529	0.0773	-0.0232
EH3	L4	1529.1	0.929	0.0421	-0.0165	-0.0256	0.0528	0.0773	-0.0238