

## Replacing Arcane Words with Common Synonyms

PMID	SID	Source	Target
32365314	6	Intervention: The intervention group used a nighttime orthosis on the second or third finger of the dominant hand.	The <i>treatment</i> group used a nighttime <i>brace</i> on the second or third finger of the dominant hand.

## Changing Passive Voice to Active Voice

PMID	SID	Source	Target
29857264	6	A comprehensive search was conducted on PubMed and Google Scholar.	We searched common online resources for papers.

## Omitting Subheadings

PMID	SID	Source	Target
15228825	1	Objective: To evaluate management of foreign bodies in the upper gastrointestinal tract.	Our objective is to rate treatment of foreign objects stuck in the upper digestive tract.

## Splitting Sentences

PMID	SID	Source	Target
32659844	13	Simple cysts are very rare in children and ADPKD in a parent should be excluded.	Simple fluid-filled sacs are very rare in children. ADPKD in parents should be excluded.

## Omitting Confidence Intervals and P-Values

PMID	SID	Source	Target
15156437	2	The summary odds ratio (OR) for bacteriologic cure rate significantly favored cephalosporins, compared with penicillin (OR,1.83; 95% confidence interval [CI], 1.37-2.44); the bacteriologic failure rate was nearly 2 times higher for penicillin therapy than it was for cephalosporin therapy (P=.00004).	Results favored cephalosporins (antibacterial antibiotics) over penicillin (another antibiotic).

## Carrying Over Understandable Sentences

PMID	SID	Source	Target
27852615	9	The interaction between dietary n-6:n-3 ratio and n-3 predicted the number of moves required to solve the most difficult planning problems in children aged 7-9 y and those aged 10-12 y, similar to results from the previous study.	The interaction between dietary n-6:n-3 ratio and n-3 predicted the number of moves required to solve the most difficult planning problems in children aged 7-9 y and those aged 10-12 y, similar to results from the previous study.

## Ignoring Sentences Not Relevant to Consumer Understanding

PMID	SID	Source	Target
32956536	3	Magnesium supplements are marketed for the prophylaxis of cramps but the efficacy of magnesium for this purpose remains unclear.	Magnesium supplements are used for preventing cramps, but their effectiveness is unclear.
	4	This is an update of a Cochrane Review first published in 2012, and performed to identify and incorporate more recent studies.	

## Resolving Anaphora and Pronouns

PMID	SID	Source	Target
30838456	1	This chapter covers antidepressants that fall into the class of serotonin (5-HT) and norepinephrine (NE) reuptake inhibitors.	This work covers antidepressants that block removal of the chemical messengers, serotonin (5-HT) and norepinephrine (NE).
	2	That is, <i>they</i> bind to the 5-HT and NE transporters with varying levels of potency and binding affinity ratios.	<i>These antidepressants</i> bind to 5-HT and NE transporters with varying effect.

## Explaining Complex Terms and Abbreviations only in the First Mention

PMID	SID	Source	Target
15316838	7	<i>Duloxetine</i> is a combined serotonin/norepinephrine reuptake inhibitor currently under clinical investigation for the treatment of women with stress urinary incontinence.	<i>Duloxetine</i> ( <i>a common antidepressant</i> ) blocks removal of serotonin/norepinephrine (chemical messengers) and is studied for treating women with bladder control loss from stress.
	Ex. 1	<i>It</i> is only available on prescription and commonly taken orally via capsules.	<i>This antidepressant</i> is prescribed and usually swallowed as a pill.
	Ex. 2	<i>Duloxetine</i> is only available on prescription and commonly taken orally via capsules.	<i>Duloxetine</i> is prescribed and usually swallowed as a pill.