



Production




Decay

$$\mathcal{L} \subset \frac{e^2}{16\pi^2 f_\pi} \pi^0 F_{\mu\nu} \hat{F}^{\mu\nu} + \frac{g}{M_X} \cdot \bar{X}_2 \sigma_{\mu\nu} X_1 F^{\mu\nu}$$

- M_X = TeV particle giving 5-point interaction, coupling X_2 , X_1 and photon
- Coupling g , will govern production and decay rates




- Decay length contour in $\Delta M = m_2 - m_1$ vs g plane
- In following slides I'll pick **$\Delta M = 20$ MeV** as a benchmark. ($m_1 = 30$ MeV, $m_2 = 50$ MeV).




- Production of pions (from FASER paper)
- Expect roughly **10^{17}** pions with \sim TeV energies headed towards FASER in Run3
- Can also think about η 's (or other mesons) later

Differential decay rate, vs Energy of outgoing χ_2 state
in Pion Rest frame. Normalized by g^2 .
 $m_1=30$ MeV, $m_2=50$ MeV




- Here I plot the differential decay rate.
- In pion rest frame, χ_2 produced with $E_2 \sim m_2$
- Integrating this curve gives a partial width for pion $\rightarrow (\chi_2 \chi_1 \gamma)$ of $\sim 10^{-20} \text{ GeV } g^2$
- Using a total pion width of ~ 10 eV, giving $\text{BR} \sim 10^{-12} g^2$
- So for Run3, expect about $10^5 g^2$ DM pairs

Number of decays in faser



- Here I plot* the number of decays inside FASER for:
 $m_1 = 30 \text{ MeV}$,
 $m_2 = 50 \text{ MeV}$,
 $M_x = 1 \text{ TeV}$,
 $N_\pi = 10^{17}$
- Need to fully explore (m_1, m_2) parameter space
- For HL-LHC, $N_{\text{decays}} \rightarrow N_{\text{decays}} * 20$
- Main bottleneck is production. If we can decouple the production/decay parameter ($g \rightarrow g_{\text{production}}, g_{\text{decay}}$) then this will free us quite a bit.
- $M_x = 1 \text{ TeV}$ was chosen somewhat arbitrarily. How small can we make this?

*To get an estimate, I assume that the momentum of the heavy state, $p_2 = 1 \text{ TeV}$. Need to make this plot properly, integrating over pion and x_1 momenta



To do:

- Need to look at annihilation rate ($x_1 x_1 \rightarrow \text{gamma gamma}$) and CMB bounds. Naively, however, these should be evaded due to the heavy $Mx^2 = (1 \text{ TeV})^2$ suppression.
- Need to think about constraints investigating these dipole interactions. Haven't put much thought into this yet.
- Need to full explore (m_1, m_2) parameter space (or start adding a ladder of states)
- Can start thinking about other meson decays. Heavier B, eta, eta' mesons are produced with a much lower flux ($\sim 10^{13}$ for run3), but will allow us to probe heavier m_2 masses.

