

Doping asymmetry in the three-band Hamiltonian for cuprate ladders: failure of the standard model of superconductivity in cuprates

Jeong-Pil Song,¹ S. Mazumdar,¹ and R. Torsten Clay²

¹Department of Physics, University of Arizona, Tucson, AZ 85721

²Department of Physics and Astronomy and HPC² Center for Computational Sciences, Mississippi State, MS 39762

(Dated: May 18, 2023)

S.1. LATTICE STRUCTURE




FIG. S1: Lattice structure of the three-band ladder. Colors indicate opposite signs of the Cu $d_{x^2-y^2}$ and O p orbitals. We take $t_{dp}=1$ and $t_{pp}=\{0.5, 0.6\}$. The DMRG calculation used open boundary conditions, with a Cu-O-Cu rung at each end of the lattice as shown in the figure.

S.2. LUTTINGER EXPONENTS

TABLE I: Luttinger correlation exponents K_ρ estimated from fitting charge density oscillations for electron (δ_e) and hole (δ_h) doping. Error bars are estimated from fitting the amplitude of charge oscillations in ladders of length up to $L=96$ rungs (see Fig. 3(d)).

U_p	t_{pp}	δ_e	K_ρ
3	0.5	0.0625	1.40(2)
3	0.5	0.0833	1.399(7)
3	0.5	0.1000	1.33(2)
3	0.5	0.1250	1.32(2)
3	0.5	0.2000	1.29(2)
3	0.5	0.2500	1.311(9)
4	0.6	0.0625	1.410(6)
4	0.6	0.0833	1.417(7)
4	0.6	0.1000	1.35(2)
4	0.6	0.1250	1.33(1)
4	0.6	0.2000	1.330(9)
4	0.6	0.2500	1.35(1)
U_p	t_{pp}	δ_h	K_ρ
3	0.5	0.0625	0.97(1)
3	0.5	0.1250	0.80(4)
4	0.6	0.0625	0.93(5)
4	0.6	0.1250	0.75(8)