

We want: $\Gamma = \Gamma_n \bar{e}^S \gtrsim H^4$ (during the phase transition)

$T_n \sim \mu_r^4$; (μ_r : the field tunnels to this value).

$$\alpha, \quad \boxed{S \lesssim 4 \ln \left(\frac{\mu_r}{\mu_{\min}^2} M_{\text{pl}} \right)} \quad \parallel \quad H \approx \frac{\mu_{\min}^2}{M_{\text{pl}}}$$

↓
valid for
temp. near
critical point
and below it.

We have:

$$V_{\text{GW}}(\mu_{\min}) \simeq -\epsilon^{3/2} v_{\text{IR}}^2 \mu_{\min}^4$$

$$T_c = \left(\frac{-8 V_{\text{GW}}(\mu_{\min})}{\pi^2 N^2} \right)^{1/4} \simeq \left(\frac{8 \epsilon^{3/2}}{\pi^2} \right)^{1/4} \sqrt{\frac{v_{\text{IR}}}{N}} \mu_{\min}$$

Characteristic length scale of the bubble:

$$R_{\text{bubble}} \simeq \frac{1}{\mu_{\min}} \frac{N}{v_{\text{IR}} \epsilon^{3/4}}$$

$\parallel N \gtrsim 4.4$
for higher
Ricci scalar
terms to be
neglected

For, $\frac{v_{\text{IR}}}{N} \simeq \frac{1}{5}, \epsilon \simeq \frac{1}{20}$

$$R_{\text{bubble}} \simeq 50 \mu_{\min}^{-1}$$

$\parallel \frac{v_{\text{IR}}}{N}, \frac{v_{\text{UV}}}{N} \ll 1$
 $\epsilon \simeq \frac{1}{20}$

We estimate: $T_c \simeq \mu_{\min}/2$.

So, in this case, the $O(4)$ symmetric solution would dominate. Further, assume that $\mu_r \simeq \mu_{\min}$.

$$S_4 \simeq \frac{9N^4}{8\pi^2} \frac{\mu_r^4}{-V_{\text{GW}}(\mu_r)}$$

[at zero temp.
dominating piece]

$$S_4 \simeq \frac{9N^4}{8\pi^2 \epsilon^{3/2} v_{\text{IR}}^2}$$

Therefore, we want :-

$$\frac{9N^4}{8\pi^2 \epsilon^{3/2} v_{IR}^2} \lesssim 4 \ln \left(\frac{M_{Pl}}{\mu_{min}} \right)$$

Constraints.
 $N \gtrsim 4.4$
 $\frac{v_{IR}}{N} \ll 1, \frac{v_{UV}}{N} \ll 1$
 we choose $\epsilon > 0$, and $\epsilon \ll 1$.

Remember that :

$$\mu_{min} \approx M_{Pl} \left(\frac{v_{IR}}{v_{UV}} \right)^{1/\epsilon} \left[\frac{8+6\epsilon+\epsilon^2+(2+\epsilon)\sqrt{4\epsilon+\epsilon^2}}{8+8\epsilon+2\epsilon^2} \right]^{1/\epsilon}$$

So we want :

$$\frac{9N^4}{32\pi^2 \epsilon^{3/2} v_{IR}^2 \ln \left(\frac{M_{Pl}}{\mu_{min}} \right)} \lesssim 1$$

Let's apply this to the intermediate brane scale:-
 $v_{IR} \rightarrow v_I$; $\mu_{min} \rightarrow \mu_I$. Choose ϵ based on that.

$$\boxed{\frac{9N^4}{32\pi^2 \epsilon^{3/2} v_I^2 \Omega_I} \lesssim 1}$$

let's assume, $\Omega_I \approx 30$.

where,
 $\mu_I \equiv M_{Pl} e^{-\Omega_I}$
 For 100 TeV intermediate brane say,
 $\Omega_I \approx 30$.

v_{UV} can be expressed as:-

$$v_{UV} \approx v_I \left(\frac{M_{Pl}}{\mu_I} \right)^\epsilon \left[\frac{8+6\epsilon+\epsilon^2+(2+\epsilon)\sqrt{4\epsilon+\epsilon^2}}{8+8\epsilon+2\epsilon^2} \right]$$

We should have $v_{UV} < N$.