

Question

You are an expert radiation oncologist from an academic center.

Please give your recommendation on the following case:

".....(text description)"

Question for the expert:

Please give a specific recommendation of your most favored therapeutic approach in this patient case based on the available information and your reasoning for it.

ChatGPT-4

Based on the available information, the patient has **locally recurrent breast cancer** with contralateral axillary lymph node involvement. Considering the patient's history and pathology findings, my recommendation for the therapeutic approach would include the following:

Systemic therapy: Given that the patient has hormone receptor-positive (ER/PR+) and HER2-negative disease, initiating endocrine therapy with an aromatase inhibitor (AI) would be my first choice. In addition, considering the prior discontinuation of tamoxifen due to intolerance, AI would be a better option for this patient. Additionally, since the patient has locally advanced and recurrent disease, I would also recommend adding a CDK4/6 inhibitor, such as palbociclib, ribociclib, or abemaciclib, in combination with the AI to improve the efficacy of treatment.

Radiation therapy: As the patient has not received radiation therapy in the past and given the involvement of contralateral axillary lymph nodes with extranodal extension, I would recommend considering regional nodal irradiation, which includes the chest wall (in the case of a mastectomy) and regional lymph nodes (axillary, supraclavicular, and internal mammary). This will help in reducing the risk of further local-regional recurrence and may improve overall survival.

Close monitoring: The patient should undergo regular follow-ups with physical examinations, and imaging as needed, to closely monitor her response to the treatment and to detect any signs of disease progression or new metastases.

Supportive care: It is essential to address any potential side effects or complications of the treatments mentioned above, such as fatigue, bone health issues, and lymphedema. This would involve coordinating with a multidisciplinary team of healthcare professionals including medical oncologists, radiation oncologists, surgeons, physical therapists, and other supportive care services.

In summary, my favored therapeutic approach for this patient would be the combination of endocrine therapy with an aromatase inhibitor and a CDK4/6 inhibitor, regional nodal irradiation, close monitoring, and appropriate supportive care. This recommendation is based on the patient's hormone receptor-positive, HER2-negative status, prior treatment history, and the need to address local-regional disease control while minimizing the risk of further recurrence.

Question

Please summarize the five experts' recommendations in short sentences.

ChatGPT-4

Expert 1: Recommends right postmastectomy radiation therapy (PMRT) to the chest wall and regional nodes, regardless of systemic therapy or axillary surgery. Omission of left-side radiation is suggested.

Expert 2: Suggests bilateral comprehensive radiation therapy to the chest wall and regional nodal irradiation (RNI), with discussion of additional systemic therapy.

Expert 3: Advises radiation to the right-sided axillary levels 1 to 3 and supraclavicular nodes only, with systemic therapy as a critical component for reducing the chance of metastases.

Expert 4: Recommends postoperative comprehensive radiation therapy with curative intent, targeting bilateral chest walls and lymph node areas, alongside subsequent aromatase inhibitor and CDK 4/6 inhibitor.

Expert 5: Favors right-sided locoregional radiation therapy with a hypofractionated dose to the chest wall and regional lymph nodes. The provision of chemotherapy would not alter this recommendation.

Question

As an expert radiation oncologist, which of the above experts' recommendation do you think the most proper for the patient described above?

ChatGPT-4

As an expert radiation oncologist, **I believe the recommendation provided**

by Expert 2 is the most appropriate for the patient described above.

This recommendation suggests bilateral comprehensive radiation therapy to the chest wall and regional nodal irradiation (RNI), with a discussion of additional systemic therapy.

This approach considers the patient's suboptimal initial treatment and the potential benefits of aggressive locoregional therapy in controlling the disease. By targeting both sides, it addresses the risk of residual disease on the left side and the possibility of aberrant lymphatic drainage after the axillary lymph node dissection. Furthermore, the recommendation includes a thorough assessment for potential primary or recurrent tumors through bilateral magnetic resonance imaging.

Adding systemic therapy to the treatment plan is essential to minimize the risk of distant metastases and maximize progression-free survival and overall survival. The multidisciplinary approach mentioned in this recommendation ensures a comprehensive treatment plan tailored to the patient's needs and overall prognosis.

Question

Which expert's recommendation is your initial suggestion closest to?

ChatGPT-4

My initial recommendation aligns most closely with Expert 3's recommendation. Expert 3 suggests radiating the right-sided axillary levels 1 to 3 and supraclavicular nodes only and emphasizes the critical role of systemic therapy in reducing the chance of metastases. This approach focuses on treating the current contralateral axillary nodal metastases and involves the use of endocrine therapy and a CDK4/6 inhibitor as part of the systemic therapy.

While Expert 3's recommendation omits irradiation of the bilateral chest walls and the left axilla/supraclavicular nodes to reduce morbidity, it acknowledges the importance of long-term follow-up and potential salvage treatment in case of isolated recurrence. This recommendation balances the need for effective treatment of the current disease presentation while minimizing potential side effects and considering the patient's overall prognosis.

Question

After seeing the other experts' recommendations, are you going to update your own recommendation of your most favored therapeutic approach in this patient case? If so, what is your updated recommendation?

ChatGPT-4

After reviewing the other five experts' recommendations, I would like to update my initial recommendation to provide a more comprehensive approach for the patient, **drawing inspiration from Expert 2's recommendation.**

Updated recommendation:

Perform thorough imaging, including bilateral magnetic resonance imaging, to assess for potential primary or recurrent tumors in the residual ipsilateral or contralateral breast tissue.

Administer bilateral comprehensive radiation therapy to the chest wall and regional nodal irradiation (RNI), targeting undissected axillary nodal basins, supraclavicular nodes, and internal mammary nodes.

Initiate systemic therapy, including endocrine therapy with an aromatase inhibitor, a CDK4/6 inhibitor, and discuss the possibility of chemotherapy to mitigate the risk of distant recurrence and maximize progression-free survival and overall survival.

Conduct close monitoring and long-term follow-up to detect any signs of recurrence or progression early and provide appropriate supportive care.

This updated recommendation is primarily based on Expert 2's suggestion. It takes into account the patient's suboptimal initial treatment and the potential benefits of aggressive locoregional therapy in controlling the disease. By targeting both sides, it addresses the risk of residual disease on the left side and the possibility of aberrant lymphatic drainage after the axillary lymph node dissection. Furthermore, the recommendation includes a thorough assessment for potential primary or recurrent tumors through bilateral magnetic resonance imaging.

Adding systemic therapy to the treatment plan is essential to minimize the risk of distant metastases and maximize progression-free survival and overall survival. The multidisciplinary approach mentioned in this recommendation ensures a comprehensive treatment plan tailored to the patient's needs and overall prognosis.