

Input Abstract

The Kinect video game (KVG) has received attention as an intervention method for cerebral palsy (CP). However, evidence remains limited. **PURPOSE** To investigate the effects of training using Xbox Kinect on lower extremity motor function, balance, and gait in adolescents with spastic diplegia CP. **METHODS** This study was a pilot randomized controlled trial. Ten participants were randomly allocated to either the KVG training group (n = 5) or the conventional training (CT) group (n = 5). The Selective Control Assessment of the Lower Extremity (SCALE) tool, Pediatric Balance Scale (PBS), and GAITRite were used for the outcome measurements. **RESULTS** In the comparison between the groups, the KVG group showed significant improvements in all the items in the SCALE (except for right hip abduction) and PBS score as compared with the CT group. **CONCLUSIONS** KVG training might be an effective intervention for the rehabilitation of adolescents with spastic diplegia CP.

**OBJECTIVE** To evaluate the effect of antrectomy in which resection was started from 2 cm or closer to the pylorus on % excess weight loss (EWL), nausea, vomiting, and complication rates. **STUDY DESIGN** Comparative study. **PLACE AND DURATION OF STUDY** Antalya Training and Research Hospital, from April 2018 to December 2018. **METHODOLOGY** Patients in whom laparoscopic sleeve gastrectomy (LSG) were done starting at a level of 2 cm or closer to pylorus were included in the study. Patients were divided into one of the two groups based on the distance between the pylorus and the resection margin: group 1 having resection ≤10 mm and group 2 at 11-20 mm. Above mentioned parameters were compared in both groups. **RESULTS** Ninety-two patients were included. Postoperative nausea and vomiting rates were similar in both groups. At the end of the first year, % EWL was 82.9% and 73.5% in groups 1 and 2 (p=0.003). **CONCLUSION** Starting antrectomy at a distance of 2 cm or less from the pylorus is safe and effective. Starting antrectomy at a distance of 1 cm or less from the pylorus in LSG provides effective weight loss without increasing complications. **KEY WORDS** Bariatric surgery, Antrectomy, Laparoscopic sleeve gastrectomy, Complications.

**OBJECTIVE** To comparatively examine the effectiveness and safety of the combination therapy of traditional Chinese medicine formula Bushen-Shugan granule and psychotherapy (BSSG-P) and Chinese herbal medicine Bushen-Shugan granule (BSSG) alone in the treatment of moderate to serious mood disorder in menopausal women. **METHODS** In our previous clinical studies, BSSG-P had been proved to be superior to BSSG, psychological treatment, and placebo in improving mild mood disorder in menopausal women. In this study, we analyzed the efficacy of BSSG-P and BSSG in the treatment of moderate to serious mood disorder. Eighty-five eligible participants, who were diagnosed as menopausal women with moderate to serious mood disorder and categorized as kidney deficiency and liver-qi stagnation pattern, were randomly assigned into two groups and treated with BSSG-P or BSSG. They were subjected to an 8-week treatment period and a 4-week follow-up study. The primary outcome instrument was the Greene Climacteric Scale, Self-Rating Depression Scale (SDS), and Self-Rating Anxiety Scale (SAS), respectively. **RESULTS** When comparing all time points with baseline, both BSSG-P and BSSG markedly decreased the total score of Greene, SDS, and SAS and the score of each dimension, in which BSSG-P exerted superior effect after 8-week treatment and 4-week follow-up ( P <0.05). Furthermore, BSSG-P also showed great advantage in reducing the score of Greene, SDS, and SAS for menopausal women with moderate mood disorder at the end of the 8th and 12th week when compared with BSSG ( P < 0.05), whereas there was no significant difference between groups at any time point for patients with serious mood disorder ( P >0.05). No serious event occurred in both groups, and no significant difference was found between groups in adverse event proportion. **CONCLUSIONS** BSSG-P was superior to BSSG in improving the physical and psychological symptoms of menopausal women with mood disorder. For patients with moderate mood disorder, BSSG-P showed obvious advantages; however, no superiority was observed for serious mood disorder.

Generated Summary

This study investigated the effects of training using Xbox Kinect on lower extremity motor function, balance, and gait in adolescents with spastic diplegia CP. Results showed that the KVG group had significant improvements in all the items in the SCALE (except for right hip abduction) and PBS score compared to the conventional training group. The findings suggest that KVG training may be an effective intervention for the rehabilitation of adolescents with spastic diplegia CP.

This study evaluated the impact of antrectomy in which resection was started from 2 cm or closer to the pylorus on % excess weight loss (EWL), nausea, vomiting, and complication rates. Ninety-two patients were included and divided into two groups based on the distance between the pylorus and the resection margin. Postoperative nausea and vomiting rates were similar in both groups. At the end of the first year, % EWL was significantly higher in group 1 which had resection ≤10 mm compared to group 2 at 11-20 mm (p=0.003). Overall, starting antrectomy at a distance of 2 cm or less from the pylorus in laparoscopic sleeve gastrectomy is safe and effective and provides effective weight loss without increasing complications.

This study examined the effectiveness and safety of a combination therapy of traditional Chinese medicine formula Bushen-Shugan granule and psychotherapy (BSSG-P) and Chinese herbal medicine Bushen-Shugan granule (BSSG) alone in the treatment of moderate to serious mood disorder in menopausal women. The study found that BSSG-P was superior to BSSG in improving the physical and psychological symptoms of menopausal women with mood disorder, especially for those with moderate mood disorder. No serious event occurred in both groups and no significant difference was found between groups in adverse event proportion.