

Flan-T5 Summary	GPT-3 Summary (One Sentence)	GPT-3 Summary
Population-based informatics systems can enable sustained increases in mammography screening rates beyond rates seen with office-based visit reminders.	The use of population-based informatics systems increased mammography screening rates, with intervention patients having significantly higher completion rates than control patients after 3 years.	This study examined the ability of a health information technology system to facilitate population-based breast cancer screening. The study was a two-year follow-up after a one-year cluster randomized trial of 32,688 women aged 42-69 receiving care within a 12-practice primary care network. The system identified women overdue for mammograms, connected them to primary care providers, generated outreach letters, monitored mammography scheduling and completion, and provided practice delegates with a list of overdue patients for reminder phone calls. Results showed that adjusted completion rates for mammography screening were significantly higher for intervention patients in the prevalent cohort after 3 years (51.7% vs. 45.8%) and for the incident cohort after 2 years (53.8% vs. 48.7%). These results suggest that population-based informatics systems can enable sustained increases in mammography screening rates.
The result of CBT for SAD was maintained through lockdown period and was associated with significantly less COVID-19 related fear.	Cognitive-Behavioral Therapy (CBT) for Social Anxiety Disorder (SAD) was shown to be more effective at reducing social anxiety than Psychoeducational-Supportive Therapy (PST), and was associated with significantly less fear of COVID-19.	The study compared the effects of cognitive-behavioral therapy (CBT) and psychoeducational-supportive therapy (PST) on social anxiety disorder (SAD) before and during the COVID-19 pandemic. Results showed that the CBT group had significantly better outcomes than the PST group at post-intervention, and this effect was maintained during the lockdown period. In addition, the CBT group had significantly less fear of COVID-19 than the PST group. The study also found that social anxiety severity was positively correlated with fear of COVID-19, and that individuals with comorbidities had significantly more fear. The hierarchical multiple regression revealed that SPIN post-intervention, COVID-19 fear, and duration of SAD predicted social anxiety severity during the post-lockdown period.
Combining aerobic exercise training with moderate- to high-calorie restriction has greater effects on proximal aortic stiffness than exercise alone in older adults with obesity.	A combination of aerobic exercise and moderate caloric restriction led to greater improvements in proximal aortic stiffness compared to aerobic exercise alone in older adults with obesity.	In this study, 160 older adults with obesity were assigned to three groups: aerobic exercise only, aerobic exercise with moderate caloric restriction, or aerobic exercise with more intensive caloric restriction. Results showed that the greatest improvements in aortic stiffness were found in the aerobic exercise with moderate caloric restriction group, with increased distensibility and decreased pulse wave velocity. Increases in aortic distensibility were correlated with improvements in body weight and body fat distribution, but these associations were not statistically significant after adjusting for multiple comparisons. These results suggest that combining aerobic exercise with moderate caloric restriction may be more effective than exercise alone in improving aortic stiffness in older adults with obesity.