

Input:

To investigate the optimal blood pressure (BP) in patients with coronary artery disease (CAD), we conducted subgroup analysis using SPRINT data. The study sample included 1206 participants with CAD (of whom 692 underwent coronary revascularization) and 8127 participants without CAD. Participants were randomized into two groups (systolic BP target of 140 mm Hg vs. 120 mm Hg). The primary outcome was a composite of cardiovascular events. After a median follow-up of 3.9 years, the hazard ratios (HRs) for the primary outcome were 0.65 (95% confidence interval (CI) 0.53-0.79) and 1.05 (95% CI 0.76-1.46) among those in the non-CAD and CAD subgroups, respectively (P value for interaction 0.02). Intensive BP treatment was a protective factor for all-cause death (HR 0.60, 95% CI 0.37-0.96) in the CAD subgroup, compared with standard BP treatment. The HRs (95% CI) for stroke were 3.57 (1.17-10.85) and 1.03 (0.29-3.62) among those in the coronary revascularization and non-revascularization subgroups, respectively (P value for interaction 0.13). For safety events, intensive BP treatment increased the risk of hypotension (HR 2.00, 95% CI 1.06-3.79) and electrolyte abnormalities (HR 2.38, 95% CI 1.25-4.56) in the CAD subgroup, while the risk of serious adverse events did not increase (HR 1.03, 95% CI 0.88-1.20). These results suggest that positive benefits from intensive BP treatment might be attenuated in patients with CAD who are under better secondary prevention. The risk of stroke might increase at the systolic BP target of 120 mm Hg in case of coronary revascularization, although the confidence interval was wide.

Generated Technical Summary:

The study investigated the optimal blood pressure for patients with coronary artery disease (CAD) and found that after a median follow-up of 3.9 years, intensive BP treatment was a protective factor for all-cause death in the CAD subgroup. However, the risk of stroke might increase in patients who underwent coronary revascularization. Intensive BP treatment was also associated with an increased risk of hypotension and electrolyte abnormalities in the CAD subgroup, while the risk of serious adverse events did not increase.

Generated Simplified Summary:

This study looked at people with and without coronary artery disease (CAD), a type of heart disease. They wanted to find out if lowering the blood pressure to a certain level (120 mmHg) was better for their health than having it at a slightly higher level (140 mmHg). They found that for people without CAD, lowering their blood pressure was helpful. But for those with CAD, lowering their blood pressure did not help as much. They also found that people who had had a special surgery to help their CAD (called coronary revascularization) had a higher chance of having a stroke if their blood pressure was lowered too much. In conclusion, it is important to keep a healthy blood pressure, but for people with CAD, it is important to not lower it too much.