

Select a page

Document-Level Cluster View

Which document would you like to edit?

701

Read the Passage

dobutamine and exercise induced myocardial ischaemia . objective : to determine whether pharmacological stress leads to prolonged but reversible left ventricular dysfunction in patients with coronary artery disease , similar to that seen after exercise. design : a randomised crossover study of recovery time of systolic and diastolic left ventricular function after exercise and dobutamine induced ischaemia . subjects : 10 patients with stable angina , angiographically proven coronary artery disease , and normal left ventricular function. interventions : treadmill exercise and dobutamine stress were performed on different days. quantitative assessment of systolic and diastolic left ventricular function was performed using transthoracic echocardiography at baseline and at regular intervals after each test.

Cluster4

Cluster Name:('Heart-related conditions,') with explanation: These questions have been clustered together because they all pertain to heart-related conditions, specifically focusing on terms and diseases such as myocardial stunning, ischaemia, coronary artery disease, left ventricular dysfunction, and angina.

Questions belonging to this cluster

What is the result of dobutamine and exercise induced ischaemia?::myocardial stunning
What is the term for the condition that causes reversible left ventricular dysfunction?::myocardial ischaemia
What does dobutamine induced ischaemia cause?::left ventricular dysfunction

Cluster5

Cluster6

Cluster Name:('cause,') with explanation: These questions should be related in such a way that they talk about what drugs cause which disease

Questions belonging to this cluster

What pharmacological stress is believed to cause myocardial stunning?::dobutamine
What dobutamine stress is believed to cause myocardial stunning?::ischaemia