

Table. Classification of architectural types, including pennate and non-pennate muscles, in human forearm muscles. R^2 of the pennate muscles are higher than 0.9, while lower than 0.9 for the non-pennate muscles. Only for the type of palmaris longus, this study differs from the previous studies (Lee et al. (1), Matulionis (2)). (Abbreviations: non-pennate (N), pennate (P), confidence interval (CI).)

Muscle	Non-Pennate /Pennate Muscle (1, 2)	Non-Pennate/Pennate Muscle (this study)
Anconeus	N	N ($R^2=0.82\pm 0.06$, $P=0.001$)
Abductor pollicis longus	P	P ($R^2=0.95\pm 0.02$, $P<0.001$)
Extensor carpi radialis brevis	P	P ($R^2=0.92\pm 0.10$, $P=0.566$, 95% CI: [-0.05, 0.09])
Extensor carpi radialis longus	N	N ($R^2=0.85\pm 0.13$, $P=0.254$, 95% CI: [-0.14, 0.04])
Extensor carpi ulnaris	P	P ($R^2=0.97\pm 0.01$, $P<0.001$)
Extensor digitorum communis	P	P ($R^2=0.96\pm 0.02$, $P<0.001$)
Extensor digiti minimi	P	P ($R^2=0.94\pm 0.08$, $P=0.186$, 95% CI: [-0.02, 0.09])
Extensor pollicis longus	P	P ($R^2=0.96\pm 0.02$, $P<0.001$)
Flexor carpi radialis	P	P ($R^2=0.97\pm 0.02$, $P<0.001$)
Flexor carpi ulnaris	P	P ($R^2=0.98\pm 0.01$, $P<0.001$)

Flexor digitorum profundus	P	P ($R^2=0.97\pm 0.01$, $P<0.001$)
Flexor digitorum superficialis	P	P ($R^2=0.98\pm 0.01$, $P<0.001$)
Flexor pollicis longus	P	P ($R^2=0.95\pm 0.09$, $P=0.113$, 95% CI: [-0.01, 0.11])
Palmaris longus	N	P ($R^2=0.95\pm 0.02$, $P<0.001$)
Pronator quadratus	N	N ($R^2=0.70\pm 0.10$, $P<0.001$)
Pronator teres	P	P ($R^2=0.95\pm 0.01$, $P<0.001$)
Supinator	N	N ($R^2=0.76\pm 0.04$, $P<0.001$)

References:

1. Lee D, Li Z, Sohail QZ, Jackson K, Fiume E, Agur A. A three-dimensional approach to pennation angle estimation for human skeletal muscle. *Computer Methods in Biomechanics and Biomedical Engineering* 2015;18(13):1474-1484.
2. Matulionis DH. A Functional Study of the Forearm Musculature of the Human and *Macaca mulatta*. *Bios* 1966;37(1):3-14.