

	$A_0 = \{a_1, a_2, a_3, a_4\}$	$B_0 = \{b_1, b_2, b_3, b_4\}$	Inverse	$A_1 = \{a_1, a_2, a_3, a_4\}$	$B_1 = \{b_1, b_2, 0, 0\}$
	$A_2 = \{a_1, a_2, 0, 0\}$	$B_2 = \{b_1, b_2, 0, 0\}$	N-M	$A_1 = \{a_1, a_2, a_3, 0\}$	$B_1 = \{b_1, b_2, b_3, 0\}$
	$a_1 > a_2 > a_3 > a_4 > b_1 > b_2 > b_3 > b_4 > 0$				
Default	$A_1 = \{a_1, a_2, 0, 0\}$	$B_1 = \{b_1, b_2, b_3, b_4\}$	LBC	$A_1$ $(a_1, a_2)$ <del><math>(a_1, a_4)</math></del> <del><math>(a_2, a_4)</math></del>	$B_1$ $(b_1, b_2)$ <del><math>(b_1, b_4)</math></del> <del><math>(b_2, b_4)</math></del>
				$(a_1, a_3)$ $(a_2, a_3)$ <del><math>(a_3, a_4)</math></del>	$(b_1, b_3)$ $(b_2, b_3)$ <del><math>(b_3, b_4)</math></del>