

Intrinsic toroidal rotation driven by turbulent and neoclassical processes in tokamak plasmas from global gyrokinetic simulations: Supplemental Material

Hongxuan Zhu,^{1,2} T. Stoltzfus-Dueck,¹ R. Hager,¹ S. Ku,¹ and C. S. Chang¹

¹Princeton Plasma Physics Laboratory, Princeton, NJ 08540

²Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544

Here, we provide an ordering estimate for the momentum fluxes driven by turbulence. From equation (7) in the main text, the kinetic stress is calculated as

$$\Pi_{\parallel}^D = \frac{2\pi}{Z_i e m_i^2} \int d\mu d\mathcal{P}_{\varphi} d\tilde{H} \oint d\varphi \Delta F_i p_{i\varphi}. \quad (1)$$

Considering the effects from turbulence, we have $\Delta F_i = \int d\tau (-\tilde{\mathbf{R}} \cdot \nabla F_i - \tilde{p}_{\parallel} \partial_{p_{\parallel}} F_i)$. Since $(\tilde{\mathbf{R}}, \tilde{p}_{\parallel})$ are derived from the Hamiltonian \tilde{H} and conserve phase space, we have $\tilde{\mathbf{R}} \cdot \nabla F_i + \tilde{p}_{\parallel} \partial_{p_{\parallel}} F_i = [\nabla \cdot (B_{\parallel}^* \tilde{\mathbf{R}} F_i) + \partial_{p_{\parallel}} (B_{\parallel}^* \tilde{p}_{\parallel} F_i)] / B_{\parallel}^*$, where the spatial-divergence term dominates because $k_{\parallel} \ll k_{\perp}$ for turbulent fluctuations. Since only the non-axisymmetric $\tilde{F}_i \sim \rho_* F_i$ contributes to the toroidal average, we have $\Delta F_i \sim \tau_{\text{orb}} k_r |\mathbf{v}_E| \rho_* F_i$, where $\tau_{\text{orb}} \sim qR/v_t$, q is the safety factor, $|\mathbf{v}_E| \sim \rho_i^{-1} \rho_* T_e / eB$, $F_i \sim n_i / v_i^3$, and k_r is some wavenumber corresponding to the spatial variation of the turbulence. Replacing $\int p_{i\varphi} d\mu d\mathcal{P}_{\varphi} d\tilde{H}$ with $\mu v_{i\varphi}^2 \tilde{H}$ at thermal velocity v_t , we obtain

$$\Pi_{\parallel}^D \sim q(k_r \rho_i) \rho_*^2 n_i T_i R^3. \quad (2)$$

Next, from $\Pi_{\parallel}^E = \sum_s \langle \int d^3v F_s p_{s\varphi} (\mathbf{v}_E \cdot \nabla V) \rangle$ and assuming $p_{i\varphi} \sim \rho_* m_i R v_t$ from the turbulent parallel electric fields, the fluid stress can be estimated as

$$\Pi_{\parallel}^E \sim \alpha \rho_*^2 n_i T_i a R^2, \quad (3)$$

where α is a small number representing a symmetry-breaking mechanism. Therefore, Π_{\parallel}^D can be comparable to Π_{\parallel}^E if $k_r \rho_i \sim \alpha a / qR \ll 1$, so that even a small spatial variation of turbulence can result in a nonnegligible Π_{\parallel}^D .

Using the usual gyroBohm estimate, we have $a Q_i / v_t \sim \rho_*^2 n_i T_i a^2 R$. Therefore, both Π_{\parallel}^D and Π_{\parallel}^E estimated above should in principle be able to drive significant intrinsic toroidal rotation in the sense that $|v_t \Pi_{\parallel} / a Q_i| \sim 1$ [1]. The assumed ρ_* dependence of Π_{\parallel}^D and Π_{\parallel}^E are important for this conclusion. To investigate the ρ_* scaling, we also simulated a turbulent plasma in a larger machine ITER with a smaller ρ_* . The density and temperature profiles are shown in Fig. 1, which correspond to a numerically modelled full-current ITER H-mode plasma [2, 3]. Compared to DIII-D, ρ_* is 4 times smaller for the simulated ITER plasma, so that the spatial resolution needed in the simulation is 16 times higher. To reduce computational costs, we use the adiabatic-electron model.

Simulation results are shown in Fig. 2. We found that turbulence is active in both the core and the edge despite the H-mode E_r well, consistent with Refs. [2, 3],




FIG. 1. Equilibrium density and temperature profile.




FIG. 2. (a) E_r along the outboard midplane in the presence of turbulence. (b) Π_{\parallel}^D and Π_{\parallel}^E at the same time. (c) $e\tilde{\Phi}/T_e$ near the outboard midplane. The dashed curve and the solid curve are respectively the $\psi_n = 0.5$ and the $\psi_n = 1$ flux surface.

even though the adiabatic-electron model is used here. In the core, Π_{\parallel}^D and Π_{\parallel}^E become comparable as soon as turbulence develops, when zonal flows are also generated. Compared to the zonal flows in DIII-D, here they have many more radial periods due to the larger machine size. This suggests that $k_r \rho_i$ can still remain finite in ITER, assuming k_r is the characteristic wavenumber of zonal flows. In fact, similar to the DIII-D simulation, here $v_t \Pi_{\parallel}^D / a Q_i$ also reaches 0.5, suggesting that Π_{\parallel}^D may still be important for intrinsic toroidal rotation in the core region of ITER, where ρ_* is much smaller.

-
- [1] W. A. Hornsby *et al.* Nucl. Fusion **57**, 046008 (2017).
 [2] C. S. Chang *et al.*, Nucl. Fusion **57**, 116023 (2017).
 [3] C. S. Chang *et al.* Phys. Plasmas **28**, 022501 (2021).