

Article:
A university researcher sent 1,500 identical CVs to German firms - except that some bore the name Meryem Ozturk and others the name Sandra Bauer. In 18.8% of cases Sandra Bauer was invited for interview, whereas the figure for Meryem was just 13.5%. When the photo of Meryem showed her in a headscarf only 4.2% invited her. The study was published by the Institute for the Study of Labour, in Bonn. The researcher was Doris Weichselbaumer from the University of Linz, in Austria. The findings are especially significant in light of Germany's current efforts to integrate record numbers of Muslim migrants, many of them refugees from the conflicts in Syria, Iraq and Afghanistan. More than a million asylum seekers reached Germany last year, and there has been a backlash from nationalist groups, especially Pegida and Alternative for Germany (AfD). With an estimated three million people of Turkish origin, Germany is home to the largest Turkish diaspora. In her study, Ms Weichselbaumer said that "a very modern binding of the headscarf was chosen to signal that the applicant was a young, modern woman who could easily fit into a secular environment". It is normal in Germany to attach a photo to a job application, she pointed out. The result of the experiment "implies that the candidate with the headscarf had to send 4.5 times as many applications as an identical applicant with a German name and no headscarf to receive the same number of callbacks for interview", she said. ...

Ground truth summary:
An experiment in Germany has found evidence of job discrimination against women with Turkish names - and even more if they wear an Islamic headscarf.

BART summary:
Women who apply for jobs in Germany wearing headscarves are much less likely to be invited for interview, a study suggests.

TriSum rationale:
<rationale> <aspects> Job discrimination against women with Turkish names in Germany. Job discrimination against women wearing an Islamic headscarf in Germany. <triples> [women with Turkish names | face discrimination in | job applications in Germany], [women wearing an Islamic headscarf | face even more discrimination in | job applications in Germany].

TriSum summary:
Women with Turkish names in Germany face discrimination in job applications, and this discrimination is intensified for women wearing an Islamic headscarf.

Article:
Common indications for adult tonsillectomy include tonsillar hypertrophy, obstructive sleep apnea, and chronic tonsillitis. Although there are many studies looking into how best to prevent and treat post-operative pain and nausea, there are very few that give recommendations to the patients pre-operatively. This study's aim is to evaluate if pre-operative oral hydration has an impact in reducing post-operative pain, nausea and vomiting after adult tonsillectomies. If pre-operative hydration can reduce the pain associated with the procedure, then this could also be an adjunct to help reduce the amount of narcotic pain medication that is required.~The goal of this study is to evaluate if the amount of pre-operative oral hydration can reduce the amount of pain and nausea/vomiting in patients undergoing adult tonsillectomies. This study will be measuring post-operative pain and nausea scores at several points after the adult tonsillectomy procedure for participants that have consumed different amounts of clear fluid in the 24 hours leading up to their nothing by mouth (NPO) status. Participants will be randomized to one of three categories of preoperative fluid: 0.5 - 1.5 Liters (L), 1.5 - 3 L, and 3 - 4.5 L, which will correlate to the low, medium and high fluid intake groups, respectively. Participants will be given a 1000 mL measuring water bottle that they can use to accurately record their fluid consumption in the 24 hours prior to their pre-operative NPO status. The fluid consumption that they record will include all liquids (i.e., water, flavored water, juice, coffee, tea, soda, milk, alcohol). They will be asked to keep a record of the volume of caffeinated beverage versus un-caffeinated beverages.~After surgery the participants will be given a form to record their pain and nausea scores. Participants will record their pain and nausea by using two separate 10-point scales at 7 different points after the procedure. These scores will be measured immediately after surgery, and at 4 hours, 8 hours, 24 hours, 72 hours, 5 days and 7 days postoperative. This form, along with a count of opioid medication used, will be returned to the physician at the first post-operative follow-up visit.

Ground truth summary:
Tonsillectomy is commonly described as one of the most painful procedures that an adult can undergo. This study's aim is to evaluate if pre-operative oral hydration has an impact in reducing post-operative pain, nausea and vomiting, and opioid medication use after adult tonsillectomies. Participants will be randomized to a low, medium and high fluid intake groups and will record fluid intake on the day before surgery. After surgery, participants will record their pain and nausea at timepoints up to 7 days post-operative. Medication use will be recorded up to the first post-operative follow-up visit with their physician.

BART summary:
The goal of this study is to evaluate if the amount of pre-operative oral hydration can reduce the amount of pain and nausea/vomiting in patients undergoing adult tonsillectomies.

TriSum rationale:
<rationale> <aspects> Purpose of the study. Concern about tonsillectomy pain. Role of pre-operative oral hydration. Randomization of participants into fluid intake groups. Recording of fluid intake prior to surgery. Post-operative recording of pain and nausea. Recording of opioid medication use. <triples> [Study | aims to| evaluate impact of pre-operative hydration on post-operative outcomes], [Tonsillectomy | is described as| painful procedure for adults], [Participants | will be randomized to | fluid intake groups: low, medium, high], [Participants | will record | fluid intake before surgery], [Participants | will record | pain and nausea post-operatively], [Participants | will record | opioid medication use until first follow-up].

TriSum summary:
This study focuses on the possible influence of pre-operative oral hydration on post-operative pain, nausea, vomiting, and the use of opioid medication. Individuals taking part in the study will be divided into low, medium, and high fluid intake categories. On the day before their procedure, they will note down their fluid consumption. After undergoing the tonsillectomy, these participants will monitor and record their pain, nausea for up to seven days, and also their opioid medication intake until their initial post-operative check-up with the doctor.