

README file for the ancillary files to A dispersive study of final-state interactions in $K \rightarrow \pi\pi\pi$ amplitudes

Véronique Bernard^{1*}, Sébastien Descotes-Genon^{1†}, Marc Knecht^{2‡}, Bachir Moussallam^{1§}

¹ *Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France*

² *Centre de Physique Théorique, CNRS/Aix-Marseille Univ./Univ. de Toulon (UMR7332)
CNRS-Luminy Case 907, 13288 Marseille Cedex 9, France*

The present text provides a description of the content of the ancillary files to the solution of the Omnès-Khuri-Treiman-Sawyer-Wali integral equations for the $K \rightarrow \pi\pi\pi$ amplitudes. These solutions are expressed as linear combinations of a set of fundamental functions (I is an isospin label)

$$\begin{aligned} \begin{pmatrix} M_I(s) \\ N_I(s) \end{pmatrix} &= \begin{pmatrix} \mu_0 \\ \nu_0 \end{pmatrix} \mathcal{S}_I^{(0)}(s) + \begin{pmatrix} \mu_1 \\ \nu_1 \end{pmatrix} \mathcal{S}_I^{(1)}(s) + \begin{pmatrix} \mu_2 \\ \nu_2 \end{pmatrix} \mathcal{S}_I^{(2)}(s) + \begin{pmatrix} \mu_3 \\ \nu_3 \end{pmatrix} \mathcal{S}_I^{(3)}(s), \quad I = 0, 1, 2 \\ \tilde{M}_1(s) &= \tilde{\mu}_1 \tilde{\mathcal{S}}_1(s), \quad \tilde{N}_1(s) = \tilde{\nu}_0 \tilde{\mathcal{S}}_1^{(0)} + \tilde{\nu}_1 \tilde{\mathcal{S}}_1^{(1)}, \quad I = 1, 2, \end{aligned} \tag{1}$$

modulated by the complex coefficients $\mu_n, \nu_n, \tilde{\nu}_{0,1}, \tilde{\mu}_1$. Some of these coefficients and fundamental functions carry dimensions, and will be given in units of GeV^N , with N as given in the following Table:

N	coefficients	fundamental functions
-4	$\mu_2, \mu_3, \nu_2, \nu_3, \tilde{\mu}_1, \tilde{\nu}_1$	-
-2	$\mu_1, \nu_1, \tilde{\nu}_0$	$\mathcal{S}_1^{(0)}$
0	μ_0, ν_0	$\mathcal{S}_0^{(0)}, \mathcal{S}_2^{(0)}, \mathcal{S}_1^{(1)}, \tilde{\mathcal{S}}_1^{(0)}$
+2	-	$\mathcal{S}_0^{(1)}, \mathcal{S}_2^{(1)}, \mathcal{S}_1^{(2)}, \mathcal{S}_1^{(3)}, \tilde{\mathcal{S}}_1^{(1)}, \tilde{\mathcal{S}}_2^{(0)}, \tilde{\mathcal{S}}_1$
+4	-	$\mathcal{S}_0^{(2)}, \mathcal{S}_2^{(2)}, \mathcal{S}_0^{(3)}, \mathcal{S}_2^{(3)}, \tilde{\mathcal{S}}_2^{(1)}$

The numerical values of the complex coefficients are given in these units by (the coefficient $\tilde{\mu}_1$ remains undetermined so far):

n	μ_n	ν_n	$\tilde{\nu}_n$	$\tilde{\mu}_n$
0	$+3.8(4) - i0.570(2)$	$-4.7(3) - i2.37(10) \cdot 10^{-2}$	$-2.04(1.2) + i4.8(5) \cdot 10^{-4}$	-
1	$-676.1(4.0) + i7.27(2)$	$+26.7(2.8) + i0.30(1)$	$+433.2(12.2) - i6.0(1.3) \cdot 10^{-4}$	n.d.
2	$+559.7(18.5) - i16.80(5)$	$-46.0(14.2) - i0.74(3)$	-	-
3	$-1072.6(13.6) + i7.57(2)$	$+123.9(12.6) - i0.28(1)$	-	-

The ancillary files provide the numerical tables for the set of fundamental solutions as functions of s according to the following correspondence:

$$\begin{aligned} \text{S_n.dat} & \quad \mathcal{S}_I^{(n)}(s) & \quad I = 0, 1, 2, n = 0, 1, 2, 3 \\ \text{St_n.dat} & \quad \tilde{\mathcal{S}}_I^{(n)}(s) & \quad I = 1, 2, n = 0, 1 \\ \text{St1.dat} & \quad \tilde{\mathcal{S}}_1(s) \end{aligned}$$

Each of the files S_n.dat displays seven columns. The first column gives the value of s in GeV^2 . The remaining six columns give successively the corresponding values of the functions $\text{Re } \mathcal{S}_0^{(n)}(s)$, $\text{Im } \mathcal{S}_0^{(n)}(s)$, $\text{Re } \mathcal{S}_1^{(n)}(s)$, $\text{Im } \mathcal{S}_1^{(n)}(s)$, $\text{Re } \mathcal{S}_2^{(n)}(s)$,

* veronique.bernard@ijclab.in2p3.fr

† sebastien.descotes-genon@ijclab.in2p3.fr

‡ knecht@cpt.univ-mrs.fr

§ bachir.moussallam@ijclab.in2p3.fr

$\text{Im } \mathcal{S}_2^{(n)}(s)$, in units of GeV^N , with N as specified above. The two files **St_n.dat** display five columns each, the first one giving the value of s in GeV^2 . The four remaining columns give successively the values of the functions $\text{Re } \tilde{\mathcal{S}}_1^{(n)}(s)$, $\text{Im } \tilde{\mathcal{S}}_1^{(n)}(s)$, $\text{Re } \tilde{\mathcal{S}}_2^{(n)}(s)$, $\text{Im } \tilde{\mathcal{S}}_2^{(n)}(s)$ in the appropriate units of GeV^N . Finally, the last file, **St1.dat**, contains three columns displaying successively the values of s , $\text{Re } \tilde{\mathcal{S}}_1(s)$ and $\text{Im } \tilde{\mathcal{S}}_1(s)$, all in units of GeV^2 . The range of values of s that is covered in the files is $-2 \text{ GeV}^2 \leq s \leq 10 \text{ GeV}^2$.

Also provided are the numerical values of the S and P -wave $\pi\pi$ phase-shifts $\delta_0(s)$, $\delta_1(s)$, $\delta_2(s)$ from which the solutions of the KT equations have been computed. They are given in the three files: **phase00+omnes00.dat**, **phase11+omnes11.dat** and **phase02+omnes02.dat**. In each of these files the numbers in the first column correspond to s (in GeV^2), those in the second and third column to the phase-shift (in radians) and to the absolute value of the Omnès function (dimensionless).