

Chirostatic response of microswimmers in fluids with odd viscosity

Supplemental Material

Yuto Hosaka,^{1,*} Michalis Chatzittofi,¹ Ramin Golestanian,^{1,2,†} and Andrej Vilfan^{1,3,‡}

¹Max Planck Institute for Dynamics and Self-Organization (MPI-DS), Am Fassberg 17, 37077 Göttingen, Germany

²Rudolf Peierls Centre for Theoretical Physics, University of Oxford, Oxford OX1 3PU, United Kingdom

³Jožef Stefan Institute, 1000 Ljubljana, Slovenia

ANGULAR VELOCITY OF A DISK-SHAPED SWIMMER IN TWO-DIMENSIONAL ODD FLOWS

Here, we derive the angular velocity of a disk-shaped swimmer in a two-dimensional (2D) odd flow. The 2D flat and thin fluid layer is assumed to be perpendicular to the odd viscosity axis \mathbf{e} . By defining the effective pressure $\hat{P} = \hat{p} - \hat{\eta}^o \mathbf{e} \cdot (\nabla \times \hat{\mathbf{v}})$, the stress tensor can be expressed as

$$\hat{\sigma}_{ij} = -\hat{P}\delta_{ij} + \hat{\eta}^e(\partial_i\hat{v}_j + \partial_j\hat{v}_i) + 2\hat{\eta}^o e_\ell \epsilon_{\ell jk} \partial_k \hat{v}_i. \quad (\text{S1})$$

First, we solve the flow around a no-slip disk of radius a rotating with the angular velocity $\hat{\Omega} = \hat{\Omega}\mathbf{e}$. From the no-slip boundary condition $\hat{\mathbf{v}} = \hat{\Omega} \times \mathbf{r} = a\hat{\Omega} \times \mathbf{n}$ at $r = a$, where \mathbf{n} is the normal vector at the disk perimeter, the solution is given by a rotlet singularity

$$\hat{\mathbf{v}} = \frac{a^2\hat{\Omega}}{r^2} \mathbf{e} \times \mathbf{r}. \quad (\text{S2})$$

Applying the pressureless condition ($\hat{P} = 0$) simplifies the stress tensor (S1) to give the traction

$$\hat{\mathbf{f}} = \hat{\boldsymbol{\sigma}} \cdot \mathbf{n} = -2\hat{\Omega} \frac{a^2}{r^2} (\hat{\eta}^e \mathbf{e} \times \mathbf{n} + \hat{\eta}^o \mathbf{n}). \quad (\text{S3})$$

The net torque acting on the disk contour reads

$$\hat{\mathbf{T}} = -4\pi a^2 \hat{\eta}^e \hat{\Omega} \mathbf{e}, \quad (\text{S4})$$

without any modifications due to the odd viscosity, which is consistent with the result of the torque acting on no-slip rigid objects in 2D odd flows [S1].

Applying the obtained traction and torque to the Lorentz reciprocal theorem and using the volume conservation within the swimmer ($\int_C ds \mathbf{n} \cdot \mathbf{v}^s = 0$), we find

$$\hat{\Omega} = -\frac{1}{2\pi a^2} \int_C ds \mathbf{n} \times \mathbf{v}^s, \quad (\text{S5})$$

which contains only the intrinsic rotation and is independent of the odd viscosity, as derived for a classical fluid [S2]. In contrast to the precession that occurs in 3D, odd viscosity therefore cannot contribute to the rotation of a swimmer in a 2D incompressible fluid.

DEFINITION OF THE EULER ANGLES

To describe the angular dynamics of a chiral microswimmer with an intrinsic angular velocity determined in the body frame, it is convenient to work in the swimmer-fixed frame with the basis $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ oriented such that $\mathbf{e}_3 \equiv \mathbf{p}$. We describe the rotation of the swimmer relative to the laboratory frame with the basis $(\mathbf{e}_x, \mathbf{e}_y, \mathbf{e}_z)$, where $\mathbf{e}_z \equiv \mathbf{e}$, using the Euler angles (ϕ, θ, ψ) [S3] (see Fig. S1). The rotation follows the zxz -convention, which is composed of a rotation of angle ϕ about \mathbf{e}_z , a rotation of θ about the x axis of the resulting reference frame and, a rotation of ψ about the body fixed axis \mathbf{e}_3 . Then the rotation matrix \mathbf{R} , which transforms the laboratory frame basis vectors to the body-fixed basis vectors (active transformation) via the relation

$$(\mathbf{e}_1 \ \mathbf{e}_2 \ \mathbf{e}_3) = \mathbf{R} \cdot (\mathbf{e}_x \ \mathbf{e}_y \ \mathbf{e}_z), \quad (\text{S6})$$




FIG. S1. Definition of the Euler angles (ϕ, θ, ψ) in the $z x z$ -convention with the swimmer orientation $\mathbf{e}_3 \equiv \mathbf{p}$ and $\mathbf{e}_z \equiv \mathbf{e}$.

is given by [S3]

$$\mathbf{R} = \begin{pmatrix} \cos \phi \cos \psi - \sin \phi \cos \theta \sin \psi & -\cos \phi \sin \psi - \sin \phi \cos \theta \cos \psi & \sin \phi \sin \theta \\ \sin \phi \cos \psi + \cos \phi \cos \theta \sin \psi & -\sin \phi \sin \psi + \cos \phi \cos \theta \cos \psi & -\cos \phi \sin \theta \\ \sin \theta \sin \psi & \sin \theta \cos \psi & \cos \theta \end{pmatrix}. \quad (\text{S7})$$

The inverse transformation is given by

$$(\mathbf{e}_x \ \mathbf{e}_y \ \mathbf{e}_z) = \mathbf{R}^T \cdot (\mathbf{e}_1 \ \mathbf{e}_2 \ \mathbf{e}_3). \quad (\text{S8})$$

In particular, \mathbf{e}_z can be expressed in the swimmer frame as $\mathbf{e}_z = \sin \theta \sin \psi \mathbf{e}_1 + \sin \theta \cos \psi \mathbf{e}_2 + \cos \theta \mathbf{e}_3$.

The angular velocity of the swimmer can be expressed with the derivatives of the Euler angles as [S3]

$$\boldsymbol{\Omega} = (\dot{\phi} \sin \theta \sin \psi + \dot{\theta} \cos \psi) \mathbf{e}_1 + (\dot{\phi} \sin \theta \cos \psi - \dot{\theta} \sin \psi) \mathbf{e}_2 + (\dot{\phi} \cos \theta + \dot{\psi}) \mathbf{e}_3. \quad (\text{S9})$$

APPROXIMATE EXPRESSION FOR THE CRITICAL ANGULAR VELOCITY Ω_c^e

The critical angular velocity, at which the saddle-node bifurcation takes place in the dynamical system determined by Eq. (12) from the main text can be derived analytically in the limit $|\lambda| \ll 1$. The fixed points $(\bar{\theta}, \bar{\psi})$ correspond to the solutions of equations $\dot{\theta} = 0$ and $\dot{\psi} = 0$, leading to the conditions

$$0 = \frac{3g_2(\lambda)B_2}{10a} \sin(2\bar{\theta}) + \Omega^e \sin \gamma \cos \bar{\psi}, \quad (\text{S10})$$

$$0 = \frac{3g_1(\lambda)B_2}{5a} \cos \bar{\theta} + \Omega^e \left(\cos \gamma - \frac{\sin \gamma \sin \bar{\psi}}{\tan \bar{\theta}} \right). \quad (\text{S11})$$

Because $g_2(\lambda) \ll g_1(\lambda)$, Eqs. (S10) and (S11) can only be solved simultaneously if $\cos \bar{\psi} \approx 0$. With $B_2 < 0$ and $\gamma < \pi/2$ (the situation shown in Fig. 3 in the main text), the saddle-node bifurcation takes place between two fixed points at $\bar{\psi} \approx 3\pi/2$, whereas the solution $\bar{\psi} \approx \pi/2$ gives a single spiral source. We therefore choose $\bar{\psi} = 3\pi/2$ to analyze the bifurcation. Equation (S11) then simplifies to

$$\Omega^e(\bar{\theta}) = -\frac{3\lambda B_2}{40a} \frac{\sin(2\bar{\theta})}{\sin(\gamma + \bar{\theta})}. \quad (\text{S12})$$




FIG. S2. Numerical solution for the critical angular velocity. The rescaled critical angular velocity Ω_c^e is calculated as a function of the odd viscosity ratio λ for three values of the angle γ . The solid lines represent the analytical approximation (S14).

For a fixed Ω^e , Eq. (S12) is solved by either 1 or 3 values of $\bar{\theta}$. The saddle-node bifurcation, at which 2 solutions vanish, corresponds to a local maximum of $\Omega^e(\bar{\theta})$. From the condition $\partial\Omega^e(\bar{\theta})/\partial\bar{\theta} = 0$, we obtain the position in phase space at which the bifurcation takes place

$$(\bar{\theta}, \bar{\psi}) = \left(\arctan\left(\tan^{1/3}\gamma\right), \frac{3}{2}\pi \right), \quad (\text{S13})$$

which gives the critical value

$$\Omega_c^e = -\frac{3\lambda B_2}{20a} (\sin^{2/3}\gamma + \cos^{2/3}\gamma)^{-3/2}, \quad (\text{S14})$$

for $0 < \gamma < \pi/2$. Expression (S14) has a minimum at $\gamma = \pi/4$. The lowest possible critical angular frequency is $\Omega_c^e = (3/2)|\lambda B_2|/(20a)$.

For larger λ , the critical angular velocity can be determined numerically by analyzing the number of fixed points in the dynamical system. Figure S2 shows Ω_c^e as a function of λ for three values of γ . The analytical expression is exact in the limit $\lambda \ll 1$ and the deviation stays within a few percent up to $\lambda = 1$.

* yuto.hosaka@ds.mpg.de

† ramin.golestanian@ds.mpg.de

‡ andrej.vilfan@ds.mpg.de

[S1] S. Ganeshan and A. G. Abanov, Odd viscosity in two-dimensional incompressible fluids, *Phys. Rev. Fluids* **2**, 094101 (2017).

[S2] T. M. Squires and M. Z. Bazant, Breaking symmetries in induced-charge electro-osmosis and electrophoresis, *J. Fluid Mech.* **560**, 65 (2006).

[S3] H. Goldstein, C. Poole, and J. Safko, *Classical Mechanics*, 3rd ed. (Pearson, Boston MA, 2001).