

<i>geodesic equation</i> approach (MTW, 1973)	<i>energy integral</i> approach (Weinberg, 1972)	hand-waving argument	new heuristic approach
<p>Hamilton-Jacobi equation in Schwarzschild geometry</p> <p>or</p> <p>null geodesic equation</p> <p>↓</p> <p>(angular momentum conservation is considered)</p> <p>↓</p> <p>Relativistic version of Binet's equation</p> <p>↓</p> <p>(perturbative expansion)</p> <p>↓</p> <p>trajectory equation obtained (first-order)</p> <p>↓</p> <div> $\theta = \frac{4GM}{c^2 R}$ </div>	<p>Schwarzschild metric</p> <p>↓</p> <p>Christoffel symbols</p> <p>↓</p> <p>geodesic equation</p> <p>↓</p> <p>energy integral for radial motion with effective potential</p> <p>↓</p> <p>trajectory equation obtained</p> <p>↓</p> <p>integral expression for θ</p> <p>↓</p> <p>(by expanding in powers before integration)</p> <p>↓</p> <p>Solvable integral form obtained</p> <p>↓</p> <div> $\theta = \frac{4GM}{c^2 R}$ </div>	<p>Newtonian attraction</p> <p>or</p> <p>equivalence principle</p> <p>↓</p> <p>(calculating a change in velocity)</p> <p>↓</p> <div> $\theta = \frac{2GM}{c^2 R}$ </div> <p>↓</p> <p>(by choosing the parameter arbitrarily)</p> <p>↓</p> <div> $\theta = \frac{4GM}{c^2 R}$ </div>	<p>Geometrized Newtonian formulation</p> <p>↓</p> <p>Schwarzschild metric</p> <p>↓</p> <p>(applying the variational principle)</p> <p>↓</p> <p>Newtonian type of equation obtained (focusing on the acceleration)</p> <p>↓</p> <p>(calculating a change in velocity by simple integration)</p> <p>↓</p> <div> $\theta = \frac{4GM}{c^2 R}$ </div>

*We denoted the deflection angle as θ