

Elements of Eq. (10)

$\Phi_m(\cdot)$


$(\mathbf{x}_j, \alpha_{w_j})$

$y(t, \mathbf{x}_j)$


$\hat{y}_m(t)$

Step 1: Create n_x -variate gPC basis functions $\Phi_m(\cdot)$

univariate basis \mathbf{x}_1




use tensor product to estimate n_x -variate bases




$$\{\Phi_m(\tilde{\mathbf{x}})\}_{m=0}^{M-1} = \otimes \left\{ \prod_{i=1}^2 \phi_k(x_i) \right\}$$

univariate basis \mathbf{x}_2




Step 2: Find quadrature nodes \mathbf{x}_j and weights α_{w_j}

$q_2 = 3$ collocation nodes



$q_1 = 3$ collocation nodes

scaled $\alpha_{w_{q1}}$



tensor product of quadrature weights

$$\otimes \left\{ \prod_{i=1}^2 \alpha_{w_{q_i}}(x_i) \right\}$$

Step 3: Solve Q deterministic problems

for $j = 1 : Q$

solve deterministic problem j

end

Q solutions
 $y(t, \mathbf{x}) \in \mathbb{R}^Q$

Step 4: Estimate Statistics

Estimate statistics using Eq.(13)

