

Question

Solve the equation
 $(x - 1)^2 - 5(x - 1) + 4 = 0$.



Simplifying the given equation: $x^2 - 2x + 1 - 5x + 5 + 4 = 0$, which simplifies further to $x^2 - 7x + 10 = 0$. Applying the quadratic formula $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, we first calculate the discriminant: $b^2 - 4ac = 49 - 40 = 9$. Therefore, the solutions are $x = \frac{7 \pm 3}{2}$, yielding the roots $x = 2$ and $x = 5$.



Solution (reflective training)

Simplifying the given equation: $x^2 - 2x + 1 \dots$ [original solution omitted] \dots yielding the roots $x = 2$ and $x = 5$.
(Original Solution)

Let's revisit the solution and consider some techniques to simplify the problem-solving process.

Given the simplified equation $x^2 - 7x + 10 = 0$, we can use a method called **factorization** to quickly get the roots by checking the coefficients b and c . According to Vieta's formulas, for $a = 1$, the sum of two roots is $-b$ and their product is c . Thus, in this equation, $x_1 + x_2 = 7$ and $x_1 x_2 = 10$. Recognizing that 2 and 5 satisfies these conditions, we factorize the original equation as $(x - 2)(x - 5) = 0$, thereby quickly finding the roots.

Taking a step further, there's another trick to solve the original equation $(x - 1)^2 - 5(x - 1) + 4 = 0$: **substitution**. By letting $x - 1 = t$, we transform the equation into $t^2 - 5t + 4 = 0$. Applying the factorization method, this becomes $(t - 1)(t - 4) = 0$. Thus, $t = 1$ or 4 . Since $t = x - 1$, we find $x = 2$ or 5 .
(Reflection)



Question

The unknown number x satisfies
 $(7x + 1)(9x + 1) = 61$. Find x .

Let's simplify the equation as $63x^2 + 7x + 9x + 1 = 61 \Rightarrow 63x^2 + 16x - 60 = 0$. The Two roots can be calculated as $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, where $\sqrt{b^2 - 4ac} = \sqrt{16^2 - 4 \times 63 \times (-60)} = \dots?$



Inference (after standard training)

Let's use substitution to solve the equation. To create a common structure for easier substitution, we multiply out the expressions: $(7x + 1) \times 9$ and $(9x + 1) \times 7$, yielding $(63x + 9)(63x + 7) = 61 \times 63$. Observing both sides, we notice that by letting $t = 63x + 8$, the equation can be rewritten as $(t + 1)(t - 1) = (62 + 1)(62 - 1)$, which simplifies to $t^2 = 62^2 \Rightarrow t = \pm 62$. Therefore, $x = \frac{6}{7}$ or $x = -\frac{10}{9}$.



Inference (after reflective training)