

Training Question: Solve the equation $(x - 99)(x - 101) = 8$.

Standard Training

Let $t = x - 100$. Then the equation becomes $(t - 1)(t + 1) = 8 \Rightarrow t^2 - 1 = 8$. Therefore, $t = \pm 3$, and accordingly we get $x = 97$ or 103 .

Training with Reflective Augmentation

Let $t = x - 100$ [omitted, same as standard training] ... we get $x = 97$ or 103 .

(Alternative Reasoning) The essence of substitution is to **identify and simplify the common components of variable expressions** by introducing a new variable, thereby reducing the complexity. Let's revisit the original equation. Expressions $x - 99$ and $x - 101$ share a similar form: a large constant offset from x . Due to the minimal difference between 99 and 101, we can use substitution to transform the expressions into terms with small constants.

Therefore, an alternative approach is to substitute $t = x - 99$, which transforms the equation into $t(t - 2) = 8 \Rightarrow t^2 - 2t - 8 = 0$. This can be easily factorized into $(t - 4)(t + 2) = 0$. Hence, $t = 4$ or -2 , leading to the same results $x = 97$ or 103 . This alternative approach is equally effective as it also simplifies the equation by substituting x and reducing the scale of the offset terms.

(Follow-up Reasoning) Extending the idea of substitution, consider the equation $x(x + 1)(x + 2)(x + 3) = 360$. We notice that $x(x + 3) = x^2 + 3x$, and $(x + 1)(x + 2) = x^2 + 3x + 2$. Therefore, to simplify the expression, we set the common term $x^2 + 3x$ as t , which transforms the equation into $t(t + 2) = 360 \Rightarrow t^2 + 2t - 360 = 0 \Rightarrow t = -20$ or 18 . If $t = -20$, then $x^2 + 3x + 20 = 0$. Here, the discriminant $\Delta = -71 < 0$, resulting in no real solutions for x . If $t = 18$, then $x^2 + 3x - 18 = 0$, so $x = 3$ or -6 . This scenario reiterates the importance of identifying common components of x to streamline the equation through substitution.

Test Question: The unknown number x satisfies $(7x + 1)(9x + 1) = 61$. Find x .

After Standard Training

Let $t = 8x + 1$. The equation becomes $(t - x + 1)(t + x + 1) = 61$. Next, we...???
If we set $t = 7x + 1$, the original equation turns into $t(t + 2x + 1) = 61$. Then we can get...???

After Reflective Augmentation

Let's use substitution to solve the equation. **To create a common structure for easier substitution**, we multiply the expressions to unify the coefficient of x : $(7x + 1) \times 9$ and $(9x + 1) \times 7$, yielding $(63x + 9)(63x + 7) = 61 \times 63$. Observing the equation, we notice that by letting $t = 63x + 8$, it can be rewritten as $(t + 1)(t - 1) = (62 + 1)(62 - 1)$, which simplifies to $t^2 = 62^2 \Rightarrow t = \pm 62$. Therefore, $x = \frac{6}{7}$ or $x = -\frac{10}{9}$. (Early stopping: no reflection needed during inference)