

Train

Question

Solve the equation
 $(x - 1)^2 - 5(x - 1) + 4 = 0$.



Solution (standard training)

Simplifying the given equation: $x^2 - 2x + 1 - 5x + 5 + 4 = 0$, which simplifies further to $x^2 - 7x + 10 = 0$. Applying the quadratic formula $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, we first calculate the discriminant: $b^2 - 4ac = 49 - 40 = 9$. Therefore, the solutions are $x = \frac{7 \pm 3}{2}$, yielding the roots $x = 2$ and $x = 5$.



Solution (with reflective augmentation)

Simplifying the given equation: $x^2 - 2x + 1 \dots$ [omitted, same as above] ... yielding the roots $x = 2$ and $x = 5$.

[Original Solution]

[Methodological reflection] Let's revisit the problem and find alternative ways to simplify the solution. We can use a method called **substitution** trick to solve the original equation $(x - 1)^2 - 5(x - 1) + 4 = 0$. By letting $x - 1 = t$, we transform the equation into $t^2 - 5t + 4 = 0$. Solving this equation leads to $t = 1$ or 4 . Since $t = x - 1$, we find $x = 2$ or 5 .

[Knowledge Extension] Extending the idea of substitution, let's consider solving $(x - 2021)(x - 2023) = 8$. Directly applying the quadratic formula results in complicated calculation. Thus, considering the symmetry of terms on the left, we let $t = x - 2022$, which transforms the equation to $(t + 1)(t - 1) = 8 \Rightarrow t^2 - 1 = 8$. Therefore, $t = \pm 3$, and accordingly we get $x = 2025$ or 2019 .

[Reflection]



Test

Question

The unknown number x satisfies
 $(7x + 1)(9x + 1) = 61$. Find x .

Let's simplify the equation as $63x^2 + 7x + 9x + 1 = 61 \Rightarrow 63x^2 + 16x - 60 = 0$. The Two roots can be calculated as $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, where $\sqrt{b^2 - 4ac} = \sqrt{16^2 - 4 \times 63 \times (-60)} = \dots$?



Inference (after standard training)

Let's use substitution to solve the equation. To create a common structure for easier substitution, we multiply out the expressions: $(7x + 1) \times 9$ and $(9x + 1) \times 7$, yielding $(63x + 9)(63x + 7) = 61 \times 63$. Observing the equation, we notice that by letting $t = 63x + 8$, it can be rewritten as $(t + 1)(t - 1) = (62 + 1)(62 - 1)$, which simplifies to $t^2 = 62^2 \Rightarrow t = \pm 62$. Therefore, $x = \frac{6}{7}$ or $x = -\frac{10}{9}$.



Inference (after reflective training)