

Based on the medical history and imaging features you provided, I will answer your questions in three steps.

Step One: Analysis and Recommendations Based on Medical History:

1. Tumor Marker AFP: The patient's Alpha-fetoprotein (AFP) level is 56448.33 ng/mL, which is above normal, suggesting the possibility of liver cancer. Combined with PIVKA-I and CA-199 levels, this also supports the diagnosis of hepatocellular carcinoma.
2. Portal Vein Tumor Thrombus: Imaging features show that the grayscale difference between the tumor and surrounding tissues is low, indicating a significant difference in blood supply between the tumor and surrounding tissues, suggesting the presence of a portal vein tumor thrombus.
3. Treatment Plan: The patient is currently undergoing neoadjuvant therapy with oral Lenvatinib and Sindilizumab. For hepatocellular carcinoma, surgery, radiation therapy, and chemotherapy are all possible treatment options. Given the patient's elevated AFP, surgical resection should be considered. However, it is important to note that due to the large size of the tumor, which may involve surrounding tissues and blood vessels, the surgical risk is high.

Step Two: Analysis of Imaging Features:

1. Tumor Diameter: 12.1 cm, classified as a large hepatocellular carcinoma.
2. Tumor Edge Shape: Irregular, indicating a high possibility of tumor aggressiveness.
3. Grayscale Differences Between Tumor and Surrounding Tissues: Low, indicating little difference in blood supply between the tumor and surrounding tissues.
4. Proximity of Tumor to Major Blood Vessels: Close, which requires careful consideration of the impact on blood vessels during surgery and radiation therapy.

Step Three: Treatment Recommendations:

Due to the large size of the tumor, surgical resection might be the better treatment option. However, during the surgery, attention should be paid to the impact on blood vessels, and procedures such as angiography may be necessary to assess the vascular situation. For the portal vein tumor thrombus, close monitoring of its changes is necessary, and consideration for thrombectomy or portal vein ligation may be required. Post-surgery, regular follow-ups are needed to monitor tumor markers and changes in imaging studies. The treatment plan should be adjusted based on the patient's specific circumstances, following EASL or mRECIST criteria.