

Firstly, the patient's medical records indicate that the main symptoms are pain under the xiphoid process and in the upper right abdomen, without any obvious cause. An abdominal imaging examination conducted at a local hospital revealed a liver space-occupying lesion and a defect in the filling of the portal vein. Further consultation at your hospital's Department of Hepatobiliary Surgery suggested a right-lobe liver lesion, suspected to be hepatocellular carcinoma, accompanied by portal vein tumor thrombus. Tumor marker results showed an Alpha-fetoprotein (AFP) level of 56448.33 ng/mL, suggesting the possibility of hepatocellular carcinoma due to the elevated AFP.

The imaging features show that the tumor has a diameter of 12.1 cm, is irregularly shaped, has low grayscale differences with surrounding tissues, has significant internal grayscale variations, and is located close to major blood vessels. These characteristics may suggest rapid tumor growth, blurred interfaces with surrounding tissues, and potentially severe vascular invasion.

Combining the patient's medical records and imaging features, it is likely that the patient has hepatocellular carcinoma in the right lobe of the liver, accompanied by portal vein tumor thrombus and metastasis. The rapid growth of the tumor may have a significant impact on surrounding tissues and blood vessels. The presence of portal vein tumor thrombus indicates that the tumor has invaded the portal vein, which may be associated with portal hypertension. The patient has started oral Lenvatinib and Sindilizumab for neoadjuvant therapy. Based on the patient's condition and changes in tumor markers, further imaging studies and laboratory tests might be needed to assess the treatment effectiveness and prognosis.

The above is a preliminary analysis and interpretation of the patient's medical and imaging information. Specific diagnosis and treatment should be determined in conjunction with clinical circumstances and other test results.