


Question: Is there any infiltration of the lesser curvature of the stomach or duodenum?



Answer 1: According to the axial abdominal CT scan, there is no infiltration of the lesser curvature of the stomach or duodenum.

Answer 2: No, the CT scan does not show any infiltration of the lesser curvature of the stomach or duodenum.



Answer 1



Answer 2

Question: What could be the possible cause of bilateral infiltrates?



Answer 1: Bilateral infiltrates on a chest X-ray can be caused by various conditions, including infections (such as pneumonia), inflammation, or other lung diseases. Some possible causes are bacterial, viral, or fungal infections, aspiration of foreign material, or autoimmune disorders. It is important to consider the patient's clinical history, symptoms, and additional diagnostic tests to determine the specific cause of the bilateral infiltrates and provide appropriate treatment.

Answer 2: Bilateral infiltrates on a chest X-ray can be caused by various factors, such as infections (like pneumonia), inflammation, heart failure, or lung diseases.



Answer 1



Answer 2

Question: What kind of lesion is it?



Answer 1: The lesion appears to be a multilocular, cystic retroperitoneal tumor. It is a fluid-filled mass that is situated in the retroperitoneal space, which is the area behind the peritoneum, the membrane that lines the abdominal cavity. The lesion is divided into multiple septations, which are thin walls or partitions within the cystic structure.

Answer 2: The lesion appears to be a soft tissue heterogeneously enhancing lesion with a nodular appearance. It could be suggestive of a reactive lymph node or another type of lesion.



Answer 1



Answer 2