

Question: What are positively charged, thus allowing the compaction of the negatively charged DNA?

STLLaVA-Med w/o DPO

GT: the histone subunits

In the context of the image, **histones are positively charged**, which means they have a positive charge at physiological pH. This positive charge allows them to interact with the negatively charged DNA, helping to compact and organize the DNA into a more compact structure.



STLLaVA-Med


In the context of the image, **histones are positively charged**, which means they have a positive charge at physiological pH. This positive charge allows them to interact with the negatively charged DNA, helping to compact and organize the DNA into a more compact structure. Histones are essential components of chromatin, which is the complex of DNA and histones that makes up the nucleosome, a fundamental unit of chromosome organization.

Question: What does the wall of the artery show with protein deposition and inflammation?

STLLaVA-Med w/o DPO

GT: a circumferential bright pink area of necrosis

In the context of this image, the presence of protein deposition and inflammation in the arterial wall may indicate an underlying vascular condition or injury that requires further investigation and appropriate treatment.



STLLaVA-Med

In the histopathological image, the wall of the artery shows protein deposition and inflammation. Protein deposition refers to the accumulation of proteins in the arterial wall, which can be a sign of injury, inflammation, or other pathological processes. Inflammation is the body's response to injury or infection, and it can cause redness, swelling, and pain in the affected area. In the context of this image, the presence of protein deposition and inflammation in the arterial wall may indicate an underlying vascular condition or injury that requires further investigation and appropriate treatment.
