

Supplemental Material for “Hilbert band complexes and their applications”

Zeying Zhang,^{1,*} Y. X. Zhao,² Yugui Yao,³ and Shengyuan A. Yang⁴

¹*College of Mathematics and Physics, Beijing University of Chemical Technology, Beijing 100029, China*

²*Department of Physics and HKU-UCAS Joint Institute for Theoretical and Computational Physics at Hong Kong, The University of Hong Kong, Pokfulam Road, Hong Kong, China*

³*Centre for Quantum Physics, Key Laboratory of Advanced Optoelectronic Quantum Architecture and Measurement (MOE), Beijing Institute of Technology, Beijing 100081, China*

⁴*Research Laboratory for Quantum Materials, IAPME, FST, University of Macau, Macau, China*

CONTENTS

S1. Notations	2
S2. Some applications of Hilbert basis in periodic systems	2
S3. Examples	3
S4. 133 kinds of bipartite graph 230 space groups	10
S5. Quantitative mapping between bipartite graph and 230 space groups	21
S6. Hilbert basis and bipartite graph in 230 space groups without SOC	26
S7. Hilbert basis and bipartite graph in 230 space groups with SOC	883
References	1632

* zzy@mail.buct.edu.cn

S1. NOTATIONS

In Sec. S4, we list all the possible bipartite graph 230 space groups. Each bipartite graph is labelled by $H_{m/n}^i$, where m is the number of vertex and n is the number of edge in the bipartite graph, i is the number to distinguish the bipartite graph with same number of vertex and edge but not isomorphic to each other.

In Sec. S5 A and S5 B, we list the quantitative mapping between bipartite graph and 230 space groups with and without SOC. The first column gives the label of bipartite graph, the second column gives the SG number which can host the bipartite graph.

In Sec. S6 and S7, we list the Hilbert basis for each high symmetry line in single and double SG, including the compatibility relations, balance band matrix, Hilbert basis for $\ker M \cap \mathbb{Z}_{\geq 0}^{m+n}$ and the bipartite graph(s) for each Hilbert basis. Notice the labels of IRRs and the names of high symmetry points in this work are from Bradley and Cracknell's (BC) book [1], which have been implemented in `SpaceGrouprep` and `MSGCorep` packages [2, 3]. Therefore the labels of IRRs and the names of high symmetry points may different from Bilbao Crystallographic Server (BCS) [4]. One can find the correspondence between BC and BCS labels in Supplemental Material of Ref. [2].

We use one table to represent the compatibility relations and balance band matrix. The M matrix in this section is shown as

corep		P_1	P_2	...	P_m	P_{m+1}	P_{m+1}	...	P_{n+m}
	dim	$d(P_1)$	$d(P_2)$...	$d(P_m)$	$d(P_{m+1})$	$d(P_{m+2})$...	$d(P_{m+n})$
L_1	$d(L_1)$	a_1^1	a_1^2	...	a_1^m	$-a_1^{m+1}$	$-a_1^{m+2}$...	$-a_1^{m+n}$
L_2	$d(L_2)$	a_2^1	a_2^2	...	a_2^m	$-a_2^{m+1}$	$-a_2^{m+2}$...	$-a_2^{m+n}$
...	
L_s	$d(L_s)$	a_s^1	a_s^2	...	a_s^m	$-a_s^{m+1}$	$-a_s^{m+2}$...	$-a_s^{m+n}$

where $d(P_i)$ is the dimension of P_i , and we have the compatibility relations $P_i \downarrow L = \bigoplus_j^s a_i^j L_j$. The kernel of M is the null space of

$$\begin{pmatrix} a_1^1 & a_1^2 & \dots & a_1^m & -a_1^{m+1} & -a_1^{m+2} & \dots & -a_1^{m+n} \\ a_2^1 & a_2^2 & \dots & a_2^m & -a_2^{m+1} & -a_2^{m+2} & \dots & -a_2^{m+n} \\ & & \dots & & & & \dots & \\ a_s^1 & a_s^2 & \dots & a_s^m & -a_s^{m+1} & -a_s^{m+2} & \dots & -a_s^{m+n} \end{pmatrix}$$

The Hilbert basis of $\ker M \cap \mathbb{Z}_{\geq 0}^{m+n}$ is represented by

	P_1	P_2	...	P_m	P_{m+1}	P_{m+2}	...	P_{n+m}	
	$d(P_1)$	$d(P_2)$...	$d(P_m)$	$d(P_{m+1})$	$d(P_{m+2})$...	$d(P_{m+n})$	bipartite graph
v^1	v_1^1	v_2^1	...	v_m^1	v_{m+1}^1	v_{m+2}^1	...	v_{m+n}^1	H_1
v^2	v_1^2	v_2^2	...	v_m^2	v_{m+1}^2	v_{m+2}^2	...	v_{m+n}^2	H_2
...									
v^h	v_1^h	v_2^h	...	v_m^h	v_{m+1}^h	v_{m+2}^h	...	v_{m+n}^h	H_h

where v_i is the i -th Hilbert basis, gives exactly the number of occurrences v_i^j of each IRR for P_j . H_i is the label of bipartite graph shown in S4.

S2. SOME APPLICATIONS OF HILBERT BASIS IN PERIODIC SYSTEMS

Hilbert basis build basic blocks of band connectivity in periodic system. In this section, we introduce some applications to show that Hilbert basis is a powerful tool to study the topological matter. To this end, we first introduce the method to calculate the Hilbert basis over the entire Brillouin zone (BZ).

To calculate the Hilbert basis globally, one need simply calculate the Hilbert basis of

$$S = \left(\bigcap_i \ker M_i \right) \cap \mathbb{Z}_{\geq 0}^d$$

where M_i is the balance band matrix for high symmetry line numbered by i , d is the total number of IRRs on each end of high symmetry lines. Since Hilbert basis of intersection of linear space and non-negative integer can be obtained by `Normaliz` package [5]. Therefore, the main task is to calculate $\bigcap_i \ker M_i$, this can be also easily done by Gaussian elimination or iterative simplification algorithm [6]. Thus, we can use `Normaliz` package to calculate $\text{Hilb}(\left(\bigcap_i \ker M_i\right) \cap \mathbb{Z}_{\geq 0}^d)$.

However, It is too redundant to write all the Hilbert bases of all SGs here. For example, the number of global Hilbert bases of space group $P/6mmm$ (No. 191) is 240. Therefore, we present a Mathematica script at <https://github.com/zhangzeyingvv/> to calculate and show all the global Hilbert bases for 230 SGs.

All the band representations with a set of isolated bands can be written as the positive integer combination of elements in $\text{Hilb}(S)$. Mathematically, the each coefficient of representations in the set of band is an affine monoid:

$$\text{Aff} \langle \text{HBR} \rangle_{\mathbb{Z}_{\geq 0}} = \{a_1x_1 + a_2x_2 + \dots + a_nx_n | a_i \in \mathbb{Z}_{\geq 0}, x_i \in \text{HBR}\}$$

which is different form linear space spanned by $\text{Hilb}(S)$:

$$\text{Span} \langle \text{HBR} \rangle_{\mathbb{Z}} = \{a_1x_1 + a_2x_2 + \dots + a_nx_n | a_i \in \mathbb{Z}, x_i \in \text{HBR}\}.$$

With the help of Hilbert basis, the definition and their relationships of Hilbert band representation (HBR), elementary band representations (EBR), band insulators (BI), atomic insulators (AI), filling-enforced quantum band insulators (QBI), fragile topology (FT), stable topology (ST), and obstructed atomic insulator (OAI) can be simply written as

$$\begin{aligned} \{\text{BI}\} &= \text{Aff} \langle \text{HBR} \rangle_{\mathbb{Z}_{\geq 0}} \\ \{\text{AI}\} &= \text{Aff} \langle \text{EBR} \rangle_{\mathbb{Z}_{\geq 0}} \\ \{\text{QBI}\} &= \{\text{BI}\} - \{\text{AI}\} \\ \{\text{FT}\} &= \text{Span} \langle \text{HBR} \rangle_{\mathbb{Z}} \cap \{\text{BI}\} - \{\text{AI}\} \\ \{\text{ST}\} &= \{\text{BI}\} - \text{Span} \langle \text{EBR} \rangle_{\mathbb{Z}} \\ \{\text{OAI}\} &= \{\text{AI}\} - \text{Aff} \langle \text{EBR}_p \rangle_{\mathbb{Z}_{\geq 0}} \end{aligned} \tag{S1}$$

where $A - B$ is the relative complement of B in A . and EBR_p is restricted only to the atomic-occupied Wyckoff positions. Notice the above definition is more general than the symmetry-based indicators since symmetry-based indicator in Ref. [7] treats $\{\text{BI}\}$ and $\{\text{AI}\}$ as linear spaces rather than affine monoids. In Sec. S3, we give an SG example which have trivial symmetry-based indicator but have non-trivial topological band structure.

Another application of Hilbert basis is the generalization the theory of filling-enforced gaplessness in band structures. Consider a set of isolated bands generated by $\{\text{BI}\}$ or $\{\text{AI}\}$. The numbers of bands are also affine monoids (numerical monoids):

$$N_h = \{a_1n(x_1) + a_2n(x_2) + \dots + a_n n(x_n) | a_i \in \mathbb{Z}_{\geq 0}, x_i \in \text{HBR}\}$$

$$N_e = \{a_1n(x_1) + a_2n(x_2) + \dots + a_n n(x_n) | a_i \in \mathbb{Z}_{\geq 0}, x_i \in \text{EBR}\}$$

where $n(x_i)$ represent the number of bands for x_i . Notice the above affine monoids not only can apply to the entire BZ, but also can apply to arbitrary continuous curves or planes in BZ.

S3. EXAMPLES

In this section, we give an example to show how to use Hilbert basis to study the filling-enforced gapless point and topology band structure in SG $P4_232$ (No. 208) and then we give two lattice models to illustrate the construction of Hilbert particles in realistic systems. Beside, we also give the specific usages of S6-S7 through the examples. The first example is the construction of H_{878}^2 (as shown in S4) in double space group (SG) $P4_1$ (No. 76). The second example is the construction of $H_{5/5}^2$ and $H_{5/5}^3$ in single SG $Pa\bar{3}$ (No. 205) which correspond to the same Hilbert basis. The source code to generate two models can be found in <https://github.com/zhangzeyingvv/MagneticTB> [8].

A. Lattice model for $H_{8/8}^1$ in double SG $P4_1$

$P4_1$ have only one Wyckoff position $4a$. Consider the simplest case in presence of spin-orbital coupling (SOC) : put $|s \uparrow\rangle$ and $|s \downarrow\rangle$ orbitals on each site of $4a$. Then the lattice Hamiltonian can be established as

$$H_{76} = t \begin{pmatrix} 0 & H \\ H^\dagger & 0 \end{pmatrix}$$

where

$$H = \begin{pmatrix} -ie^{-\frac{ik_z}{4}} & -\lambda e^{\frac{ik_z}{4}} & ie^{-\frac{ik_z}{4}} & i\lambda e^{-\frac{ik_z}{4}} \\ \lambda e^{\frac{ik_z}{4}} & ie^{\frac{ik_z}{4}} & i\lambda e^{-\frac{ik_z}{4}} & -ie^{-\frac{ik_z}{4}} \\ ie^{-\frac{ik_z}{4}} & -i\lambda e^{-\frac{ik_z}{4}} & -ie^{\frac{ik_z}{4}} & \lambda e^{\frac{ik_z}{4}} \\ -i\lambda e^{-\frac{ik_z}{4}} & -ie^{-\frac{ik_z}{4}} & -\lambda e^{\frac{ik_z}{4}} & ie^{\frac{ik_z}{4}} \end{pmatrix}$$

t and λ are real parameters in the lattice model. One can find for $\lambda t \neq 0$, the band structure always form the shape of $H_{8/8}^1$.

In fact, there is only one possibility for the shape of band structure along $\Gamma - Z$. We can confirm it by check Λ line of double SG $P4_1$ in S7. The M matrix in S7 is shown as

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_5	Z_6Z_8	Z_7Z_7
	dim	2	2	2	2	2
Λ_5	1	1	0	-2	0	0
Λ_6	1	0	1	0	-1	0
Λ_7	1	0	1	0	0	-2
Λ_8	1	1	0	0	-1	0

Here, ‘‘corep’’ and ‘‘dim’’ are the labels and dimension of irreducible small co-representations (IRRs) (row for the high symmetry points and column for high symmetry line). Each number in the matrix is the number of occurrences of IRRs of high symmetry line in IRRs of high symmetry point. That is, the above table can be equivalently written as

	$\Gamma_5\Gamma_8 \downarrow \Lambda$	$\Gamma_6\Gamma_7 \downarrow \Lambda$	$Z_5Z_5 \downarrow \Lambda$	$Z_6Z_8 \downarrow \Lambda$	$Z_7Z_7 \downarrow \Lambda$
$\chi^{\Gamma(Z)} \downarrow \Lambda$	$\Lambda_5 \oplus \Lambda_8$	$\Lambda_6 \oplus \Lambda_7$	$2\Lambda_5$	$\Lambda_6 \oplus \Lambda_8$	$2\Lambda_7$

where $\chi^{\Gamma(Z)}$ are the IRRs of $\Gamma(Z)$ point. The kernel of M is just the null space of

$$\begin{pmatrix} 1 & 1 & 0 & -2 & 0 & 0 \\ 1 & 0 & 1 & 0 & -1 & 0 \\ 1 & 0 & 1 & 0 & 0 & -2 \\ 1 & 1 & 0 & 0 & -1 & 0 \end{pmatrix}$$

The Hilbert basis and corresponding bipartite graph of $\ker M \cap \mathbb{Z}_+^5$ can also be found in S7:

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_5	Z_6Z_8	Z_7Z_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	2	1	$H_{8/8}^1$

One can find there is only one Hilbert basis, indicate the above 8-band model have to be formed $H_{8/8}^1$ shape of band structure on Λ (as shown in Fig. S1). The band IRs for the model with parameters $t = \lambda = 1$ are

\mathbf{k}	bands	energy	degeneracy	corep
$\Gamma(0, 0, 0)$	7 – 8	2.73	2	$\Gamma_6\Gamma_7$
	5 – 6	0.73	2	$\Gamma_5\Gamma_8$
	3 – 4	-0.73	2	$\Gamma_6\Gamma_7$
	1 – 2	-2.73	2	$\Gamma_5\Gamma_8$
$\Lambda(0, 0, \frac{1}{4})$	8	2.80	1	Λ_6
	7	2.24	1	Λ_7
	6	1.72	1	Λ_8
	5	0.37	1	Λ_7
	4	-0.37	1	Λ_5
	3	-1.72	1	Λ_6
	2	-2.24	1	Λ_5
	1	-2.80	1	Λ_8
$Z(0, 0, \frac{1}{2})$	7 – 8	2.45	2	Z_6Z_8
	5 – 6	1.41	2	Z_7Z_7
	3 – 4	-1.41	2	Z_5Z_5
	1 – 2	-2.45	2	Z_6Z_8

which consistent with the Hilbert basis.

B. Lattice model for $H_{5/5}^2$ and $H_{5/5}^3$ in single SG $Pa\bar{3}$

In this example, we show that two graphs that are not isomorphic to each other can correspond to a same Hilbert basis. Consider the $|d_z^2\rangle$ and $|d_{x^2-y^2}\rangle$ orbitals locate at $4a$ Wyckoff position of single SG $Pa\bar{3}$. Then the lattice Hamiltonian can be established as

$$H_{205} = t \begin{pmatrix} H_{11} & H_{12} \\ H_{12}^\dagger & H_{22} \end{pmatrix}$$

where

$$H_{11} = \begin{pmatrix} 0 & 0 & 4 \cos \frac{k_x}{2} \cos \frac{k_y}{2} & 2\lambda \cos \left(\frac{k_x+k_y}{2} \right) \\ 0 & 0 & 2\lambda \cos \left(\frac{k_x-k_y}{2} \right) & 0 \\ 4 \cos \frac{k_x}{2} \cos \frac{k_y}{2} & 2\lambda \cos \left(\frac{k_x-k_y}{2} \right) & 0 & 0 \\ 2\lambda \cos \left(\frac{k_x+k_y}{2} \right) & 0 & 0 & 0 \end{pmatrix},$$

$$H_{22} = \begin{pmatrix} 0 & 0 & 4 \cos \frac{k_x}{2} \cos \frac{k_y}{2} & 2\lambda \cos \left(\frac{k_x-k_y}{2} \right) \\ 0 & 0 & 2\lambda \cos \left(\frac{k_x+k_y}{2} \right) & 0 \\ 4 \cos \frac{k_x}{2} \cos \frac{k_y}{2} & 2\lambda \cos \left(\frac{k_x+k_y}{2} \right) & 0 & 0 \\ 2\lambda \cos \left(\frac{k_x-k_y}{2} \right) & 0 & 0 & 0 \end{pmatrix},$$




FIG. S1. (a) Band structure along $\Gamma - \Lambda$ of SG $P4_1$ lattice model with parameters $t = \lambda = 1$. (b)-(c) Band structure along $\Gamma - \Lambda$ of SG $Pa\bar{3}$ lattice model with parameters $t = 0.5, \lambda = \pm 0.2$. (d)-(f) The bipartite graphs correspond to (a)-(c).

$$H_{12} = \begin{pmatrix} (\sqrt{3}\lambda + 1) C_{yz} & -2\lambda S_{yz} - (\lambda + \sqrt{3}) C_{yz} & (1 - \sqrt{3}\lambda) C_{xz} & (\sqrt{3} - \lambda) C_{xz} - 2\lambda S_{xz} \\ 2\lambda S_{yz} - (\lambda + \sqrt{3}) C_{yz} & (3 - \sqrt{3}\lambda) C_{yz} & 2\lambda S_{xz} + (\sqrt{3} - \lambda) C_{xz} & (\sqrt{3}\lambda + 3) C_{xz} \\ (1 - \sqrt{3}\lambda) C_{xz} & 2\lambda S_{xz} + (\sqrt{3} - \lambda) C_{xz} & (\sqrt{3}\lambda + 1) C_{yz} & -2\lambda S_{yz} - (\lambda + \sqrt{3}) C_{yz} \\ (\sqrt{3} - \lambda) C_{xz} - 2\lambda S_{xz} & (\sqrt{3}\lambda + 3) C_{xz} & 2\lambda S_{yz} - (\lambda + \sqrt{3}) C_{yz} & (3 - \sqrt{3}\lambda) C_{yz} \end{pmatrix},$$

here, $C_{yz} = \cos \frac{k_y}{2} \cos \frac{k_z}{2}$, $S_{yz} = \sin \frac{k_y}{2} \sin \frac{k_z}{2}$, $C_{xz} = \cos \frac{k_x}{2} \cos \frac{k_z}{2}$, $S_{xz} = \sin \frac{k_x}{2} \sin \frac{k_z}{2}$, and t, λ are real parameters in the model. One can find that when $\lambda > 0$ the band structure form the shape of $H_{5/5}^3$ and when $\lambda < 0$ the band structure form the shape of $H_{5/5}^2$. We can look up Λ line of SG $Pa\bar{3}$ in S6 to confirm it. The compatibility relations and M matrix for Λ are

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_3^+	Γ_2^-	Γ_3^-	Γ_4^+	Γ_4^-	R_1^+	R_1^+	R_1^-	R_1^-	R_2^+	R_3^+	R_2^-	R_3^-
dim	1	1	2	2	3	3			4	4	4	4				
Λ_1	1	1	0	0	1	1			0	0	-2	-2				
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	1		-2	-2	-1	-1				

which is equivalent to

	$\Gamma_1^+ \downarrow \Lambda$	$\Gamma_1^- \downarrow \Lambda$	$\Gamma_2^+ \Gamma_3^+ \downarrow \Lambda$	$\Gamma_2^- \Gamma_3^- \downarrow \Lambda$	$\Gamma_4^+ \downarrow \Lambda$	$\Gamma_4^- \downarrow \Lambda$	$R_1^+ R_1^+ \downarrow \Lambda$	$R_1^- R_1^- \downarrow \Lambda$	$R_2^+ R_3^+ \downarrow \Lambda$	$R_2^- R_3^- \downarrow \Lambda$
$\chi^{\Gamma(R)} \downarrow \Lambda$	Λ_1	Λ_1	Λ_2	Λ_2	$\Lambda_1 \oplus \Lambda_2$	$\Lambda_1 \oplus \Lambda_2$	$2\Lambda_2$	$2\Lambda_2$	$2\Lambda_1 \oplus \Lambda_2$	$2\Lambda_1 \oplus \Lambda_2$

$\ker M \cap \mathbb{Z}_+^{10}$ have 80 Hilbert bases and the Hilbert basis for our lattice model is v_{53}

Γ_1^+	Γ_1^-	Γ_2^+	Γ_3^+	Γ_2^-	Γ_3^-	Γ_4^+	Γ_4^-	R_1^+	R_1^+	R_1^-	R_1^-	R_2^+	R_3^+	R_2^-	R_3^-
1	1	2	2	3	3	4	4	4	4	Bipartite graph					

v^1	0	0	0	0	0	4	0	1	0	2	$H_{7/8}^2, H_{7/8}^3$
v^2	0	0	0	0	0	4	0	1	1	1	$H_{7/8}^2, H_{7/8}^3$
...											
v^{56}	0	0	1	0	2	0	1	0	1	0	$H_{5/5}^2, H_{5/5}^3$
...											
v^{80}	2	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$

which means v_{56} can form two kinds of bipartite graph $H_{5/5}^2$ and $H_{5/5}^3$.
The band IRRs for the model with parameters $t = 0.5, \lambda = 0.2$ are

\mathbf{k}	bands	energy	degeneracy	corep
$\Gamma(0, 0, 0)$	7 – 8	3.00	2	$\Gamma_2^+ \Gamma_3^+$
	4 – 6	1.04	3	Γ_4^+
	1 – 3	-3.04	3	Γ_4^+
$\Lambda(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$	7 – 8	1.52	2	$\Lambda_2 \Lambda_3$
	6	0.69	1	Λ_1
	4 – 5	0.42	2	$\Lambda_2 \Lambda_3$
	3	-1.35	1	Λ_1
	1 – 2	-1.61	2	$\Lambda_2 \Lambda_3$
$R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$	5 – 8	0.35	4	$R_2^+ R_3^+$
	1 – 4	-0.35	4	$R_1^+ R_1^+$

and the band IRRs for the model with parameters $t = 0.5, \lambda = -0.2$ are

\mathbf{k}	bands	energy	degeneracy	corep
$\Gamma(0, 0, 0)$	7 – 8	3.00	2	$\Gamma_2^+ \Gamma_3^+$
	4 – 6	1.04	3	Γ_4^+
	1 – 3	-3.04	3	Γ_4^+
$\Lambda(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$	7 – 8	1.52	2	$\Lambda_2 \Lambda_3$
	5 – 6	0.60	2	$\Lambda_2 \Lambda_3$
	4	0.35	1	Λ_1
	2 – 3	-1.44	2	$\Lambda_2 \Lambda_3$
	1	-1.69	1	Λ_1
$R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$	5 – 8	0.35	4	$R_1^+ R_1^+$
	1 – 4	-0.35	4	$R_2^+ R_3^+$

both two cases consistent with the Hilbert basis v_{53} . Therefore, it is confirm that two non-isomorphic bipartite graphs can correspond to a same Hilbert basis (as shown in Fig. S1).

C. Materials realizing for complete bipartite graph $H_{5/6}^4$ in double SG $F\bar{4}_3c$

Based on the results presented in Secs. S4-S5, it found that only $H_{4/4}^2$ and $H_{5/6}^4$ are complete bipartite graph, where $H_{4/4}^2$ is the well know Hourglass Fermion. However, material with $H_{5/6}^4$ HBC have not been reported yet. The $H_{5/6}^4$




FIG. S2. (a) Lattice structure of Ge_3Cl_8 . (b) BZ with high symmetry points, line and plane labelled. (c) Band structure of Ge_3Cl_8 along high symmetry line. (d) Zoom in band structure of Ge_3Cl_8 along $\Gamma - L$ around Fermi energy, which has been marked within the green rectangles in (c). The band representations are also labelled.

bipartite graph can only appear on the Λ high symmetry line of the double space group $F\bar{4}_3c$. Here, by first-principles calculations, we confirm that Ge_3Cl_8 , which has been successfully synthesized [9], indeed host $H_{5/6}^4$ HBC around Fermi energy. One can observe the $H_{5/6}^4$ HBC along path $\Gamma - L$, as shown in Fig. S2(c)-(d). The calculation is done by using the *Vienna ab initio simulation package* (VASP) [10, 11]. The generalized gradient approximation with the Perdew-Burke-Ernzerhof (PBE) realization is adopted for the exchange-correlation functional [12]. The optimized lattice parameters are used in Materials Project database $a = b = c = 17.62 \text{ \AA}$ [13]. The cutoff energy was chosen as 500 eV, and BZ was sampled with Γ centred $3 \times 3 \times 3$ k -mesh. The band representations of Ge_3Cl_8 are calculated by *vasp2trace* and *SpaceGroupRep* packages [2, 14].

D. Wannier obstructions with trivial symmetry-based indicator in double SG $P4_232$

By putting each irreducible co-representation of site symmetry group in each Wyckoff position. We can list the 17 AIs in double SG $P4_232$:

	occ	Γ_6	Γ_7	Γ_8	X_6	X_7	M_6	M_7	R_6	R_7	R_8
$24m$	48	4	4	8	12	12	12	12	4	4	8
$12l$	24	2	2	4	6	6	6	6	2	2	4
$12k$	24	2	2	4	6	6	6	6	2	2	4
$12j$	24	2	2	4	6	6	6	6	2	2	4
$12i$	24	2	2	4	6	6	6	6	2	2	4




FIG. S3. (a) Band structure of $P4_232$ lattice model. (b) (111) surface states for the lattice model $P4_232$ lattice model.

$12h$	24	2	2	4	6	6	6	6	2	2	4
$8g$	16	0	0	4	4	4	4	4	0	0	4
$8g$	16	2	2	2	4	4	4	4	2	2	2
$6f$	12	1	1	2	3	3	3	3	1	1	2
$6e$	12	1	1	2	3	3	3	3	1	1	2
$6d$	12	1	1	2	3	3	3	3	1	1	2
$4c$	8	1	1	1	2	2	2	2	1	1	1
$4c$	8	0	0	2	2	2	2	2	0	0	2
$4b$	8	1	1	1	2	2	2	2	1	1	1
$4b$	8	0	0	2	2	2	2	2	0	0	2
$2a$	8	0	0	2	2	2	2	2	0	0	2
$2a$	4	1	1	0	1	1	1	1	1	1	0

where each row gives the occurrences of each irreducible co-representation. Notice 9 of them are EBR:

	occ	Γ_6	Γ_7	Γ_8	X_6	X_7	M_6	M_7	R_6	R_7	R_8
$6f$	12	1	1	2	3	3	3	3	1	1	2
$6e$	12	1	1	2	3	3	3	3	1	1	2
$6d$	12	1	1	2	3	3	3	3	1	1	2
$4c$	8	1	1	1	2	2	2	2	1	1	1
$4c$	8	0	0	2	2	2	2	2	0	0	2
$4b$	8	1	1	1	2	2	2	2	1	1	1
$4b$	8	0	0	2	2	2	2	2	0	0	2
$2a$	8	0	0	2	2	2	2	2	0	0	2
$2a$	4	1	1	0	1	1	1	1	1	1	0

By doing the Smith decomposition of 17 AI vectors. We can get 2 AI bases vectors a_1, a_2 :

	occ	Γ_6	Γ_7	Γ_8	X_6	X_7	M_6	M_7	R_6	R_7	R_8
a_1	8	1	1	1	2	2	2	2	1	1	1
a_2	4	1	1	0	1	1	1	1	1	1	0

which gives a trivial symmetry-based indicator. The Hilbert basis over the entire Brillouin zone can be obtained by calculate the kernel of the balance band matrix in globally:


	Γ_6	Γ_7	Γ_8	X_6	X_7	M_6	M_7	R_6	R_7	R_8
Δ_5	1	0	1	-1	0	0	0	0	0	0
Δ_6	0	1	1	-1	0	0	0	0	0	0
Δ_7	0	1	1	0	-1	0	0	0	0	0
Δ_8	1	0	1	0	-1	0	0	0	0	0
Σ_3	1	1	2	0	0	-1	-1	0	0	0
Σ_4	1	1	2	0	0	-1	-1	0	0	0
Λ_4	1	1	1	0	0	0	0	-1	-1	-1
Λ_5	1	1	1	0	0	0	0	-1	-1	-1
Λ_6	0	0	2	0	0	0	0	0	0	-2
S_3	0	0	0	1	1	0	0	-1	-1	-2
S_4	0	0	0	1	1	0	0	-1	-1	-2
Z_3	0	0	0	1	1	-1	-1	0	0	0
Z_4	0	0	0	1	1	-1	-1	0	0	0
T_5	0	0	0	0	0	0	1	0	-1	-1
T_6	0	0	0	0	0	1	0	0	-1	-1
T_7	0	0	0	0	0	1	0	-1	0	-1
T_8	0	0	0	0	0	0	1	-1	0	-1




Thus we get 2 Hilbert bases v^1, v^2 :




	occ	Γ_6	Γ_7	Γ_8	X_6	X_7	M_6	M_7	R_6	R_7	R_8
v^1	4	1	1	0	1	1	1	1	1	1	0
v^2	4	0	0	1	1	1	1	1	0	0	1




It clear that $v^1, v^2 \in \text{Span}\{\text{EBR}\}$, $v^1 = a_2 \in \{\text{AI}\}$, but $v^2 = a_1 - a_2 \notin \{\text{AI}\}$. Notice the AI bases can be arbitrary maximal linear independent integer vectors in AI. In a sharp contrast, the Hilbert bases is uniquely determined by the compatibility relations. Therefore, a set of isolated bands with coefficient of representations in v^1 is an FT. The lattice model for v^1 can be realized by putting $|d_{z^2} \uparrow / \downarrow\rangle$ and $|d_{x^2-y^2} \uparrow / \downarrow\rangle$ orbitals at $2a$ Wyckoff position. The band structure and (111) surface states are shown in Fig. S3(b),(d). One can find there exists four non-trivial suspend surface states around Fermi energy.




S4. 133 KINDS OF BIPARTITE GRAPH 230 SPACE GROUPS






 $H_{5/4}^2$ $H_{5/4}^3$ $H_{5/5}^1$ $H_{5/5}^2$ $H_{5/5}^3$  $H_{5/5}^4$ $H_{5/6}^1$ $H_{5/6}^2$ $H_{5/6}^3$ $H_{5/6}^4$  $H_{5/6}^5$ $H_{5/6}^6$ $H_{5/8}^1$ $H_{5/8}^2$ $H_{5/8}^3$








$H_{7/8}^9$

$H_{7/8}^{10}$

$H_{7/8}^{11}$

$H_{7/8}^{12}$

$H_{7/8}^{13}$




$H_{7/10}^1$

$H_{7/12}^1$

$H_{7/12}^2$

$H_{8/7}^1$

$H_{8/8}^1$






$H_{8/8}^2$



$H_{8/10}^1$



$H_{8/10}^2$



$H_{8/10}^3$



$H_{8/10}^4$

 $H_{8/10}^5$ $H_{8/10}^6$ $H_{8/10}^7$ $H_{8/10}^8$ $H_{8/10}^9$  $H_{8/10}^{10}$ $H_{8/10}^{11}$ $H_{8/10}^{12}$ $H_{9/8}^1$ $H_{9/10}^1$  $H_{9/12}^1$ $H_{9/12}^2$ $H_{9/12}^3$ $H_{9/12}^4$ $H_{9/12}^5$









$H_{9/12}^3$	199	214
$H_{9/12}^4$	214	
$H_{9/12}^5$	214	
$H_{9/12}^6$	199	214
$H_{9/12}^7$	199	214
$H_{9/12}^8$	206	214
$H_{9/12}^9$	206	
$H_{9/12}^{10}$	206	
$H_{9/12}^{11}$	220	230
$H_{9/12}^{12}$	220	230
$H_{9/12}^{13}$	220	230
$H_{9/12}^{14}$	220	230
$H_{10/12}^1$	206	
$H_{10/12}^2$	206	
$H_{10/12}^3$	206	
$H_{10/12}^4$	206	
$H_{10/12}^5$	206	
$H_{10/12}^6$	206	
$H_{10/12}^7$	206	
$H_{10/12}^8$	230	
$H_{10/12}^9$	230	
$H_{10/12}^{10}$	230	
$H_{10/12}^{11}$	230	
$H_{10/12}^{12}$	230	
$H_{11/10}^1$	222	
$H_{11/12}^1$	206	
$H_{11/12}^2$	206	
$H_{11/12}^3$	220	230
$H_{11/12}^4$	220	230
$H_{11/14}^1$	220	230
$H_{11/14}^2$	220	230
$H_{11/14}^3$	220	230
$H_{11/14}^4$	220	230
$H_{11/14}^5$	220	230
$H_{11/14}^6$	220	230
$H_{13/16}^1$	220	230
$H_{13/16}^2$	220	230
$H_{13/16}^3$	220	230
$H_{13/16}^4$	220	230
$H_{13/16}^5$	220	230

$H_{3/6}^1$	218																
$H_{3/6}^2$	198	212	213														
$H_{4/3}^1$	198	212	213	220													
$H_{4/3}^2$	152	154	171	172	180	181											
$H_{4/4}^1$	159	190	220	230													
$H_{4/4}^2$	4	17	18	19	20	28	29	30	31	32	33	34	36	40	41	43	45
	52	54	56	57	60	61	62	76	77	78	90	91	92	93	94	95	96
	100	102	104	106	109	110	113	114	117	118	120	122	130	133	138	158	159
	169	170	173	178	179	182	188	190	198	208	210	214	219	220	222	226	228
$H_{4/4}^4$	198	199	205	206	212	213	214	220	230								
$H_{4/6}^1$	199	214	230														
$H_{4/8}^1$	198	212	213														
$H_{5/4}^1$	159	190	199	214	218	220	223	230									
$H_{5/4}^3$	80	92	96	98	109	122	142	159	190	210	222	228					
$H_{5/6}^1$	220																
$H_{5/6}^2$	205																
$H_{5/6}^4$	219																
$H_{6/6}^1$	220																
$H_{6/6}^5$	144	145	151	152	153	154	171	172	180	181							
$H_{6/8}^1$	212	213															
$H_{6/8}^2$	220																
$H_{7/6}^1$	220																
$H_{7/6}^3$	169	170	178	179													
$H_{7/8}^1$	220																
$H_{7/8}^2$	220																
$H_{7/8}^3$	220																
$H_{7/8}^9$	212	213															
$H_{7/8}^{10}$	220																
$H_{7/8}^{11}$	212	213															
$H_{8/8}^1$	76	78	91	92	95	96	110	212	213								
$H_{12/12}^1$	169	170	178	179													

S6. HILBERT BASIS AND BIPARTITE GRAPH IN 230 SPACE GROUPS WITHOUT SOC

SG 1

Space group $P1$, No. 1, without SOC

SG 2

Space group $P\bar{1}$, No. 2, without SOC

SG 3

Space group $P2$, No. 3, without SOC

SG 4

Space group $P2_1$, No. 4, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	Γ_2	$Z_1 Z_2$
dim	1	1	2
Λ_1	1	1	0
Λ_2	1	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_1	Γ_2	$Z_1 Z_2$	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	B_1	B_2	$D_1 D_2$
dim	1	1	2
V_1	1	1	0
V_2	1	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	B_1	B_2	$D_1 D_2$	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	C_1C_2
	dim	1	1	2
W_1	1	1	0	-1
W_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	Y_2	C_1C_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U: $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	E_1E_2
	dim	1	1	2
U_1	1	1	0	-1
U_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_1	A_2	E_1E_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 5

Space group $C2$, No. 5, without SOC

SG 6

Space group Pm , No. 6, without SOC

SG 7

Space group Pc , No. 7, without SOC

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V: $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	B_1B_2	D_1D_2
dim	2	2
V_1V_1	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	B_1B_2	D_1D_2	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U: $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	E_1E_2
dim	2	2
U_1U_1	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_1A_2	E_1E_2	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 8

Space group Cm , No. 8, without SOC

SG 9

Space group Cc , No. 9, without SOC

U

High symmetry line $U(-\frac{1}{2}, -u, u)$, high symmetry points at both ends of U: $A(-\frac{1}{2}, 0, 0), M(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	M_1M_2
dim	2	2
U_1U_1	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$A_1 A_2$	$M_1 M_2$	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 10

Space group $P2/m$, No. 10, without SOC

SG 11

Space group $P2_1/m$, No. 11, without SOC

 Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Z_1
	dim	1	1	1	1	2
Λ_1	1	1	1	0	0	-1
Λ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Z_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

 V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		B_1^+	B_1^-	B_2^+	B_2^-	D_1
	dim	1	1	1	1	2
V_1	1	1	1	0	0	-1
V_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	B_1^+	B_1^-	B_2^+	B_2^-	D_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	C_1
	dim	1	1	1	1	2
W_1	1	1	1	0	0	-1
W_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	C_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U : $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	E_1
	dim	1	1	1	1	2
U_1	1	1	1	0	0	-1
U_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_1^+	A_1^-	A_2^+	A_2^-	E_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

SG 12

Space group $C2/m$, No. 12, without SOC

SG 13

Space group $P2/c$, No. 13, without SOC

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		B_1	D_1
	dim	2	2
V_1V_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	B_1	D_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U : $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	E_1
	dim	2	2
U_1U_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_1	E_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 14

Space group $P2_1/c$, No. 14, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Z_1
	dim	1	1	1	1	2
Λ_1	1	1	1	0	0	-1
Λ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Z_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		B_1	$D_1^+ D_2^+$	$D_1^- D_2^-$
	dim	2	2	2
$V_1 V_2$	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	B_1	$D_1^+ D_2^+$	$D_1^- D_2^-$	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	C_1
	dim	1	1	1	1	2
W_1	1	1	1	0	0	-1
W_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	C_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U: $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	$E_1^+ E_2^+$	$E_1^- E_2^-$
dim	2	2	2
$U_1 U_2$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1	$E_1^+ E_2^+$	$E_1^- E_2^-$	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

SG 15

Space group $C2/c$, No. 15, without SOC

U

High symmetry line $U(-\frac{1}{2}, -u, u)$, high symmetry points at both ends of U: $A(-\frac{1}{2}, 0, 0), M(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	M_1
dim	2	2
$U_1 U_2$	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_1	M_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 16

Space group $P222$, No. 16, without SOC

SG 17

Space group $P222_1$, No. 17, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1
	dim	1	1	1	1	2
Λ_1	1	1	0	1	0	-1
Λ_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H : $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	Y_3	Y_4	T_1
	dim	1	1	1	1	2
H_1	1	1	0	1	0	-1
H_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	Y_3	Y_4	T_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	U_1
	dim	1	1	1	1	2
G_1	1	1	0	1	0	-1
G_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	X_3	X_4	U_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1	T_1
	dim	2	2
$B_1 B_2$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	U_1
	dim	2	2
$A_1 A_2$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	U_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_1	R_1
dim		2	2
P_1P_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_1	R_1
dim		2	2
E_1E_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_1	S_2	S_3	S_4	R_1
dim		1	1	1	1	2
Q_1	1	1	0	1	0	-1
Q_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_1	S_2	S_3	S_4	R_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

SG 18

Space group $P2_12_12$, No. 18, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
	dim	1	1	1	1	2
Δ_1	1	1	1	0	0	-1
Δ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	X_1
	dim	1	1	1	1	2
Σ_1	1	1	0	0	1	-1
Σ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	X_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_1	S_1S_2	S_3S_4
dim	2	2	2
C_1C_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_1	T_1
dim	2	2
H_1H_2	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_1	S_1S_2	S_3S_4
dim	2	2	2
D_1D_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1	U_1
dim	2	2
G_1G_2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	U_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	Z_3	Z_4	T_1
dim	1	1	1	1	2
B_1	1	1	1	0	-1
B_2	1	0	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	Z_3	Z_4	T_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	U_1
	dim	1	1	1	1	2
A_1	1	1	0	0	1	-1
A_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	Z_3	Z_4	U_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_1	R_1R_2	R_3R_4
	dim	2	2	2
P_1P_2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_1	R_1R_2	R_3R_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_1	R_1R_2	R_3R_4
	dim	2	2	2
E_1E_2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_1	R_1R_2	R_3R_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1S_2	S_3S_4	R_1R_2	R_3R_4
	dim	2	2	2	2
Q_1Q_1	2	1	0	-1	0
Q_2Q_2	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1S_2	S_3S_4	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 19

Space group $P2_12_12_1$, No. 19, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
	dim	1	1	1	1	2
Δ_1	1	1	1	0	0	-1
Δ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	X_1
	dim	1	1	1	1	2
Σ_1	1	1	0	0	1	-1
Σ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	X_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

 Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1
	dim	1	1	1	1	2
Λ_1	1	1	0	1	0	-1
Λ_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

 C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_1	S_1S_2	S_3S_4
dim	2	2	2
C_1C_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_1	T_1T_2	T_3T_4
dim	2	2	2
H_1H_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	T_1T_2	T_3T_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	S_1S_2	S_3S_4
dim	2	2	2
D_1D_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1	U_1U_2	U_3U_4
dim	2	2	2
G_1G_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	U_1U_2	U_3U_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_1	T_1T_2	T_3T_4
dim	2	2	2
B_1B_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	T_1T_2	T_3T_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	U_1U_2	U_3U_4
dim	2	2	2
A_1A_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	U_1U_2	U_3U_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_1U_2	U_3U_4	R_1R_1
	dim	2	2	4
P_1P_1	2	1	0	-1
P_2P_2	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_1U_2	U_3U_4	R_1R_1	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_1T_2	T_3T_4	R_1R_1
	dim	2	2	4
E_1E_1	2	1	0	-1
E_2E_2	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1T_2	T_3T_4	R_1R_1	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_1S_2	S_3S_4	R_1R_1
dim	2	2	4
Q_1Q_1	2	0	1
Q_2Q_2	2	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1S_2	S_3S_4	R_1R_1	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 20

Space group $C222_1$, No. 20, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Z_1
dim	1	1	1	1	2
Λ_1	1	1	0	1	0
Λ_2	1	0	1	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Y_1	Y_2	Y_3	Y_4	T_1
dim	1	1	1	1	2
H_1	1	1	0	1	0
H_2	1	0	1	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	Y_3	Y_4	T_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	T_1
dim	2	2
$A_1 A_2$	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_1	S_2	$R_1 R_2$
dim	1	1	2
D_1	1	1	0 -1
D_2	1	0	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	S_2	$R_1 R_2$	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 21

Space group $C222$, No. 21, without SOC

SG 22

Space group $F222$, No. 22, without SOC

SG 23

Space group $I222$, No. 23, without SOC

SG 24

Space group $I2_12_12_1$, No. 24, without SOC

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q : $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep		R_1	R_2	W_1
	dim	1	1	2
Q_1	1	1	0	-1
Q_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	R_2	W_1	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D : $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep		S_1	S_2	W_1
	dim	1	1	2
D_1	1	0	1	-1
D_2	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	S_2	W_1	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of *P*: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	T_1	T_2	W_1
dim	1	1	2
P_1	1	1	0
P_2	1	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	T_2	W_1	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 25

Space group *Pmm2*, No. 25, without SOC

SG 26

Space group *Pmc2₁*, No. 26, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1
Λ_2	1	0	1	0	0	0
Λ_3	1	0	0	1	0	0
Λ_4	1	0	0	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4
dim		1	1	1	1	2	2
H_1	1	1	0	0	0	-1	0
H_2	1	0	1	0	0	0	-1
H_3	1	0	0	1	0	0	-1
H_4	1	0	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	U_1U_2	U_3U_4
dim		1	1	1	1	2	2
G_1	1	1	0	0	0	-1	0
G_2	1	0	1	0	0	0	-1
G_3	1	0	0	1	0	0	-1
G_4	1	0	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	X_3	X_4	U_1U_2	U_3U_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4
dim	2	2	2	2
B_1B_1	2	1	0	-1
B_2B_2	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4
dim	2	2	2	2
A_1A_2	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1U_2	U_3U_4	R_1R_2	R_3R_4
dim	2	2	2	2
P_1P_1	2	1	0	-1
P_2P_2	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_1U_2	U_3U_4	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_1T_2	T_3T_4	R_1R_2	R_3R_4
dim	2	2	2	2
E_1E_2	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1T_2	T_3T_4	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_1	S_2	S_3	S_4	R_1R_2	R_3R_4
dim	1	1	1	1	2	2
Q_1	1	1	0	0	0	-1
Q_2	1	0	1	0	0	0
Q_3	1	0	0	1	0	0
Q_4	1	0	0	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1	S_2	S_3	S_4	R_1R_2	R_3R_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

SG 27

Space group $Pcc2$, No. 27, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	-1
Λ_3	1	0	0	1	0	-1	0
Λ_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H : $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4
	dim	1	1	1	1	2	2
H_1	1	1	0	0	0	-1	0
H_2	1	0	1	0	0	0	-1
H_3	1	0	0	1	0	-1	0
H_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	U_1U_2	U_3U_4
	dim	1	1	1	1	2	2
G_1	1	1	0	0	0	-1	0
G_2	1	0	1	0	0	0	-1
G_3	1	0	0	1	0	-1	0
G_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	X_3	X_4	U_1U_2	U_3U_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4
	dim	2	2	2	2
B_1B_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4
dim		2	2	2	2
A_1A_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2})$, $R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_1U_2	U_3U_4	R_1R_2	R_3R_4
dim		2	2	2	2
P_1P_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1U_2	U_3U_4	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2})$, $R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_1T_2	T_3T_4	R_1R_2	R_3R_4
dim		2	2	2	2
E_1E_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1T_2	T_3T_4	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1	S_2	S_3	S_4	R_1R_2	R_3R_4
	dim	1	1	1	1	2	2
Q_1	1	1	0	0	0	-1	0
Q_2	1	0	1	0	0	0	-1
Q_3	1	0	0	1	0	-1	0
Q_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1	S_2	S_3	S_4	R_1R_2	R_3R_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

SG 28

Space group $Pma2$, No. 28, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
	dim	1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_1	S_1
dim		2	2
C_1C_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	S_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_1	T_1
dim		2	2
H_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1	X_2	X_3	X_4	S_1
	dim	1	1	1	1	2
D_1	1	1	0	0	1	-1
D_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	X_3	X_4	S_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	T_1
	dim	1	1	1	1	2
B_1	1	1	0	0	1	-1
B_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	Z_3	Z_4	T_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_1	U_2	U_3	U_4	R_1
	dim	1	1	1	1	2
P_1	1	1	0	0	1	-1
P_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	U_2	U_3	U_4	R_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_1	R_1
dim	2	2
E_1E_2	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_1	R_1
dim	2	2
Q_1	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 29

Space group $Pca2_1$, No. 29, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
dim		1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	Bipartite graph
	1	1	1	1	2	
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
dim		1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	-1
Λ_3	1	0	0	1	0	0	-1
Λ_4	1	0	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	Bipartite graph
	1	1	1	1	2	2	
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

—————

corep	Y_1	S_1
dim	2	2
$C_1 C_2$	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	S_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_1	$T_1 T_1$
dim	2	4
H_1	2	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	$T_1 T_1$	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	X_3	X_4	S_1
dim	1	1	1	1	2
D_1	1	1	0	0	1 -1
D_2	1	0	1	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	X_3	X_4	S_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	U_1U_2	U_3U_4
	dim	1	1	1	1	2	2
G_1	1	1	0	0	0	-1	0
G_2	1	0	1	0	0	0	-1
G_3	1	0	0	1	0	0	-1
G_4	1	0	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	X_3	X_4	U_1U_2	U_3U_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1Z_2	Z_3Z_4	T_1T_1
	dim	2	2	4
B_1B_1	2	1	0	-1
B_2B_2	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1Z_2	Z_3Z_4	T_1T_1	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4
	dim	2	2	2	2
A_1A_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_1U_2	U_3U_4	R_1R_1
	dim	2	2	4
P_1P_1	2	1	0	-1
P_2P_2	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_1U_2	U_3U_4	R_1R_1	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_1T_1	R_1R_1
	dim	4	4
E_1E_1	2	1	-1
E_2E_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1T_1	R_1R_1	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_1	$R_1 R_1$
dim		2	4
Q_1	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	$R_1 R_1$	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

SG 30

Space group *Pnc2*, No. 30, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
dim		1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	-1
Λ_3	1	0	0	1	0	-1	0
Λ_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	Bipartite graph
	1	1	1	1	2	2	
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_1	S_1
	dim	2	2
C_1C_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	S_1	Bipartite graph
	2	2	
v^1	1	1	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_1	T_1
	dim	2	2
H_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	T_1	Bipartite graph
	2	2	
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_1	X_2	X_3	X_4	S_1
	dim	1	1	1	1	2
D_1	1	1	0	0	1	-1
D_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	X_3	X_4	S_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	U_1U_2	U_3U_4
	dim	1	1	1	1	2	2
G_1	1	1	0	0	0	-1	0
G_2	1	0	1	0	0	0	-1
G_3	1	0	0	1	0	-1	0
G_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	X_3	X_4	U_1U_2	U_3U_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	T_1
dim	2	2	2
B_1B_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	T_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4
dim	2	2	2	2
A_1A_2	2	1	1	-1
			-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1U_2	U_3U_4	R_1
dim	2	2	2
P_1P_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_1U_2	U_3U_4	R_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_1	R_1
	dim	2	2
E_1	1	1	-1
E_2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1	R_1
	dim	2	2
Q_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 31

Space group $Pmn2_1$, No. 31, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
	dim	1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	-1	0
Λ_3	1	0	0	1	0	0	-1
Λ_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	1	1	0	1	$H_{3/2}^1$
v^2	1	1	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_1	S_1
dim	2	2
C_1C_2	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	S_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_1	T_1
dim		2	2
H_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_1	X_2	X_3	X_4	S_1
dim		1	1	1	1	2
D_1	1	1	0	0	1	-1
D_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	X_3	X_4	S_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	U_1U_2	U_3U_4
dim		1	1	1	1	2	2
G_1	1	1	0	0	0	-1	0
G_2	1	0	1	0	0	-1	0
G_3	1	0	0	1	0	0	-1
G_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	X_3	X_4	U_1U_2	U_3U_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	1	1	0	1	$H_{3/2}^1$
v^2	1	1	0	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	T_1
dim	2	2	2
B_1B_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	T_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4
dim	2	2	2	2
A_1A_1	2	1	0	-1
A_2A_2	2	0	1	0
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	U_1U_2	U_3U_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1U_2	U_3U_4	R_1
dim	2	2	2
P_1P_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_1U_2	U_3U_4	R_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_1	R_1
dim	2	2
E_1E_2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_1	R_1
dim	2	2
Q_1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 32

Space group $Pba2$, No. 32, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
	dim	1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	X_1
	dim	1	1	1	1	2
Σ_1	1	1	1	0	0	-1
Σ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	X_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_1	S_1S_2	S_3S_4
dim	2	2	2
C_1C_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_1	T_1
dim	2	2
H_1	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_1	S_1S_2	S_3S_4
dim	2	2	2
D_1D_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	U_1
dim		2	2
G_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	U_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	T_1
dim		1	1	1	1	2
B_1	1	1	0	0	1	-1
B_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	Z_3	Z_4	T_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	U_1
dim		1	1	1	1	2
A_1	1	1	1	0	0	-1
A_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	Z_3	Z_4	U_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1	R_1R_2	R_3R_4
dim	2	2	2
P_1P_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_1	R_1R_2	R_3R_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_1	R_1R_2	R_3R_4
dim	2	2	2
E_1E_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_1	R_1R_2	R_3R_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1S_2	S_3S_4	R_1R_2	R_3R_4
	dim	2	2	2	2
Q_1	1	1	0	-1	0
Q_2	1	0	1	0	-1
Q_3	1	1	0	-1	0
Q_4	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1S_2	S_3S_4	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 33

Space group $Pna2_1$, No. 33, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
	dim	1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	X_1
	dim	1	1	1	1	2
Σ_1	1	1	1	0	0	-1
Σ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	X_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

 Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	-1
Λ_3	1	0	0	1	0	0	-1
Λ_4	1	0	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

 C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_1	S_1S_2	S_3S_4
dim	2	2	2
C_1C_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_1	T_1T_1
dim	2	4
H_1	2	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	T_1T_1	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	S_1S_2	S_3S_4
dim	2	2	2
D_1D_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1	U_1
dim	2	2
G_1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	U_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	T_1T_1
dim	2	2	4
B_1B_1	2	1	-1
B_2B_2	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1Z_2	Z_3Z_4	T_1T_1	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	U_1
dim	2	2	2
A_1A_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	U_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_1	R_1R_2	R_3R_4
	dim	2	2	2
P_1P_2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

		U_1	R_1R_2	R_3R_4	
		2	2	2	Bipartite graph
v^1	1	0	1		$H_{2/1}^1$
v^2	1	1	0		$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_1T_1	R_1R_2	R_3R_4
	dim	4	2	2
E_1E_1	2	1	0	-1
E_2E_2	2	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		T_1T_1	R_1R_2	R_3R_4	
		4	2	2	Bipartite graph
v^1	1	1	1		$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1S_2	S_3S_4	R_1R_2	R_3R_4
	dim	2	2	2	2
Q_1	1	1	0	0	-1
Q_2	1	0	1	0	-1
Q_3	1	1	0	-1	0
Q_4	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1S_2	S_3S_4	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 34

Space group $Pnn2$, No. 34, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
dim		1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	X_1
dim		1	1	1	1	2
Σ_1	1	1	1	0	0	-1
Σ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	X_1	
	1	1	1	1	2	Bipartite graph

v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	-1
Λ_3	1	0	0	1	0	-1	0
Λ_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_1	S_1S_2	S_3S_4
	dim	2	2	2
C_1C_2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_1	T_1
dim	2	2
H_1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	S_1S_2	S_3S_4
dim	2	2	2
D_1D_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	S_1S_2	S_3S_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	U_1
dim	2	2
G_1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	U_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	T_1
dim	2	2	2
B_1B_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	T_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	U_1
dim	2	2	2
A_1A_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	U_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_1	R_1	R_2	R_3	R_4
dim	2	1	1	1	1
P_1	1	1	-1	0	0
P_2	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	R_1	R_2	R_3	R_4	
	2	1	1	1	1	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E : $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_1	R_1	R_2	R_3	R_4
dim		2	1	1	1	1
E_1	1	1	0	0	-1	-1
E_2	1	1	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1	R_1	R_2	R_3	R_4	
	2	1	1	1	1	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1S_2	S_3S_4	R_1	R_2	R_3	R_4
dim		2	2	1	1	1	1
Q_1	1	1	0	0	0	-1	0
Q_2	1	0	1	0	0	0	-1
Q_3	1	1	0	-1	0	0	0
Q_4	1	0	1	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1S_2	S_3S_4	R_1	R_2	R_3	R_4	
	2	2	1	1	1	1	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

SG 35

Space group $Cmm2$, No. 35, without SOC

SG 36

Space group $Cmc2_1$, No. 36, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	-1	0
Λ_3	1	0	0	1	0	0	-1
Λ_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	1	1	0	1	$H_{3/2}^1$
v^2	1	1	0	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4
	dim	1	1	1	1	2	2
H_1	1	1	0	0	0	-1	0
H_2	1	0	1	0	0	-1	0
H_3	1	0	0	1	0	0	-1
H_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	1	1	0	1	$H_{3/2}^1$
v^2	1	1	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4	
dim	2	2	2	2	
A_1A_1	2	1	0	-1	0
A_2A_2	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_1	S_2	R_1R_2	
dim	1	1	2	
D_1	1	1	0	-1
D_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	S_2	R_1R_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 37

Space group $Ccc2$, No. 37, without SOC

 Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	-1
Λ_3	1	0	0	1	0	-1	0
Λ_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	Bipartite graph
	1	1	1	1	2	2	
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4
	dim	1	1	1	1	2	2
H_1	1	1	0	0	0	-1	0
H_2	1	0	1	0	0	0	-1
H_3	1	0	0	1	0	-1	0
H_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	Y_2	Y_3	Y_4	T_1T_2	T_3T_4	Bipartite graph
	1	1	1	1	2	2	
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4
	dim	2	2	2	2
A_1A_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_2	Z_3Z_4	T_1T_2	T_3T_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 38

Space group $Amm2$, No. 38, without SOC

SG 39

Space group $Abm2$, No. 39, without SOC

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D : $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_1S_2	R_1R_2
dim	2	2
D_1D_1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1S_2	R_1R_2	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 40

Space group $Ama2$, No. 40, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Z_1
dim	1	1	1	1	2
Λ_1	1	1	1	0	-1
Λ_2	1	0	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	Y_3	Y_4	T_1
	dim	1	1	1	1	2
H_1	1	1	1	0	0	-1
H_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	Y_3	Y_4	T_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	T_1
	dim	2	2
$A_1 A_2$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 41

Space group *Aba2*, No. 41, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1
	dim	1	1	1	1	2
Λ_1	1	1	1	0	0	-1
Λ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	Y_3	Y_4	T_1
	dim	1	1	1	1	2
H_1	1	0	0	1	1	-1
H_2	1	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	Y_3	Y_4	T_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	T_1
dim	2	2
$A_1 A_2$	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	T_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$S_1 S_2$	$R_1 R_2$
dim	2	2
$D_1 D_1$	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$S_1 S_2$	$R_1 R_2$	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 42

Space group $Fmm2$, No. 42, without SOC

SG 43

Space group $Fdd2$, No. 43, without SOC

Λ

High symmetry line $\Lambda(u, u, 0)$, high symmetry points at both ends of Λ: $\Gamma(0, 0, 0), Z(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4
	dim	1	1	1	1	2	2
Λ_1	1	1	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	-1
Λ_3	1	0	0	1	0	-1	0
Λ_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	Γ_4	Z_1Z_2	Z_3Z_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	X_1
	dim	1	1	1	1	2
Σ_1	1	1	1	0	0	-1
Σ_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	X_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, -u, -u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(0, -\frac{1}{2}, -\frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Y_1
	dim	1	1	1	1	2
Δ_1	1	1	0	0	1	-1
Δ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Y_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

SG 44

Space group *Imm2*, No. 44, without SOC

SG 45

Space group *Iba2*, No. 45, without SOC

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	R_1R_2	W_1W_1	W_2W_2
dim	2	2	2
Q_1Q_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1R_2	W_1W_1	W_2W_2	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	S_1S_2	W_1W_1	W_2W_2
dim	2	2	2
D_1D_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$S_1 S_2$	$W_1 W_1$	$W_2 W_2$	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		T_1	T_2	$W_1 W_1$	$W_2 W_2$
	dim	1	1	2	2
P_1	1	1	0	-2	0
P_2	1	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_1	T_2	$W_1 W_1$	$W_2 W_2$	
	1	1	2	2	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	2	0	1	0	$H_{3/2}^1$

SG 46

Space group *Ima2*, No. 46, without SOC

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		R_1	R_2	$W_1 W_2$
	dim	1	1	2
Q_1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	R_1	R_2	$W_1 W_2$	
	1	1	2	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		S_1S_2	W_1W_2
dim		2	2
D_1D_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1S_2	W_1W_2	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		T_1	T_2	W_1W_2
dim		1	1	2
P_1	1	1	0	-1
P_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	T_2	W_1W_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 47

Space group $Pmmm$, No. 47, without SOC

SG 48

Space group $Pn\bar{3}n$, No. 48, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	-1	0
Δ_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	1	0	0	1	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	-1	0
Σ_4	1	0	0	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
dim		1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_1	Y_2	S_1	S_2
dim		2	2	2	2
C_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	S_1	S_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	T_1	T_2
dim		2	2	2	2
H_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	T_1	T_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1	X_2	S_1	S_2
dim		2	2	2	2
D_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	S_1	S_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	U_1	U_2
dim		2	2	2	2
G_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	T_1	T_2
	dim	2	2	2	2
B_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	U_1	U_2
	dim	2	2	2	2
A_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_1	U_2	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-
dim	2	2	1	1	1	1	1	1	1	1
P_1	1	1	0	-1	0	0	-1	0	0	0
P_2	1	1	0	0	-1	-1	0	0	0	0
P_3	1	0	1	0	0	0	0	-1	0	-1
P_4	1	0	1	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	U_1	U_2	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	Bipartite graph
	2	2	1	1	1	1	1	1	1	1	
v^1	0	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^3	0	1	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^6	1	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^7	1	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^8	1	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_1	T_2	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-
dim	2	2	1	1	1	1	1	1	1	1
E_1	1	1	0	-1	0	0	0	0	0	-1
E_2	1	1	0	0	-1	0	0	0	0	-1
E_3	1	0	1	0	0	0	-1	-1	0	0
E_4	1	0	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	T_1	T_2	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	Bipartite graph
	2	2	1	1	1	1	1	1	1	1	
v^1	0	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^2	0	1	0	0	0	1	0	1	0	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	1	0	0	0	$H_{3/2}^1$

Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
H_1	1	1	0	0	0	0	1	0	0	-1	0
H_2	1	0	0	1	0	0	0	0	1	0	-1
H_3	1	0	1	0	0	1	0	0	0	-1	0
H_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	U_1	U_2
	dim	1	1	1	1	1	1	1	1	2	2
G_1	1	1	0	0	0	0	1	0	0	-1	0
G_2	1	0	0	1	0	0	0	0	1	0	-1
G_3	1	0	1	0	0	1	0	0	0	-1	0
G_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	U_1	U_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	T_1	T_2
	dim	2	2	2	2
B_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	U_1	U_2
	dim	2	2	2	2
A_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_1	U_2	R_1	R_2
	dim	2	2	2	2
P_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	U_2	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_1	T_2	R_1	R_2
	dim	2	2	2	2
E_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1	T_2	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1^+	S_1^-	S_2^+	S_2^-	S_3^+	S_3^-	S_4^+	S_4^-	R_1	R_2
	dim	1	1	1	1	1	1	1	1	2	2
Q_1	1	1	0	0	0	0	1	0	0	-1	0
Q_2	1	0	0	1	0	0	0	0	1	0	-1
Q_3	1	0	1	0	0	1	0	0	0	-1	0
Q_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	S_1^+	S_1^-	S_2^+	S_2^-	S_3^+	S_3^-	S_4^+	S_4^-	R_1	R_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

SG 50

Space group $Pban$, No. 50, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	-1	0
Δ_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	-1	0
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y ₁	Y ₂	S ₁	S ₂
dim	2	2	2	2
C_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y ₁	Y ₂	S ₁	S ₂	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y ₁	Y ₂	T ₁	T ₂
dim	2	2	2	2
H_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y ₁	Y ₂	T ₁	T ₂	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1	X_2	S_1	S_2
	dim	2	2	2	2
D_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	S_1	S_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	U_1	U_2
	dim	2	2	2	2
G_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
B_1	1	1	0	0	1	0	0	0	0	-1	0
B_2	1	0	0	0	0	0	1	1	0	0	-1
B_3	1	0	1	1	0	0	0	0	0	-1	0
B_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2
	dim	1	1	1	1	1	1	1	1	2	2
A_1	1	1	0	0	0	0	0	0	1	-1	0
A_2	1	0	0	0	1	1	0	0	0	0	-1
A_3	1	0	1	0	0	0	0	1	0	-1	0
A_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_1	U_2	R_1	R_2
	dim	2	2	2	2
P_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	U_2	R_1	R_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_1	T_2	R_1	R_2
	dim	2	2	2	2
E_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1	T_2	R_1	R_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1	S_2	R_1	R_2
	dim	2	2	2	2
Q_1	1	1	0	-1	0
Q_2	1	1	0	-1	0
Q_3	1	0	1	0	-1
Q_4	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

H_1	1	1	0	0	0	0	1	0	0	-1	0
H_2	1	0	0	1	0	0	0	0	1	-1	0
H_3	1	0	1	0	0	1	0	0	0	0	-1
H_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G : $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	U_1	U_2
	dim	1	1	1	1	1	1	1	1	2	2
G_1	1	1	0	0	0	0	1	0	0	-1	0
G_2	1	0	0	1	0	0	0	0	1	-1	0
G_3	1	0	1	0	0	1	0	0	0	0	-1
G_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	U_1	U_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	T_1	T_2
	dim	2	2	2	2
B_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	U_1	U_2
	dim	2	2	2	2
A_1A_2	2	1	0	-1	0
A_3A_4	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	Z_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_1	U_2	R_1	R_2
	dim	2	2	2	2
P_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	U_2	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_1	T_2	R_1	R_2
dim	2	2	2	2
E_1E_2	2	1	0	-1
E_3E_4	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_1	T_2	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_1^+	S_1^-	S_2^+	S_2^-	S_3^+	S_3^-	S_4^+	S_4^-	R_1	R_2
dim	1	1	1	1	1	1	1	1	2	2
Q_1	1	1	0	0	0	0	1	0	-1	0
Q_2	1	0	0	1	0	0	0	0	-1	0
Q_3	1	0	1	0	0	1	0	0	0	-1
Q_4	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

S_1^+	S_1^-	S_2^+	S_2^-	S_3^+	S_3^-	S_4^+	S_4^-	R_1	R_2	
1	1	1	1	1	1	1	1	2	2	Bipartite graph

v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

SG 52

Space group $Pnna$, No. 52, without SOC Δ High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$ **A** matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	-1	0
Δ_4	1	0	0	0	0	1	0	0	1	0	-1

 $\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

 Σ High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$ **A** matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	-1	0
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	-1	0
Λ_3	1	0	1	0	0	1	0	0	0	0	-1
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y ₁	Y ₂	S ₁	S ₂
dim	2	2	2	2
C_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y ₁	Y ₂	S ₁	S ₂	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y ₁	Y ₂	T ₁ ⁺	T ₁ ⁻
dim	2	2	2	2
H_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y ₁	Y ₂	T ₁ ⁺	T ₁ ⁻	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1	X_2	S_1	S_2
	dim	2	2	2	2
D_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	S_1	S_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	U_1U_2
	dim	2	2	4
G_1	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	U_1U_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	T_1^+	T_1^-
	dim	2	2	2	2
B_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1^+	T_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	U_1U_2
dim	2	2	4
A_1A_2	2	1	0
A_3A_4	2	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	Z_2	U_1U_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1U_2	R_1	R_2
dim	4	2	2
P_1P_4	2	1	-1
P_2P_3	2	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_1U_2	R_1	R_2	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_1^+	T_1^-	R_1	R_2
dim		2	2	2	2
E_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1^+	T_1^-	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1	S_2	R_1	R_2
dim		2	2	2	2
Q_1	1	1	0	-1	0
Q_2	1	1	0	0	-1
Q_3	1	0	1	0	-1
Q_4	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	S_2	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 53

Space group $Pmna$, No. 53, without SOC

 Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	-1	0
Δ_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	-1	0
Λ_3	1	0	1	0	0	1	0	0	0	0	-1
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y ₁	Y ₂	S ₁	S ₂
dim	2	2	2	2
C_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y ₁	Y ₂	S ₁	S ₂	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y ₁	Y ₂	T ₁ ⁺	T ₁ ⁻
dim	2	2	2	2
H_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y ₁	Y ₂	T ₁ ⁺	T ₁ ⁻	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	S_1	S_2
	dim	1	1	1	1	1	1	1	1	2	2
D_1	1	1	0	0	1	0	0	0	0	-1	0
D_2	1	0	0	0	0	0	1	1	0	0	-1
D_3	1	0	1	1	0	0	0	0	0	-1	0
D_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	S_1	S_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G : $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	U_1	U_2
	dim	1	1	1	1	1	1	1	1	2	2
G_1	1	1	0	0	0	0	1	0	0	-1	0
G_2	1	0	0	1	0	0	0	0	1	-1	0
G_3	1	0	1	0	0	1	0	0	0	0	-1
G_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	U_1	U_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	T_1^+	T_1^-
dim	2	2	2	2
B_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1^+	T_1^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	U_1	U_2
dim	2	2	2	2
A_1A_2	2	1	0	-1 0
A_3A_4	2	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	Z_2	U_1	U_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1	U_2	R_1^+	R_1^-
dim	2	2	2	2
P_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	U_2	R_1^+	R_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_1^+	T_1^-	R_1^+	R_1^-
dim	2	2	2	2
E_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1^+	T_1^-	R_1^+	R_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_1	S_2	R_1^+	R_1^-
dim	2	2	2	2
Q_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_1	S_2	R_1^+	R_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 54

Space group $Pcca$, No. 54, without SOC

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	-1	0
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	-1	0
Λ_3	1	0	1	0	0	1	0	0	0	0	-1
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0)$, $S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	S_1	S_2
	dim	1	1	1	1	1	1	1	1	2	2
C_1	1	1	0	0	0	0	0	0	1	-1	0
C_2	1	0	0	0	1	1	0	0	0	0	-1
C_3	1	0	1	0	0	0	0	1	0	-1	0
C_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	S_1	S_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
H_1	1	1	0	0	0	0	1	0	0	-1	0
H_2	1	0	0	1	0	0	0	0	1	-1	0
H_3	1	0	1	0	0	1	0	0	0	0	-1
H_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_1	X_2	S_1	S_2
	dim	2	2	2	2
D_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	S_1	S_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	U_1U_2
	dim	2	2	4
G_1	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	U_1U_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	T_1	T_2
	dim	2	2	2	2
B_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	U_1U_2
	dim	2	2	4
A_1A_2	2	1	0	-1
A_3A_4	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	Z_2	U_1U_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_1U_2	R_1R_2
	dim	4	4
P_1P_3	2	1	-1
P_2P_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_1U_2	R_1R_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_1	T_2	R_1R_2
	dim	2	2	4
E_1E_2	2	1	0	-1
E_3E_4	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	T_2	R_1R_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_1	S_2	$R_1 R_2$
	dim	2	2	4
Q_1	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	S_1	S_2	$R_1 R_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 55

Space group *Pbam*, No. 55, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	0	-1
Δ_4	1	0	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	-1	0
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_1	Y_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$
dim	2	2	2	2	2	2
$C_1 C_2$	2	1	0	-1	0	0
$C_3 C_4$	2	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_1	Y_2	T_1	T_2
dim		2	2	2	2
H_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1	X_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$
dim		2	2	2	2	2	2
$D_1 D_2$	2	1	0	-1	0	0	-1
$D_3 D_4$	2	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	U_1	U_2
	dim	2	2	2	2
G_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
B_1	1	1	0	0	1	0	0	0	0	-1	0
B_2	1	0	0	0	0	0	1	1	0	0	-1
B_3	1	0	1	1	0	0	0	0	0	0	-1
B_4	1	0	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2
	dim	1	1	1	1	1	1	1	1	2	2
A_1	1	1	0	0	0	0	0	0	1	-1	0
A_2	1	0	0	0	1	1	0	0	0	-1	0
A_3	1	0	1	0	0	0	0	1	0	0	-1
A_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_1	U_2	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$
	dim	2	2	2	2	2	2
$P_1 P_2$	2	1	0	-1	0	0	-1
$P_3 P_4$	2	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	U_2	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_1	T_2	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	
dim	2	2	2	2	2	2	
$E_1 E_2$	2	1	0	-1	0	0	-1
$E_3 E_4$	2	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1	T_2	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	
dim	2	2	2	2	2	2	2	2	
$Q_1 Q_2$	2	1	1	0	0	-1	-1	0	0
$Q_3 Q_4$	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 56

Space group $Pccn$, No. 56, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	-1	0
Δ_3	1	0	1	1	0	0	0	0	0	0	-1
Δ_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$

 Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$

v^4	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_1	Y_2	S_1	S_2
	dim	2	2	2	2
C_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	S_1	S_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_1	Y_2	T_1	T_2
dim	2	2	4	
H_1H_4	2	1	0	-1
H_2H_3	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	Y_2	T_1	T_2	
	2	2	4		Bipartite graph
v^1	1	1	1		$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	S_1	S_2
dim	2	2	2	2
D_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	S_1	S_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1	X_2	U_1U_2
dim	2	2	4
G_1G_2	2	1	0
G_3G_4	2	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	X_2	U_1U_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	T_1T_2
dim	2	2	4
B_1	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	T_1T_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	U_1U_2
dim	2	2	4
A_1	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	U_1U_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1U_2	$R_1^+R_2^+$	$R_1^-R_2^-$	$R_3^+R_4^+$	$R_3^-R_4^-$
dim	4	2	2	2	2
P_1P_4	2	1	0	-1	-1
P_2P_3	2	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1U_2	$R_1^+R_2^+$	$R_1^-R_2^-$	$R_3^+R_4^+$	$R_3^-R_4^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_1T_2	$R_1^+R_2^+$	$R_1^-R_2^-$	$R_3^+R_4^+$	$R_3^-R_4^-$
dim	4	2	2	2	2
E_1E_4	2	1	0	-1	-1
E_2E_3	2	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1T_2	$R_1^+R_2^+$	$R_1^-R_2^-$	$R_3^+R_4^+$	$R_3^-R_4^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_1	S_2	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$
dim	2	2	2	2	2	2
$Q_1 Q_3$	2	1 0	-1	-1	0	0
$Q_2 Q_4$	2	0 1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_1	S_2	$R_1^+ R_2^+$	$R_1^- R_2^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

SG 57

Space group *Pbcm*, No. 57, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	-1
Δ_4	1	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$

v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_1	Y_2	S_1S_2
	dim	2	2	4
C_1C_2	2	1	0	-1
C_3C_4	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	Y_2	S_1S_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_1	Y_2	T_1	T_2
	dim	2	2	2	2
H_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	T_1	T_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_1	X_2	S_1	S_2
	dim	2	2	4	
D_1	2	1	1	-2	

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	S_1	S_2	Bipartite graph
	2	2	4		
v^1	0	2	1		$H_{3/2}^1$
v^2	1	1	1		$H_{3/2}^1$
v^3	2	0	1		$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	U_1	U_2
	dim	2	2	2	2
G_1G_2	2	1	0	-1	0
G_3G_4	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
B_1	1	1	0	0	1	0	0	0	0	-1	0
B_2	1	0	0	0	0	0	1	1	0	0	-1
B_3	1	0	1	1	0	0	0	0	0	0	-1
B_4	1	0	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2
	dim	1	1	1	1	1	1	1	1	2	2
A_1	1	1	0	0	0	0	0	0	1	-1	0
A_2	1	0	0	0	1	1	0	0	0	0	-1
A_3	1	0	1	0	0	0	0	1	0	0	-1
A_4	1	0	0	1	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_1	U_2	R_1R_2
dim	2	2	4
P_1	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_1	U_2	R_1R_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_1	T_2	R_1R_2
dim	2	2	4
E_1E_2	2	1	0
E_3E_4	2	0	1
			-1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	T_2	R_1R_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1S_2	R_1R_2
dim	4	4	4
Q_1Q_1	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		S_1S_2	R_1R_2	Bipartite graph
	4	4	4	
v^1	1	1	1	$H_{2/1}^1$

SG 58

Space group Pnm , No. 58, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
dim	1	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	0	-1
Δ_4	1	0	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	Bipartite graph
	1	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
dim		1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	-1	0
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

 Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
dim		1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$

v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_1	Y_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$
dim	2	2	2	2	2	2
$C_1 C_2$	2	1	0	-1	0	0
$C_3 C_4$	2	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H : $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_1	Y_2	T_1^+	T_1^-
dim	2	2	2	2
H_1	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	T_1^+	T_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$
dim	2	2	2	2	2	2
$D_1 D_2$	2	1 0	-1	0	0	-1
$D_3 D_4$	2	0 1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	U_1^+	U_1^-
dim	2	2	2	2
G_1	2	1 1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	U_1^+	U_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	T_1^+	T_1^-
	dim	2	2	2	2
B_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1^+	T_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	U_1^+	U_1^-
	dim	2	2	2	2
A_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	U_1^+	U_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_1^+	U_1^-	R_1	R_2
	dim	2	2	2	2
P_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1^+	U_1^-	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_1^+	T_1^-	R_1	R_2
dim	2	2	2	2
E_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1^+	T_1^-	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	R_1	R_2
dim	2	2	2	2	2	2
$Q_1 Q_2$	2	1	1	0	0	-1 0
$Q_3 Q_4$	2	0	0	1	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$S_1^+ S_2^+$	$S_1^- S_2^-$	$S_3^+ S_4^+$	$S_3^- S_4^-$	R_1	R_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$

SG 59

Space group $Pm\bar{m}n$, No. 59, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	-1	0
Δ_3	1	0	1	1	0	0	0	0	0	0	-1
Δ_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_1	Y_2	S_1	S_2
dim	2	2	2	2
C_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	S_1	S_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_1	Y_2	T_1	T_2
dim	2	2	2	2
H_1H_4	2	1	0	-1 0
H_2H_3	2	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_1	Y_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_1	X_2	S_1	S_2
dim	2	2	2	2
D_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	S_1	S_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1	X_2	U_1	U_2
dim	2	2	2	2
G_1G_2	2	1	0	-1 0
G_3G_4	2	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	U_1	U_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
B_1	1	1	0	0	1	0	0	0	0	-1	0
B_2	1	0	0	0	0	0	1	1	0	-1	0
B_3	1	0	1	1	0	0	0	0	0	0	-1
B_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	T_1	T_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2
	dim	1	1	1	1	1	1	1	1	2	2
A_1	1	1	0	0	0	0	0	0	1	-1	0
A_2	1	0	0	0	1	1	0	0	0	0	-1
A_3	1	0	1	0	0	0	0	1	0	0	-1
A_4	1	0	0	1	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	U_1	U_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$

v^4	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_1	U_2	R_1	R_2
dim	2	2	2	2
P_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_1	U_2	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_1	T_2	R_1	R_2
dim	2	2	2	2
E_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_1	T_2	R_1	R_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_1	S_2	R_1	R_2
dim	2	2	2	2
Q_1Q_3	2	1	0	-1
Q_2Q_4	2	0	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1	S_2	R_1	R_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 60

Space group $Pbcn$, No. 60, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	-1	0
Δ_3	1	0	1	1	0	0	0	0	0	-1
Δ_4	1	0	0	0	0	1	0	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
dim		1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	-1	0
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

 Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
dim		1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$

v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_1	Y_2	S_1S_2
dim	2	2	4
C_1	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_1	Y_2	S_1S_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_1	Y_2	T_1T_2
dim	2	2	4
H_1H_4	2	1	0
			-1
H_2H_3	2	0	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	Y_2	T_1T_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	$S_1 S_2$
dim	2	2	4
$D_1 D_2$	2	1	0
$D_3 D_4$	2	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	X_2	$S_1 S_2$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	U_1^+	U_1^-
dim	2	2	2	2
G_1	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	U_1^+	U_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	$T_1 T_2$
dim	2	2	4
B_1	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	$T_1 T_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	U_1^+	U_1^-
dim	2	2	2	2
A_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	U_1^+	U_1^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_1^+	U_1^-	$R_1 R_2$
dim	2	2	4
P_1	2	1	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_1^+	U_1^-	$R_1 R_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of *E*: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$T_1 T_2$	$R_1 R_2$
dim	4	4
$E_1 E_4$	2	1
$E_2 E_3$	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$T_1 T_2$	$R_1 R_2$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of *Q*: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$S_1 S_2$	$R_1 R_2$
dim	4	4
$Q_1 Q_1$	4	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$S_1 S_2$	$R_1 R_2$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 61

Space group *Pbca*, No. 61, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	1	1	0	-1	0
Δ_3	1	0	1	1	0	0	0	0	0	-1
Δ_4	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	-1	0
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	-1	0
Λ_3	1	0	1	0	0	1	0	0	0	0	-1
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_1	Y_2	$S_1 S_2$
	dim	2	2	4
C_1	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_1	Y_2	$S_1 S_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix



corep	Y_1	Y_2	T_1T_2
dim	2	2	4
$H_1^+H_2^+$	2	1	0
$H_1^-H_2^-$	2	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	Y_2	T_1T_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	S_1S_2
dim	2	2	4
D_1D_2	2	1	0
D_3D_4	2	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	X_2	S_1S_2	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	U_1U_2
dim	2	2	4
G_1	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	U_1U_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	$T_1 T_2$
dim	2	2	4
B_1	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	$T_1 T_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	$U_1 U_2$
dim	2	2	4
$A_1 A_2$	2	1	0
$A_3 A_4$	2	0	1
			-1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_1	Z_2	$U_1 U_2$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$U_1 U_2$	$R_1^+ R_1^+$	$R_1^- R_1^-$
dim	4	4	4
$P_1 P_1$	4	1	-1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_1U_2	$R_1^+ R_1^+$	$R_1^- R_1^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_1T_2	$R_1^+ R_1^+$	$R_1^- R_1^-$
dim	4	4	4
E_1E_1	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_1T_2	$R_1^+ R_1^+$	$R_1^- R_1^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_1S_2	$R_1^+ R_1^+$	$R_1^- R_1^-$
dim	4	4	4
Q_1Q_1	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1S_2	$R_1^+ R_1^+$	$R_1^- R_1^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

SG 62

Space group $Pnma$, No. 62, without SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
dim		1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	0	-1
Δ_4	1	0	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
dim		1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	0	-1
Σ_4	1	0	0	1	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$

v^4	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	-1	0
Λ_3	1	0	1	0	0	1	0	0	0	0	-1
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_1	Y_2	$S_1 S_2$
	dim	2	2	4
$C_1 C_2$	2	1	0	-1
$C_3 C_4$	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_1	Y_2	$S_1 S_2$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_1	Y_2	T_1	T_2
dim	2	2	2	2
H_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_1	Y_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_1	X_2	$S_1 S_2$
dim	2	2	4
D_1	2	1	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	$S_1 S_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	$U_1^+U_2^+$	$U_1^-U_2^-$	$U_3^+U_4^+$	$U_3^-U_4^-$
dim	2	2	2	2	2	2
G_1G_2	2	1	0	-1	0	-1
G_3G_4	2	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	$U_1^+U_2^+$	$U_1^-U_2^-$	$U_3^+U_4^+$	$U_3^-U_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	T_1	T_2
dim	2	2	2	2
B_1	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	$U_1^+U_2^+$	$U_1^-U_2^-$	$U_3^+U_4^+$	$U_3^-U_4^-$
dim	2	2	2	2	2	2
A_1A_2	2	1	0	-1	0	-1
A_3A_4	2	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	$U_1^+U_2^+$	$U_1^-U_2^-$	$U_3^+U_4^+$	$U_3^-U_4^-$	Bipartite graph
v^1	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	0	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$U_1^+U_2^+$	$U_1^-U_2^-$	$U_3^+U_4^+$	$U_3^-U_4^-$	R_1R_2	
dim	2	2	2	2	4	
P_1P_3	2	1	1	0	0	-1
P_2P_4	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$U_1^+U_2^+$	$U_1^-U_2^-$	$U_3^+U_4^+$	$U_3^-U_4^-$	R_1R_2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_1	T_2	R_1R_2	
dim	2	2	4	
E_1E_3	2	1	0	-1
E_2E_4	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	T_2	$R_1 R_2$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$S_1 S_2$	$R_1 R_2$
dim	4	4
$Q_1 Q_1$	4	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$S_1 S_2$	$R_1 R_2$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 63

Space group $Cmcm$, No. 63, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1
Λ_4	1	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	
1	1	1	1	1	1	1	1	2	2	Bipartite graph

v^1	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
H_1	1	1	0	0	0	0	1	0	0	-1	0
H_2	1	0	0	1	0	0	0	0	1	0	-1
H_3	1	0	1	0	0	1	0	0	0	0	-1
H_4	1	0	0	0	1	0	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	T_1	T_2
	dim	2	2	2	2
A_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1^+	S_1^-	S_2^+	S_2^-	R_1
	dim	1	1	1	1	2
D_1	1	1	1	0	0	-1
D_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_1^+	S_1^-	S_2^+	S_2^-	R_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

SG 64

Space group *Cmca*, No. 64, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	0	-1
Λ_4	1	0	0	0	1	0	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
H_1	1	0	1	0	0	1	0	0	0	0	-1
H_2	1	0	0	0	1	0	0	1	0	-1	0
H_3	1	1	0	0	0	0	1	0	0	-1	0
H_4	1	0	0	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	T_1	T_2
	dim	2	2	2	2
A_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_1	$R_1^+ R_2^+$	$R_1^- R_2^-$
	dim	2	2	2
$D_1 D_2$	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1	$R_1^+ R_2^+$	$R_1^- R_2^-$	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

SG 65

Space group $Cmmm$, No. 65, without SOC

SG 66

Space group $Cccm$, No. 66, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
H_1	1	1	0	0	0	0	1	0	0	-1	0
H_2	1	0	0	1	0	0	0	0	1	0	-1
H_3	1	0	1	0	0	1	0	0	0	-1	0
H_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z ₁	Z ₂	T ₁	T ₂
dim		2	2	2	2
A ₁	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z ₁	Z ₂	T ₁	T ₂	
	2	2	2	2	Bipartite graph
v ¹	0	1	0	1	$H_{2/1}^1$
v ²	0	1	1	0	$H_{2/1}^1$
v ³	1	0	0	1	$H_{2/1}^1$
v ⁴	1	0	1	0	$H_{2/1}^1$

SG 67

Space group *Cmma*, No. 67, without SOC

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S ₁	R ₁
dim		2	2
D ₁ D ₂	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S ₁	R ₁	
	2	2	Bipartite graph
v ¹	1	1	$H_{2/1}^1$

SG 68

Space group *Ccca*, No. 68, without SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2
	dim	1	1	1	1	1	1	1	1	2	2
H_1	1	0	1	0	0	1	0	0	0	-1	0
H_2	1	0	0	0	1	0	0	1	0	0	-1
H_3	1	1	0	0	0	0	1	0	0	-1	0
H_4	1	0	0	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Y_1^+	Y_1^-	Y_2^+	Y_2^-	Y_3^+	Y_3^-	Y_4^+	Y_4^-	T_1	T_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$

v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	T_1	T_2
dim		2	2	2	2
A_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	T_1	T_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_1	R_1
dim		2	2
$D_1 D_2$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	R_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 69

Space group $Fmmm$, No. 69, without SOC

SG 70

Space group $Fddd$, No. 70, without SOC

Λ

High symmetry line $\Lambda(u, u, 0)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2
	dim	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	0	0	1	0	0	-1	0
Λ_2	1	0	0	1	0	0	0	0	1	0	-1
Λ_3	1	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Z_1	Z_2	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	0	1	-1	0
Σ_2	1	0	0	0	1	1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	0	-1	0
Σ_4	1	0	0	1	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, -u, -u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(0, -\frac{1}{2}, -\frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2
	dim	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	1	0	0	0	0	0	-1	0
Δ_4	1	0	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Y_1	Y_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

SG 71

Space group *Immm*, No. 71, without SOC

SG 72

Space group *Ibam*, No. 72, without SOC

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	R_1	W_1W_3	W_2W_4
dim	2	2	2
Q_1Q_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1	W_1W_3	W_2W_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	S_1	W_1W_3	W_2W_4
dim	2	2	2
D_1D_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_1	W_1W_3	W_2W_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		T_1^+	T_1^-	T_2^+	T_2^-	W_1W_3	W_2W_4
dim		1	1	1	1	2	2
P_1	1	1	1	0	0	-2	0
P_2	1	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	T_1^+	T_1^-	T_2^+	T_2^-	W_1W_3	W_2W_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

SG 73

Space group *Ibca*, No. 73, without SOC

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of *Q*: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	R_1	W_1W_1
dim	2	4
Q_1Q_2	2	1
		-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	W_1W_1	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of *D*: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	S_1	W_1W_1
dim	2	4
D_1D_2	2	1
		-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_1	W_1W_1	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	T_1	W_1W_1
dim	2	4
P_1P_2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

T_1	W_1W_1	Bipartite graph
2	4	
v^1	2	$H_{3/2}^1$

SG 74

Space group *Imma*, No. 74, without SOC

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	R_1^+	R_1^-	R_2^+	R_2^-	W_1
dim	1	1	1	1	2
Q_1	1	1	0	0	-1
Q_2	1	0	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

R_1^+	R_1^-	R_2^+	R_2^-	W_1	Bipartite graph
1	1	1	1	2	
v^1	0	1	0	1	$H_{3/2}^1$
v^2	0	1	1	0	$H_{3/2}^1$
v^3	1	0	0	1	$H_{3/2}^1$
v^4	1	0	1	0	$H_{3/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		S_1^+	S_1^-	S_2^+	S_2^-	W_1
	dim	1	1	1	1	2
D_1	1	0	0	1	1	-1
D_2	1	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_1^+	S_1^-	S_2^+	S_2^-	W_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		T_1	W_1
	dim	2	2
P_1P_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_1	W_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 75

Space group $P4$, No. 75, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2
	dim	1	2	1	1	1
Δ_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3
	dim	1	2	1	1	2	1
Λ_1	1	1	0	0	-1	0	0
Λ_2	1	0	1	0	0	-1	0
Λ_3	1	0	0	1	0	0	-1
Λ_4	1	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3
	dim	1	2	1	1	2	1
Σ_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2M_4	M_3	A_1	A_2A_4	A_3
	dim	1	2	1	1	2	1
V_1	1	1	0	0	-1	0	0
V_2	1	0	1	0	0	-1	0
V_3	1	0	0	1	0	0	-1
V_4	1	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2M_4	M_3	A_1	A_2A_4	A_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1	M_2M_4	M_3	X_1	X_2
	dim	1	2	1	1	1
Y_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	M_1	M_2M_4	M_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2Z_4	Z_3	R_1	R_2
	dim	1	2	1	1	1
U_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1	Z_2Z_4	Z_3	R_1	R_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2Z_4	Z_3	A_1	A_2A_4	A_3
	dim	1	2	1	1	2	1
S_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Z_1	Z_2Z_4	Z_3	A_1	A_2A_4	A_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_4	A_3	R_1	R_2
dim		1	2	1	1	1
T_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2A_4	A_3	R_1	R_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

SG 76

Space group $P4_1$, No. 76, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2
	dim	1	2	1	1	1
Δ_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	Bipartite graph
	1	2	1	1	1	
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1Z_4	Z_2Z_3
	dim	1	2	1	2	2
Λ_1	1	1	0	0	-1	0
Λ_2	1	0	1	0	-1	0
Λ_3	1	0	0	1	0	-1
Λ_4	1	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1Z_4	Z_2Z_3	Bipartite graph
	1	2	1	2	2	
v^1	1	1	1	1	1	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3
	dim	1	2	1	1	2	1
Σ_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2M_4	M_3	A_1A_4	A_2A_3
	dim	1	2	1	2	2
V_1	1	1	0	0	-1	0
V_2	1	0	1	0	-1	0
V_3	1	0	0	1	0	-1
V_4	1	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_1	M_2M_4	M_3	A_1A_4	A_2A_3	
	1	2	1	2	2	Bipartite graph
v^1	1	1	1	1	1	$H_{5/4}^3$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2M_4	M_3	X_1	X_2
	dim	1	2	1	1	1
Y_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	M_1	M_2M_4	M_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_4	Z_2Z_3	R_1R_2
dim	2	2	2
U_1U_1	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_4	Z_2Z_3	R_1R_2	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_4	Z_2Z_3	A_1A_4	A_2A_3
dim	2	2	2	2
S_1S_1	2	1	1	-1
			-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_4	Z_2Z_3	A_1A_4	A_2A_3	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_1 A_4$	$A_2 A_3$	$R_1 R_2$
dim	2	2	2
$T_1 T_1$	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$A_1 A_4$	$A_2 A_3$	$R_1 R_2$	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	$R_1 R_2$	X_1	X_2
dim	2	1	1
W_1	1	1	-1
W_2	1	1	0
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$R_1 R_2$	X_1	X_2	
	2	1	1	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 77

Space group $P4_2$, No. 77, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	$\Gamma_2 \Gamma_4$	Γ_3	X_1	X_2
dim	1	2	1	1	1
Δ_1	1	1	2	1	-1
				-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3
dim	1	2	1	1	2	1
Λ_1	1	1	0	0	0	-1
Λ_2	1	0	1	0	-1	0
Λ_3	1	0	0	1	0	-1
Λ_4	1	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3
dim	1	2	1	1	2	1
Σ_1	1	1	2	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2M_4	M_3	A_1	A_2A_4	A_3
	dim	1	2	1	1	2	1
V_1	1	1	0	0	0	-1	0
V_2	1	0	1	0	-1	0	0
V_3	1	0	0	1	0	-1	0
V_4	1	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1	M_2M_4	M_3	A_1	A_2A_4	A_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2M_4	M_3	X_1	X_2
	dim	1	2	1	1	1
Y_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	M_1	M_2M_4	M_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2Z_4	Z_3	R_1	R_2
	dim	1	2	1	1	1
U_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1	Z_2Z_4	Z_3	R_1	R_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2Z_4	Z_3	A_1	A_2A_4	A_3
	dim	1	2	1	1	2	1
S_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Z_1	Z_2Z_4	Z_3	A_1	A_2A_4	A_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_4	A_3	R_1	R_2
dim		1	2	1	1	1
T_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2A_4	A_3	R_1	R_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

SG 78

Space group $P4_3$, No. 78, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2
	dim	1	2	1	1	1
Δ_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	Bipartite graph
	1	2	1	1	1	
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1Z_4	Z_2Z_3
	dim	1	2	1	2	2
Λ_1	1	1	0	0	0	-1
Λ_2	1	0	1	0	-1	0
Λ_3	1	0	0	1	-1	0
Λ_4	1	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1Z_4	Z_2Z_3	Bipartite graph
	1	2	1	2	2	
v^1	1	1	1	1	1	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3
	dim	1	2	1	1	2	1
Σ_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2M_4	M_3	A_1A_4	A_2A_3
	dim	1	2	1	2	2
V_1	1	1	0	0	0	-1
V_2	1	0	1	0	-1	0
V_3	1	0	0	1	-1	0
V_4	1	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_1	M_2M_4	M_3	A_1A_4	A_2A_3	
	1	2	1	2	2	Bipartite graph
v^1	1	1	1	1	1	$H_{5/4}^3$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2M_4	M_3	X_1	X_2
	dim	1	2	1	1	1
Y_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	M_1	M_2M_4	M_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_4	Z_2Z_3	R_1R_2
dim	2	2	2
U_1U_1	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_4	Z_2Z_3	R_1R_2	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_4	Z_2Z_3	A_1A_4	A_2A_3
dim	2	2	2	2
S_1S_1	2	1	1	-1
			-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1Z_4	Z_2Z_3	A_1A_4	A_2A_3	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$A_1 A_4$	$A_2 A_3$	$R_1 R_2$
dim	2	2	2
$T_1 T_1$	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$A_1 A_4$	$A_2 A_3$	$R_1 R_2$	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	$R_1 R_2$	X_1	X_2
dim	2	1	1
W_1	1	-1	0
W_2	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$R_1 R_2$	X_1	X_2	
	2	1	1	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 79

Space group $I4$, No. 79, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2 \Gamma_4$	Γ_3	Z_1	$Z_2 Z_4$	Z_3
dim	1	2	1	1	2	1
Σ_1	1	1	2	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2
dim	1	2	1	1	1
Δ_1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_1	P_1	$P_2 P_2$
	dim	1	1	2
Q_1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	N_1	P_1	$P_2 P_2$	
	1	1	2	Bipartite graph
v^1	1	1	0	$H_{2/1}^1$
v^2	2	0	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1	X_2	P_1	$P_2 P_2$
	dim	1	1	1	2
W_1	1	1	0	-1	0
W_2	1	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	P_1	$P_2 P_2$	
	1	1	1	2	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	Z_1	$Z_2 Z_4$	Z_3
	dim	1	1	1	2	1
Y_1	1	1	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_1	X_2	Z_1	$Z_2 Z_4$	Z_3	
	1	1	1	2	1	Bipartite graph

v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	$H_{2/1}^1$
v^3	0	2	0	1	0	$H_{3/2}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	1	0	0	$H_{2/1}^1$
v^6	1	1	0	1	0	$H_{3/2}^1$
v^7	2	0	0	1	0	$H_{3/2}^1$

SG 80

Space group $I4_1$, No. 80, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	
dim	1	2	1	1	2	1	
Σ_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	Bipartite graph
	1	2	1	1	2	1	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2
dim		1	2	1	1	1
Δ_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_1	P_1P_2
dim		1	2
Q_1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	N_1	P_1P_2	
	1	2	Bipartite graph
v^1	2	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_1	X_2	P_1P_2
dim		1	1	2
W_1	1	1	0	-1
W_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	X_2	$P_1 P_2$	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	Z_1	$Z_2 Z_4$	Z_3
dim	1	1	1	2	1
Y_1	1	1	1	-1	-2 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_1	X_2	Z_1	$Z_2 Z_4$	Z_3	
	1	1	1	2	1	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	$H_{2/1}^1$
v^3	0	2	0	1	0	$H_{3/2}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	1	0	0	$H_{2/1}^1$
v^6	1	1	0	1	0	$H_{3/2}^1$
v^7	2	0	0	1	0	$H_{3/2}^1$

SG 81

Space group $P\bar{4}$, No. 81, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	$\Gamma_2 \Gamma_4$	Γ_3	X_1	X_2
dim	1	2	1	1	1
Δ_1	1	1	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3
dim	1	2	1	1	2	1
Λ_1	1	1	0	1	-1	0
$\Lambda_2\Lambda_2$	2	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3
dim	1	2	1	1	2	1
Σ_1	1	1	2	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	M_1	M_2M_4	M_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2M_4	M_3	A_1	A_2A_4	A_3
	dim	1	2	1	1	2	1
V_1	1	1	0	1	-1	0	-1
V_2V_2	2	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2M_4	M_3	A_1	A_2A_4	A_3	
	1	2	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1	M_2M_4	M_3	X_1	X_2
	dim	1	2	1	1	1
Y_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	M_1	M_2M_4	M_3	X_1	X_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2Z_4	Z_3	R_1	R_2
	dim	1	2	1	1	1
U_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1	Z_2Z_4	Z_3	R_1	R_2	
	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2Z_4	Z_3	A_1	A_2A_4	A_3
	dim	1	2	1	1	2	1
S_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Z_1	Z_2Z_4	Z_3	A_1	A_2A_4	A_3	Bipartite graph
	1	2	1	1	2	1	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_4	A_3	R_1	R_2
dim		1	2	1	1	1
T_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2A_4	A_3	R_1	R_2	Bipartite graph
	1	2	1	1	1	
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

SG 82

Space group $I\bar{4}$, No. 82, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	
dim	1	2	1	1	2	1	
Σ_1	1	1	2	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	Z_1	Z_2Z_4	Z_3	Bipartite graph
	1	2	1	1	2	1	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	2	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/2}^1$
v^6	0	1	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	
dim	1	2	1	1	1	
Δ_1	1	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	$\Gamma_2\Gamma_4$	Γ_3	X_1	X_2	Bipartite graph
	1	2	1	1	1	
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	0	2	0	$H_{3/2}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	Z_1	Z_2	Z_3
dim	1	1	1	2	1
Y_1	1	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_1	X_2	Z_1	Z_2	Z_3	Bipartite graph
	1	1	1	2	1	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	$H_{2/1}^1$
v^3	0	2	0	1	0	$H_{3/2}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	1	0	0	$H_{2/1}^1$
v^6	1	1	0	1	0	$H_{3/2}^1$
v^7	2	0	0	1	0	$H_{3/2}^1$

SG 83

Space group $P4/m$, No. 83, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1^+	X_1^-	X_2^+	X_2^-
dim	1	1	1	1	2	2	1	1	1	1
Δ_1	1	0	1	0	0	2	-1	0	0	-1
Δ_2	1	0	1	0	2	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1^+	X_1^-	X_2^+	X_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$

v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$
	dim	1	1	1	1	2	2	1	1	1	1	2	2
Λ_1	1	1	1	0	0	0	0	-1	-1	0	0	0	0
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	1	0	0	0	0	-1	-1
Λ_3	1	0	0	1	1	0	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$
	dim	1	1	1	1	2	2	1	1	1	1	2	2
Σ_1	1	1	0	1	0	0	2	-1	0	-1	0	0	-2
Σ_2	1	0	1	0	1	2	0	0	-1	0	-1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 22.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	2	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	2	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^6	0	0	0	0	1	0	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	2	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	2	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	2	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	2	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+A_4^+$	$A_3^-A_4^-$
	dim	1	1	1	1	2	2	1	1	1	1	2	2
V_1	1	1	1	0	0	0	0	-1	-1	0	0	0	0
V_2V_4	2	0	0	0	0	1	1	0	0	0	0	-1	-1
V_3	1	0	0	1	1	0	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+A_4^+$	$A_3^-A_4^-$	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	X_1^+	X_1^-	X_2^+	X_2^-
dim		1	1	1	1	2	2	1	1	1	1
Y_1	1	1	0	1	0	0	2	-1	0	0	-1
Y_2	1	0	1	0	1	2	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	X_1^+	X_1^-	X_2^+	X_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$

v^{12}	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	R_1^+	R_1^-	R_2^+	R_2^-
dim		1	1	1	1	2	2	1	1	1	1
U_1	1	1	0	1	0	0	2	-1	0	0	-1
U_2	1	0	1	0	1	2	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	R_1^+	R_1^-	R_2^+	R_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+ A_4^+$	$A_3^- A_4^-$
dim		1	1	1	1	2	2	1	1	1	1	2	2
S_1	1	1	0	1	0	0	2	-1	0	-1	0	0	-2
S_2	1	0	1	0	1	2	0	0	-1	0	-1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 22.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+ A_4^+$	$A_3^- A_4^-$	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	2	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	2	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^6	0	0	0	0	1	0	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	2	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	2	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	2	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	2	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+ A_4^+$	$A_3^- A_4^-$	R_1^+	R_1^-	R_2^+	R_2^-
dim		1	1	1	1	2	2	1	1	1	1
T_1	1	1	0	1	0	0	2	-1	0	0	-1
T_2	1	0	1	0	1	2	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+ A_4^+$	$A_3^- A_4^-$	R_1^+	R_1^-	R_2^+	R_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$

v^4	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 84

Space group $P4_2/m$, No. 84, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1^+	X_1^-	X_2^+	X_2^-
dim		1	1	1	1	2	2	1	1	1	1
Δ_1	1	1	0	1	0	0	2	-1	0	0	-1
Δ_2	1	0	1	0	1	2	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1^+	X_1^-	X_2^+	X_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1	Z_2
dim		1	1	1	1	2	2	2	2
Λ_1	1	1	1	0	0	0	0	0	-1
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	1	-1	0
Λ_3	1	0	0	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1	Z_2	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$
dim		1	1	1	1	2	2	1	1	1	1	2	2
Σ_1	1	1	0	1	0	0	2	-1	0	-1	0	0	-2
Σ_2	1	0	1	0	1	2	0	0	-1	0	-1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 22.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	
	1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	2	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	2	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^6	0	0	0	0	1	0	0	0	0	2	0	0	$H_{3/2}^1$

v^7	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	2	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	2	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	2	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	2	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	A_1	A_2
	dim	1	1	1	1	2	2	2	2
V_1	1	1	1	0	0	0	0	0	-1
V_2V_4	2	0	0	0	0	1	1	-1	0
V_3	1	0	0	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	A_1	A_2	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	X_1^+	X_1^-	X_2^+	X_2^-
dim		1	1	1	1	2	2	1	1	1	1
Y_1	1	1	0	1	0	0	2	-1	0	0	-1
Y_2	1	0	1	0	1	2	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	M_1^+	M_1^-	M_2^+	M_2^-	$M_3^+M_4^+$	$M_3^-M_4^-$	X_1^+	X_1^-	X_2^+	X_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U : $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	R_1^+	R_1^-	R_2^+	R_2^-
dim		2	2	1	1	1	1
U_1	1	1	1	-1	0	0	-1
U_2	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1	Z_2	R_1^+	R_1^-	R_2^+	R_2^-	Bipartite graph
	2	2	1	1	1	1	
v^1	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	1	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$

v^6	1	0	0	1	0	1	$H_{3/2}^1$
v^7	1	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	1	1	0	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	A_1	A_2
dim		2	2	2	2
S_1	1	1	1	-1	-1
S_2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	A_1	A_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	R_1^+	R_1^-	R_2^+	R_2^-
dim		2	2	1	1	1	1
T_1	1	1	1	-1	0	0	-1
T_2	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_1	A_2	R_1^+	R_1^-	R_2^+	R_2^-	
	2	2	1	1	1	1	Bipartite graph
v^1	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	1	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	1	$H_{3/2}^1$
v^7	1	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	1	1	0	0	$H_{3/2}^1$

SG 85

Space group $P4/n$, No. 85, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1
	dim	1	1	1	1	2	2	2
Δ_1	1	1	0	1	0	0	2	-1
Δ_2	1	0	1	0	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	0	1	1	2	$H_{4/4}^2$
v^2	0	0	0	2	0	1	2	$H_{5/4}^3$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	1	0	1	0	1	2	$H_{5/4}^3$
v^6	0	1	1	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	2	$H_{5/4}^3$
v^8	1	0	0	1	0	0	1	$H_{3/2}^1$
v^9	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{10}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	2	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$
	dim	1	1	1	1	2	2	1	1	1	1	2	2
Λ_1	1	1	1	0	0	0	0	-1	-1	0	0	0	0
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	1	0	0	0	0	-1	-1
Λ_3	1	0	0	1	1	0	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1	M_2
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	1	0	0	2	-1	-1
Σ_2	1	0	1	0	1	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1	M_2	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	1	0	2	$H_{4/4}^2$
v^2	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	1	1	2	0	$H_{4/4}^2$
v^4	0	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	0	0	2	0	1	1	1	$H_{5/4}^3$
v^6	0	0	0	2	0	1	2	0	$H_{5/4}^3$
v^7	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^9	0	0	2	0	1	0	0	2	$H_{5/4}^3$
v^{10}	0	0	2	0	1	0	1	1	$H_{5/4}^3$
v^{11}	0	0	2	0	1	0	2	0	$H_{5/4}^3$
v^{12}	0	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{13}	0	1	0	1	0	1	1	1	$H_{5/4}^3$

v^{14}	0	1	0	1	0	1	2	0	$H_{5/4}^3$
v^{15}	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	2	0	0	0	1	0	2	$H_{5/4}^3$
v^{18}	0	2	0	0	0	1	1	1	$H_{5/4}^3$
v^{19}	0	2	0	0	0	1	2	0	$H_{5/4}^3$
v^{20}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{21}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{22}	1	0	1	0	1	0	0	2	$H_{5/4}^3$
v^{23}	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{24}	1	0	1	0	1	0	2	0	$H_{5/4}^3$
v^{25}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{27}	2	0	0	0	1	0	0	2	$H_{5/4}^3$
v^{28}	2	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{29}	2	0	0	0	1	0	2	0	$H_{5/4}^3$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_1	M_2	A_1	A_2
dim	2	2	2	2
$V_1 V_2$	2	1	0	-1 0
V_3	1	0	1	0 -1
V_4	1	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1	M_2	A_1	A_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_1	M_2	X_1
dim	2	2	2

Y_1	1	1	1	-1
Y_2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1	M_2	X_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	R_1
	dim	1	1	1	1	2	2	2
U_1	1	1	0	1	0	0	2	-1
U_2	1	0	1	0	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	R_1	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	0	1	1	2	$H_{4/4}^2$
v^2	0	0	0	2	0	1	2	$H_{5/4}^3$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	1	0	1	0	1	2	$H_{5/4}^3$
v^6	0	1	1	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	2	$H_{5/4}^3$
v^8	1	0	0	1	0	0	1	$H_{3/2}^1$
v^9	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{10}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	2	$H_{5/4}^3$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix



corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_4^+	Z_3^-	Z_4^-	A_1	A_2
	dim	1	1	1	1	2		2		2	2
S_1	1	1	0	1	0	0		2		-1	-1
S_2	1	0	1	0	1	2		0		-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_4^+	Z_3^-	Z_4^-	A_1	A_2	
	1	1	1	1	2		2		2	2	Bipartite graph
v^1	0	0	0	0	1		1		0	2	$H_{4/4}^2$
v^2	0	0	0	0	1		1		1	1	$H_{4/4}^2$
v^3	0	0	0	0	1		1		2	0	$H_{4/4}^2$
v^4	0	0	0	2	0		1		0	2	$H_{5/4}^3$
v^5	0	0	0	2	0		1		1	1	$H_{5/4}^3$
v^6	0	0	0	2	0		1		2	0	$H_{5/4}^3$
v^7	0	0	1	1	0		0		0	1	$H_{3/2}^1$
v^8	0	0	1	1	0		0		1	0	$H_{3/2}^1$
v^9	0	0	2	0	1		0		0	2	$H_{5/4}^3$
v^{10}	0	0	2	0	1		0		1	1	$H_{5/4}^3$
v^{11}	0	0	2	0	1		0		2	0	$H_{5/4}^3$
v^{12}	0	1	0	1	0		1		0	2	$H_{5/4}^3$
v^{13}	0	1	0	1	0		1		1	1	$H_{5/4}^3$
v^{14}	0	1	0	1	0		1		2	0	$H_{5/4}^3$
v^{15}	0	1	1	0	0		0		0	1	$H_{3/2}^1$
v^{16}	0	1	1	0	0		0		1	0	$H_{3/2}^1$
v^{17}	0	2	0	0	0		1		0	2	$H_{5/4}^3$
v^{18}	0	2	0	0	0		1		1	1	$H_{5/4}^3$
v^{19}	0	2	0	0	0		1		2	0	$H_{5/4}^3$
v^{20}	1	0	0	1	0		0		0	1	$H_{3/2}^1$
v^{21}	1	0	0	1	0		0		1	0	$H_{3/2}^1$
v^{22}	1	0	1	0	1		0		0	2	$H_{5/4}^3$
v^{23}	1	0	1	0	1		0		1	1	$H_{5/4}^3$
v^{24}	1	0	1	0	1		0		2	0	$H_{5/4}^3$
v^{25}	1	1	0	0	0		0		0	1	$H_{3/2}^1$
v^{26}	1	1	0	0	0		0		1	0	$H_{3/2}^1$
v^{27}	2	0	0	0	1		0		0	2	$H_{5/4}^3$
v^{28}	2	0	0	0	1		0		1	1	$H_{5/4}^3$
v^{29}	2	0	0	0	1		0		2	0	$H_{5/4}^3$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	R_1
	dim	2	2	2
T_1	1	1	1	-1
T_2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1	A_2	R_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_1	X_1
	dim	2	2
W_1W_2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 86

Space group $P4_2/n$, No. 86, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1
	dim	1	1	1	1	2	2	2
Δ_1^+	1	1	0	1	0	0	2	-1
Δ_1^-	1	0	1	0	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	0	1	1	2	$H_{4/4}^2$
v^2	0	0	0	2	0	1	2	$H_{5/4}^3$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	1	0	1	0	1	2	$H_{5/4}^3$
v^6	0	1	1	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	2	$H_{5/4}^3$
v^8	1	0	0	1	0	0	1	$H_{3/2}^1$
v^9	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{10}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	2	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1	Z_2
	dim	1	1	1	1	2	2	2	2
Λ_1	1	1	1	0	0	0	0	0	-1
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	1	-1	0
Λ_3	1	0	0	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1	Z_2	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1	M_2
dim		1	1	1	1	2	2	2	2
Σ_1	1	1	0	1	0	0	2	-1	-1
Σ_2	1	0	1	0	1	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	M_1	M_2	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	1	0	2	$H_{4/4}^2$
v^2	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	1	1	2	0	$H_{4/4}^2$
v^4	0	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	0	0	2	0	1	1	1	$H_{5/4}^3$
v^6	0	0	0	2	0	1	2	0	$H_{5/4}^3$
v^7	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^9	0	0	2	0	1	0	0	2	$H_{5/4}^3$
v^{10}	0	0	2	0	1	0	1	1	$H_{5/4}^3$
v^{11}	0	0	2	0	1	0	2	0	$H_{5/4}^3$
v^{12}	0	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{13}	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^{14}	0	1	0	1	0	1	2	0	$H_{5/4}^3$
v^{15}	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	2	0	0	0	1	0	2	$H_{5/4}^3$
v^{18}	0	2	0	0	0	1	1	1	$H_{5/4}^3$
v^{19}	0	2	0	0	0	1	2	0	$H_{5/4}^3$
v^{20}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{21}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{22}	1	0	1	0	1	0	0	2	$H_{5/4}^3$
v^{23}	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{24}	1	0	1	0	1	0	2	0	$H_{5/4}^3$
v^{25}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{27}	2	0	0	0	1	0	0	2	$H_{5/4}^3$
v^{28}	2	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{29}	2	0	0	0	1	0	2	0	$H_{5/4}^3$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_1	M_2	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+A_4^+$	$A_3^-A_4^-$
dim	2	2	1	1	1	1	2	2
V_1V_2	2	1 0	0	0	0	0	-1	-1
V_3	1	0 1	-1	-1	0	0	0	0
V_4	1	0 1	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1	M_2	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+A_4^+$	$A_3^-A_4^-$	Bipartite graph
	2	2	1	1	1	1	2	2	
v^1	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^2	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^4	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^5	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep	M_1	M_2	X_1
dim	2	2	2
Y_1^+	1	1	1
Y_1^-	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1	M_2	X_1	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	R_1
	dim	2	2	2
U_1	1	1	1	-1
U_2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	Z_2	R_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_4^+	A_3^-	A_4^-
	dim	2	2	1	1	1	1	2		2	
S_1	1	1	1	0	-1	0	-1	-2		0	
S_2	1	1	1	-1	0	-1	0	0		-2	

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Z_1	Z_2	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_4^+	A_3^-	A_4^-	
	2	2	1	1	1	1	2		2		Bipartite graph
v^1	0	1	0	0	1	1	0		0		$H_{3/2}^1$
v^2	0	1	0	1	1	0	0		0		$H_{3/2}^1$
v^3	0	1	1	0	0	1	0		0		$H_{3/2}^1$
v^4	0	1	1	1	0	0	0		0		$H_{3/2}^1$
v^5	0	2	0	0	0	0	1		1		$H_{4/4}^2$
v^6	0	2	0	0	0	2	0		1		$H_{5/4}^3$
v^7	0	2	0	0	2	0	1		0		$H_{5/4}^3$
v^8	0	2	0	1	0	1	0		1		$H_{5/4}^3$
v^9	0	2	0	2	0	0	0		1		$H_{5/4}^3$
v^{10}	0	2	1	0	1	0	1		0		$H_{5/4}^3$
v^{11}	0	2	2	0	0	0	1		0		$H_{5/4}^3$
v^{12}	1	0	0	0	1	1	0		0		$H_{3/2}^1$
v^{13}	1	0	0	1	1	0	0		0		$H_{3/2}^1$
v^{14}	1	0	1	0	0	1	0		0		$H_{3/2}^1$
v^{15}	1	0	1	1	0	0	0		0		$H_{3/2}^1$
v^{16}	1	1	0	0	0	0	1		1		$H_{4/4}^2$
v^{17}	1	1	0	0	0	2	0		1		$H_{5/4}^3$

v^{18}	1	1	0	0	2	0	1	0	$H_{5/4}^3$
v^{19}	1	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{20}	1	1	0	2	0	0	0	1	$H_{5/4}^3$
v^{21}	1	1	1	0	1	0	1	0	$H_{5/4}^3$
v^{22}	1	1	2	0	0	0	1	0	$H_{5/4}^3$
v^{23}	2	0	0	0	0	0	1	1	$H_{4/4}^2$
v^{24}	2	0	0	0	0	2	0	1	$H_{5/4}^3$
v^{25}	2	0	0	0	2	0	1	0	$H_{5/4}^3$
v^{26}	2	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{27}	2	0	0	2	0	0	0	1	$H_{5/4}^3$
v^{28}	2	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{29}	2	0	2	0	0	0	1	0	$H_{5/4}^3$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+A_4^+$	$A_3^-A_4^-$	R_1
	dim	1	1	1	1	2	2	2
T_1	1	0	1	0	1	2	0	-1
T_2	1	1	0	1	0	0	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	A_1^+	A_1^-	A_2^+	A_2^-	$A_3^+A_4^+$	$A_3^-A_4^-$	R_1	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	0	1	1	2	$H_{4/4}^2$
v^2	0	0	0	2	0	1	2	$H_{5/4}^3$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	1	0	1	0	1	2	$H_{5/4}^3$
v^6	0	1	1	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	2	$H_{5/4}^3$
v^8	1	0	0	1	0	0	1	$H_{3/2}^1$
v^9	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{10}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	2	$H_{5/4}^3$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	R_1	X_1
dim	2	2
W_1W_2	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 87

Space group $I4/m$, No. 87, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$
dim		1	1	1	1	2	2	1	1	1	1	2	2
Σ_1	1	1	0	1	0	0	2	-1	0	-1	0	0	-2
Σ_2	1	0	1	0	1	2	0	0	-1	0	-1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 22.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	2	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	2	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^6	0	0	0	0	1	0	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	2	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	2	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$

v^{16}	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	2	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	2	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1^+	X_1^-	X_2^+	X_2^-
dim		1	1	1	1	2	2	1	1	1	1
Δ_1	1	1	0	1	0	0	2	-1	0	0	-1
Δ_2	1	0	1	0	1	2	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1^+	X_1^-	X_2^+	X_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_1^+	N_1^-	P_1	P_2P_4	P_3
	dim	1	1	1	2	1
Q_1	1	1	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	N_1^+	N_1^-	P_1	P_2P_4	P_3	Bipartite graph
	1	1	1	2	1	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	$H_{2/1}^1$
v^3	0	2	0	1	0	$H_{3/2}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	1	0	0	$H_{2/1}^1$
v^6	1	1	0	1	0	$H_{3/2}^1$
v^7	2	0	0	1	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	P_1	P_2P_4	P_3
	dim	1	1	1	1	1	2	1
W_1	1	1	1	0	0	-1	0	-1
W_2	1	0	0	1	1	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_1^+	X_1^-	X_2^+	X_2^-	P_1	P_2P_4	P_3	Bipartite graph
	1	1	1	1	1	2	1	
v^1	0	0	0	2	0	1	0	$H_{3/2}^1$
v^2	0	0	1	1	0	1	0	$H_{3/2}^1$
v^3	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$
	dim	1	1	1	1	1	1	1	1	2	2
Y_1	1	1	0	0	1	-1	0	-1	0	0	-2
Y_2	1	0	1	1	0	0	-1	0	-1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	X_1^+	X_1^-	X_2^+	X_2^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	0	0	$H_{2/1}^1$
v^3	0	0	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^6	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	2	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 88

Space group $I4_1/a$, No. 88, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1	Z_2
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	1	0	0	2	-1	-1
Σ_2	1	0	1	0	1	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	Z_1	Z_2	Bipartite graph
	1	1	1	1	2	2	2	2	

v^1	0	0	0	0	1	1	0	2	$H_{4/4}^2$
v^2	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	1	1	2	0	$H_{4/4}^2$
v^4	0	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	0	0	2	0	1	1	1	$H_{5/4}^3$
v^6	0	0	0	2	0	1	2	0	$H_{5/4}^3$
v^7	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^9	0	0	2	0	1	0	0	2	$H_{5/4}^3$
v^{10}	0	0	2	0	1	0	1	1	$H_{5/4}^3$
v^{11}	0	0	2	0	1	0	2	0	$H_{5/4}^3$
v^{12}	0	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{13}	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^{14}	0	1	0	1	0	1	2	0	$H_{5/4}^3$
v^{15}	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	2	0	0	0	1	0	2	$H_{5/4}^3$
v^{18}	0	2	0	0	0	1	1	1	$H_{5/4}^3$
v^{19}	0	2	0	0	0	1	2	0	$H_{5/4}^3$
v^{20}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{21}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{22}	1	0	1	0	1	0	0	2	$H_{5/4}^3$
v^{23}	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{24}	1	0	1	0	1	0	2	0	$H_{5/4}^3$
v^{25}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{27}	2	0	0	0	1	0	0	2	$H_{5/4}^3$
v^{28}	2	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{29}	2	0	0	0	1	0	2	0	$H_{5/4}^3$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1
	dim	1	1	1	1	2	2	2
Δ_1	1	1	0	1	0	0	2	-1
Δ_2	1	0	1	0	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	X_1	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	0	1	1	2	$H_{4/4}^2$
v^2	0	0	0	2	0	1	2	$H_{5/4}^3$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	2	0	1	0	2	$H_{5/4}^3$
v^5	0	1	0	1	0	1	2	$H_{5/4}^3$
v^6	0	1	1	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	2	$H_{5/4}^3$
v^8	1	0	0	1	0	0	1	$H_{3/2}^1$
v^9	1	0	1	0	1	0	2	$H_{5/4}^3$
v^{10}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	1	0	2	$H_{5/4}^3$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	N_1^+	N_1^-	P_1P_4	P_2P_3
dim	1	1	2	2
Q_1	1	1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	N_1^+	N_1^-	P_1P_4	P_2P_3	
	1	1	2	2	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	0	2	1	0	$H_{3/2}^1$
v^3	1	1	0	1	$H_{3/2}^1$
v^4	1	1	1	0	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$
v^6	2	0	1	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	X_1	P_1P_4	P_2P_3
dim	2	2	2
W_1W_2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	P_1P_4	P_2P_3	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	Z_1	Z_2
	dim	2	2	2
Y_1	1	1	-1	-1
Y_2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	Z_1	Z_2	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 89

Space group $P422$, No. 89, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	-1	0	0
Δ_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	1	0	0	0	-1	-1	0	0	0
Λ_2	1	0	0	0	0	1	0	0	0	0	-1
Λ_3	1	0	0	1	1	0	0	0	-1	-1	0
Λ_4	1	0	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
V_1	1	1	1	0	0	0	-1	-1	0	0	0
V_2	1	0	0	0	0	1	0	0	0	0	-1
V_3	1	0	0	1	1	0	0	0	-1	-1	0
V_4	1	0	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	0	0	-1
Y_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
U_1	1	1	0	1	0	1	-1	-1	0	0
U_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S : $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
S_1	1	1	0	0	1	1	-1	0	0	-1	-1
S_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1	A_2	A_3	A_4	A_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$

v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	A_4	A_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
T_1	1	1	0	1	0	1	-1	0	0	-1
T_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 90

Space group $P4_212$, No. 90, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	1	0	0	0	-1	-1	0	0	0
Λ_2	1	0	0	0	0	1	0	0	0	0	-1
Λ_3	1	0	0	1	1	0	0	0	-1	-1	0
Λ_4	1	0	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
dim	1	1	1	1	2	2	2	2	
Σ_1	1	1	0	0	1	1	-2	0	-1
Σ_2	1	0	1	1	0	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5	
dim	2	2	2	2	2	2	
V_1V_3	2	0	0	1	0	0	-1
V_2V_4	2	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_1M_2	M_3M_4	M_5	X_1
dim	2	2	2	2
Y_1Y_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	Z_3	Z_4	Z_5	R_1
dim	1	1	1	1	2	2
U_1	1	1	0	1	0	1
U_2	1	0	1	0	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5	
dim	1	1	1	1	2	2	2	2	
S_1	1	1	0	0	1	1	-2	0	-1
S_2	1	0	1	1	0	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3A_4	A_5	R_1	
dim	2	2	2	2	
T_1T_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_4	A_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	X_1
dim	2	2
W_1W_2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 91

Space group $P4_122$, No. 91, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
dim		1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	-1	0	0
Δ_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph

v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2
	dim	1	1	1	1	2	2	2
Λ_1	1	1	1	0	0	0	-1	0
Λ_2	1	0	0	0	0	1	-1	0
Λ_3	1	0	0	1	1	0	0	-1
Λ_4	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	1	$H_{5/4}^3$
v^2	0	1	1	0	1	1	1	$H_{5/4}^3$
v^3	1	0	0	1	1	1	1	$H_{5/4}^3$
v^4	1	0	1	0	1	1	1	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1	A_2
	dim	1	1	1	1	2	2	2
V_1	1	1	1	0	0	0	-1	0
V_2	1	0	0	0	0	1	-1	0
V_3	1	0	0	1	1	0	0	-1
V_4	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	M_5	A_1	A_2	Bipartite graph
	1	1	1	1	2	2	2	
v^1	0	1	0	1	1	1	1	$H_{5/4}^3$
v^2	0	1	1	0	1	1	1	$H_{5/4}^3$
v^3	1	0	0	1	1	1	1	$H_{5/4}^3$
v^4	1	0	1	0	1	1	1	$H_{5/4}^3$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	0	0	-1
Y_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	R_1
	dim	2	2	2
U_1U_2	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	Z_2	R_1	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	A_1	A_2
dim	2	2	2	2
S_1S_2	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	A_1	A_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2	R_1
dim	2	2	2
T_1T_2	2	1	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1	A_2	R_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	X_1	X_2	X_3	X_4
dim	2	1	1	1	1
W_1	1	1	-1	0	-1 0
W_2	1	1	0	-1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

R_1	X_1	X_2	X_3	X_4	
2	1	1	1	1	Bipartite graph
v^1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	1	0	0	$H_{3/2}^1$
v^4	1	1	1	0	$H_{3/2}^1$

SG 92

Space group $P4_12_12$, No. 92, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2
	dim	1	1	1	1	2	2	2
Λ_1	1	1	1	0	0	0	-1	0
Λ_2	1	0	0	0	0	1	-1	0
Λ_3	1	0	0	1	1	0	0	-1
Λ_4	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	1	$H_{5/4}^3$
v^2	0	1	1	0	1	1	1	$H_{5/4}^3$
v^3	1	0	0	1	1	1	1	$H_{5/4}^3$
v^4	1	0	1	0	1	1	1	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	0	1	1	-2	0	-1
Σ_2	1	0	1	1	0	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1M_2	M_3M_4	M_5	A_1A_2
	dim	2	2	2	4
V_1V_3	2	0	0	1	-1
V_2V_4	2	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1M_2	M_3M_4	M_5	A_1A_2	
	2	2	2	4	Bipartite graph
v^1	0	1	1	1	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1M_2	M_3M_4	M_5	X_1
	dim	2	2	2	2
Y_1Y_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	R_1R_2	R_3R_4
	dim	2	2	2	2
U_1U_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	A_1A_2
dim	2	2	4
S_1S_2	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	A_1A_2	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	R_1R_2	R_3R_4
dim	4	2	2
T_1T_1	2	1	-1
T_2T_2	2	1	0
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_1A_2	R_1R_2	R_3R_4	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1R_2	R_3R_4	X_1
dim	2	2	2
W_1W_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1R_2	R_3R_4	X_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 93

Space group $P4_222$, No. 93, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	-1	0
Δ_2	1	0	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	1	0	0	0	0	0	0	0	-1
Λ_2	1	0	0	0	0	1	-1	-1	0	0	0
Λ_3	1	0	0	1	1	0	0	0	0	0	-1
Λ_4	1	0	0	0	0	1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	

v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
V_1	1	1	1	0	0	0	0	0	0	0	-1
V_2	1	0	0	0	0	1	-1	-1	0	0	0
V_3	1	0	0	1	1	0	0	0	0	0	-1
V_4	1	0	0	0	0	1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	0	0	-1
Y_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
U_1	1	0	1	0	1	1	-1	-1	0	0
U_2	1	1	0	1	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$

v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S : $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
S_1	1	1	0	0	1	1	-1	0	0	-1	-1
S_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1	A_2	A_3	A_4	A_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
T_1	1	1	0	1	0	1	-1	0	0	-1
T_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 94

Space group $P4_22_12$, No. 94, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	1	0	0	0	0	0	0	0	-1
Λ_2	1	0	0	0	0	1	-1	-1	0	0	0
Λ_3	1	0	0	1	1	0	0	0	0	0	-1
Λ_4	1	0	0	0	0	1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	0	1	1	-2	0	-1
Σ_2	1	0	1	1	0	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
V_1V_3	2	0	0	1	-1	-1	0
V_2V_4	2	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	1	0	0	0	0	1	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_1M_2	M_3M_4	M_5	X_1
dim	2	2	2	2
Y_1Y_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	Z_3	Z_4	Z_5	R_1
dim	1	1	1	1	2	2
U_1	1	0	1	0	1	1
U_2	1	1	0	1	0	1
						-1
						-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
S_1	1	1	0	0	1	1	0	-2	-1
S_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1A_2	A_3A_4	A_5	R_1
	dim	2	2	2	2
T_1T_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_4	A_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	X_1
dim	2	2
W_1W_2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 95

Space group $P4_322$, No. 95, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
dim	1	1	1	1	2	1	1	1	1	
Δ_1	1	1	0	1	0	1	-1	-1	0	0
Δ_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2
	dim	1	1	1	1	2	2	2
Λ_1	1	1	1	0	0	0	0	-1
Λ_2	1	0	0	0	0	1	-1	0
Λ_3	1	0	0	1	1	0	-1	0
Λ_4	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	1	$H_{5/4}^3$
v^2	0	1	1	0	1	1	1	$H_{5/4}^3$
v^3	1	0	0	1	1	1	1	$H_{5/4}^3$
v^4	1	0	1	0	1	1	1	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$

v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1	A_2
	dim	1	1	1	1	2	2	2
V_1	1	1	1	0	0	0	0	-1
V_2	1	0	0	0	0	1	-1	0
V_3	1	0	0	1	1	0	-1	0
V_4	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	M_5	A_1	A_2	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	1	$H_{5/4}^3$
v^2	0	1	1	0	1	1	1	$H_{5/4}^3$
v^3	1	0	0	1	1	1	1	$H_{5/4}^3$
v^4	1	0	1	0	1	1	1	$H_{5/4}^3$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	0	0	-1
Y_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	R_1
dim	2	2	2
U_1U_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	Z_2	R_1	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	A_1	A_2
dim	2	2	2	2
S_1S_2	2	1	1	-1
			-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	A_1	A_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2	R_1
dim	2	2	2
$T_1 T_2$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1	A_2	R_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	R_1	X_1	X_2	X_3	X_4
dim	2	1	1	1	1
W_1	1	-1	0	-1	0
W_2	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_1	X_1	X_2	X_3	X_4	
	2	1	1	1	1	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

SG 96

Space group $P4_32_12$, No. 96, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2
	dim	1	1	1	1	2	2	2
Λ_1	1	1	1	0	0	0	0	-1
Λ_2	1	0	0	0	0	1	-1	0
Λ_3	1	0	0	1	1	0	-1	0
Λ_4	1	0	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	1	$H_{5/4}^3$
v^2	0	1	1	0	1	1	1	$H_{5/4}^3$
v^3	1	0	0	1	1	1	1	$H_{5/4}^3$
v^4	1	0	1	0	1	1	1	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
dim	1	1	1	1	2	2	2	2	
Σ_1	1	1	0	0	1	1	-2	0	-1
Σ_2	1	0	1	1	0	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_1M_2	M_3M_4	M_5	A_1A_2	
dim	2	2	2	4	
V_1V_3	2	0	0	1	-1
V_2V_4	2	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1M_2	M_3M_4	M_5	A_1A_2	
	2	2	2	4	Bipartite graph
v^1	0	1	1	1	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y : $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_1M_2	M_3M_4	M_5	X_1
dim	2	2	2	2
Y_1Y_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U : $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	R_1R_2	R_3R_4
dim	2	2	2	2
U_1U_2	2	1	1	-1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	R_1R_2	R_3R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	$A_1 A_2$
dim	2	2	4
$S_1 S_2$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	$A_1 A_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_1 A_2$	$R_1 R_2$	$R_3 R_4$
dim	4	2	2
$T_1 T_1$	2	1	-1
$T_2 T_2$	2	1	0
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$A_1 A_2$	$R_1 R_2$	$R_3 R_4$	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	$R_1 R_2$	$R_3 R_4$	X_1
dim	2	2	2
$W_1 W_2$	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1R_2	R_3R_4	X_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 97

Space group $I422$, No. 97, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	1	0	1	-1	0	-1	0	-1
Σ_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
dim	1	1	1	1	2	1	1	1	1	
Δ_1	1	1	0	0	1	1	-1	0	0	-1
Δ_2	1	0	1	1	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	N_1	N_2	P_1	P_2	P_3
dim	1	1	1	2	1
Q_1	1	1	1	0	0
Q_2	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	N_1	N_2	P_1	P_2	P_3	Bipartite graph
	1	1	1	2	1	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1	X_2	X_3	X_4	P_1	P_2P_4	P_3
	dim	1	1	1	1	1	2	1
W_1	1	1	0	1	0	-1	0	-1
W_2	1	0	1	0	1	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_1	X_2	X_3	X_4	P_1	P_2P_4	P_3	Bipartite graph
	1	1	1	1	1	2	1	
v^1	0	0	0	2	0	1	0	$H_{3/2}^1$
v^2	0	0	1	0	0	0	1	$H_{2/1}^1$
v^3	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	1	0	$H_{3/2}^1$
v^5	0	2	0	0	0	1	0	$H_{3/2}^1$
v^6	1	0	0	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	1	1	1	1	2
Y_1	1	1	1	0	0	-1	0	0	-1	-1
Y_2	1	0	0	1	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	1	1	1	1	2	
v^1	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^7	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$

v^9	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 98

Space group $I4_122$, No. 98, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	1	0	1	0	-1	0	-1	-1
Σ_2	1	0	1	0	1	1	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	0	1	1	-1	0	0	-1
Δ_2	1	0	1	1	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_1	N_2	P_1
	dim	1	1	2
Q_1	1	1	0	-1
Q_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	N_1	N_2	P_1	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1	X_2	X_3	X_4	P_1
	dim	1	1	1	1	2
W_1	1	1	0	1	0	-1
W_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	X_3	X_4	P_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	1	1	1	1	2
Y_1	1	1	1	0	0	0	-1	-1	0	-1
Y_2	1	0	0	1	1	-1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

SG 99

Space group $P4mm$, No. 99, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	0	0	-1
Δ_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	0	0	0	0	-1	0	0	0	0
Λ_2	1	0	1	0	0	0	0	-1	0	0	0
Λ_3	1	0	0	1	0	0	0	0	-1	0	0
Λ_4	1	0	0	0	1	0	0	0	0	-1	0
Λ_5	2	0	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
V_1	1	1	0	0	0	0	-1	0	0	0	0
V_2	1	0	1	0	0	0	0	-1	0	0	0
V_3	1	0	0	1	0	0	0	0	-1	0	0
V_4	1	0	0	0	1	0	0	0	0	-1	0
V_5	2	0	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	-1	0	0
Y_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$

v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
U_1	1	1	0	1	0	1	-1	0	0	-1
U_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
S_1	1	1	0	0	1	1	-1	0	0	-1	-1
S_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Z ₁	Z ₂	Z ₃	Z ₄	Z ₅	A ₁	A ₂	A ₃	A ₄	A ₅	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v ¹	0	0	0	0	1	0	0	0	0	1	H _{2/2} ¹
v ²	0	0	0	0	1	0	0	1	1	0	H _{3/2} ¹
v ³	0	0	0	0	1	0	1	0	1	0	H _{3/2} ¹
v ⁴	0	0	0	0	1	1	0	1	0	0	H _{3/2} ¹
v ⁵	0	0	0	0	1	1	1	0	0	0	H _{3/2} ¹
v ⁶	0	0	0	1	0	0	0	0	1	0	H _{2/1} ¹
v ⁷	0	0	0	1	0	1	0	0	0	0	H _{2/1} ¹
v ⁸	0	0	1	0	0	0	0	1	0	0	H _{2/1} ¹
v ⁹	0	0	1	0	0	0	1	0	0	0	H _{2/1} ¹
v ¹⁰	0	0	1	1	0	0	0	0	0	1	H _{3/2} ¹
v ¹¹	0	1	0	0	0	0	0	1	0	0	H _{2/1} ¹
v ¹²	0	1	0	0	0	0	1	0	0	0	H _{2/1} ¹
v ¹³	0	1	0	1	0	0	0	0	0	1	H _{3/2} ¹
v ¹⁴	1	0	0	0	0	0	0	0	1	0	H _{2/1} ¹
v ¹⁵	1	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ¹⁶	1	0	1	0	0	0	0	0	0	1	H _{3/2} ¹
v ¹⁷	1	1	0	0	0	0	0	0	0	1	H _{3/2} ¹

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A ₁	A ₂	A ₃	A ₄	A ₅	R ₁	R ₂	R ₃	R ₄
	dim	1	1	1	1	2	1	1	1	1
T ₁	1	1	0	1	0	1	-1	-1	0	0
T ₂	1	0	1	0	1	1	0	0	-1	-1

ker $M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A ₁	A ₂	A ₃	A ₄	A ₅	R ₁	R ₂	R ₃	R ₄	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v ¹	0	0	0	0	1	0	1	0	1	H _{3/2} ¹
v ²	0	0	0	0	1	0	1	1	0	H _{3/2} ¹
v ³	0	0	0	0	1	1	0	0	1	H _{3/2} ¹
v ⁴	0	0	0	0	1	1	0	1	0	H _{3/2} ¹
v ⁵	0	0	0	1	0	0	0	0	1	H _{2/1} ¹
v ⁶	0	0	0	1	0	0	0	1	0	H _{2/1} ¹
v ⁷	0	0	1	0	0	0	1	0	0	H _{2/1} ¹
v ⁸	0	0	1	0	0	1	0	0	0	H _{2/1} ¹

v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 100

Space group $P4bm$, No. 100, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	0	0	0	0	-1	0	0	0	0
Λ_2	1	0	1	0	0	0	0	-1	0	0	0
Λ_3	1	0	0	1	0	0	0	0	-1	0	0
Λ_4	1	0	0	0	1	0	0	0	0	-1	0
Λ_5	2	0	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	0	1	1	0	-2	-1
Σ_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
V_1	1	0	1	0	0	-1	0
V_2	1	0	1	0	0	-1	0
V_3	1	1	0	0	-1	0	0
V_4	1	1	0	0	-1	0	0
V_5	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	1	0	0	1	0	0	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1M_2	M_3M_4	M_5	X_1
	dim	2	2	2	2
Y_1Y_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1
	dim	1	1	1	1	2	2
U_1	1	1	0	1	0	1	-1
U_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
S_1	1	1	0	0	1	1	0	-2	-1
S_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3A_4	A_5	R_1
	dim	2	2	2	2
T_1T_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_4	A_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_1	X_1
	dim	2	2
W_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 101

Space group $P4_2cm$, No. 101, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	0	0	-1
Δ_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	0	0	-1	0	0
Λ_2	1	0	1	0	0	0	0	-1	0
Λ_3	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	-1	0	0
Λ_5	2	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
V_1	1	1	0	0	0	0	-1	0	0
V_2	1	0	1	0	0	0	0	-1	0
V_3	1	0	0	1	0	0	0	-1	0
V_4	1	0	0	0	1	0	-1	0	0
V_5	2	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	-1	0	0
Y_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1Z_2	Z_3Z_4	Z_5	R_1R_2	R_3R_4
	dim	2	2	2	2	2
U_1U_2	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_1Z_2	Z_3Z_4	Z_5	R_1R_2	R_3R_4	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
S_1	1	2	0	1	-2	0	-1
S_2	1	0	2	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	1	0	0	1	0	$H_{2/2}^1$
v^4	1	0	0	1	0	0	$H_{2/2}^1$
v^5	1	1	0	0	0	2	$H_{4/4}^2$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1A_2	A_3A_4	A_5	R_1R_2	R_3R_4
	dim	2	2	2	2	2
T_1T_2	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1A_2	A_3A_4	A_5	R_1R_2	R_3R_4	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1R_2	R_3R_4	X_1	X_2	X_3	X_4
	dim	2	2	1	1	1	1
W_1	1	1	0	-1	0	0	0
W_2	1	0	1	0	-1	0	0
W_3	1	1	0	0	0	-1	0
W_4	1	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1R_2	R_3R_4	X_1	X_2	X_3	X_4	
	2	2	1	1	1	1	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

SG 102

Space group $P4_2nm$, No. 102, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	0	0	-1	0	0
Λ_2	1	0	1	0	0	0	0	-1	0
Λ_3	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	1	0	-1	0	0
Λ_5	2	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	0	1	1	0	-2	-1
Σ_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1M_2	M_3M_4	M_5	A_1	A_2	A_3	A_4	A_5
	dim	2	2	2	1	1	1	1	2
V_1	1	1	0	0	-1	0	0	0	0
V_2	1	1	0	0	0	0	-1	0	0
V_3	1	0	1	0	0	0	0	-1	0
V_4	1	0	1	0	0	-1	0	0	0
V_5	2	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	A_1	A_2	A_3	A_4	A_5	
	2	2	2	1	1	1	1	2	Bipartite graph
v^1	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_1M_2	M_3M_4	M_5	X_1
dim	2	2	2	2
Y_1Y_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5	R_1
dim	2	2	2	2
U_1U_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1Z_2	Z_3Z_4	Z_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5	A_1	A_2	A_3	A_4	A_5
dim	2	2	2	1	1	1	1	2
S_1	1	2	0	1	0	-1	0	-1
S_2	1	0	2	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	$Z_1 Z_2$	$Z_3 Z_4$	Z_5	A_1	A_2	A_3	A_4	A_5	
	2	2	2	1	1	1	1	2	Bipartite graph
v^1	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	1	0	2	0	0	0	0	$H_{3/2}^1$
v^9	1	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	2	$H_{4/4}^2$
v^{13}	1	1	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	1	0	0	1	1	0	1	$H_{5/4}^3$
v^{15}	1	1	0	1	0	0	1	1	$H_{5/4}^3$
v^{16}	1	1	0	1	1	0	0	1	$H_{5/4}^3$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	R_1
	dim	1	1	1	1	2	2
T_1	1	0	1	1	0	1	-1
T_2	1	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1	A_2	A_3	A_4	A_5	R_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	X_1
	dim	2	2
W_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 103

Space group $P4cc$, No. 103, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	0	0	-1
Δ_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5Z_5
	dim	1	1	1	1	2	2	2	4
Λ_1	1	1	0	0	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	0	-1	0
Λ_3	1	0	0	1	0	0	-1	0	0
Λ_4	1	0	0	0	1	0	-1	0	0
Λ_5	2	0	0	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5Z_5	
	1	1	1	1	2	2	2	4	Bipartite graph
v^1	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^3	1	1	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$

v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5A_5
	dim	1	1	1	1	2	2	2	4
V_1	1	1	0	0	0	0	0	-1	0
V_2	1	0	1	0	0	0	0	-1	0
V_3	1	0	0	1	0	0	-1	0	0
V_4	1	0	0	0	1	0	-1	0	0
V_5	2	0	0	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5A_5	Bipartite graph
	1	1	1	1	2	2	2	4	
v^1	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^3	1	1	0	0	0	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	-1	0	0
Y_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5Z_5	R_1R_2	R_3R_4
dim	2	2	4	2	2
U_1U_2	2	1	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1Z_2	Z_3Z_4	Z_5Z_5	R_1R_2	R_3R_4	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5Z_5	A_1A_2	A_3A_4	A_5A_5	
dim	2	2	4	2	2	4	
S_1S_2	2	1	1	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Z_1Z_2	Z_3Z_4	Z_5Z_5	A_1A_2	A_3A_4	A_5A_5	Bipartite graph
	2	2	4	2	2	4	
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	2	0	$H_{3/2}^1$
v^3	0	0	1	1	1	0	$H_{3/2}^1$
v^4	0	0	1	2	0	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	1	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3A_4	A_5A_5	R_1R_2	R_3R_4	
dim	2	2	4	2	2	
T_1T_2	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1A_2	A_3A_4	A_5A_5	R_1R_2	R_3R_4	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1R_2	R_3R_4	X_1	X_2	X_3	X_4
	dim	2	2	1	1	1	1
W_1	1	1	0	-1	0	0	0
W_2	1	0	1	0	-1	0	0
W_3	1	1	0	0	0	-1	0
W_4	1	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1R_2	R_3R_4	X_1	X_2	X_3	X_4	
	2	2	1	1	1	1	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

SG 104

Space group $P4nc$, No. 104, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5Z_5
	dim	1	1	1	1	2	2	2	4
Λ_1	1	1	0	0	0	0	0	-1	0
Λ_2	1	0	1	0	0	0	0	-1	0
Λ_3	1	0	0	1	0	0	-1	0	0
Λ_4	1	0	0	0	1	0	-1	0	0
Λ_5	2	0	0	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5Z_5	
	1	1	1	1	2	2	2	4	Bipartite graph
v^1	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^3	1	1	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	0	1	1	0	-2	-1
Σ_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$

v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
V_1	1	1	0	0	0	-1	0
V_2	1	1	0	0	-1	0	0
V_3	1	0	1	0	0	-1	0
V_4	1	0	1	0	-1	0	0
V_5	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	1	1	0	1	1	0	$H_{4/4}^2$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1M_2	M_3M_4	M_5	X_1
	dim	2	2	2	2
Y_1Y_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5Z_5	R_1
dim	2	2	4	2
U_1U_2	2	1	1	2
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1Z_2	Z_3Z_4	Z_5Z_5	R_1	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/2}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5Z_5	A_1A_2	A_3A_4	A_5
dim	2	2	4	2	2	2
S_1S_2	2	1	1	2	-1	-1
					-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1Z_2	Z_3Z_4	Z_5Z_5	A_1A_2	A_3A_4	A_5	
	2	2	4	2	2	2	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1A_2	A_3A_4	A_5	R_1
	dim	2	2	2	2
T_1	1	2	0	1	-1
T_2	1	0	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_2	A_3A_4	A_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	1	1	0	2	$H_{4/4}^2$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	X_1
	dim	2	2
W_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 105

Space group $P4_2mc$, No. 105, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	0	0	-1
Δ_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	0	0	-1	0	0
Λ_2	1	0	1	0	0	0	0	-1	0
Λ_3	1	0	0	1	0	0	-1	0	0
Λ_4	1	0	0	0	1	0	0	-1	0
Λ_5	2	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	1	0	1	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
V_1	1	1	0	0	0	0	-1	0	0
V_2	1	0	1	0	0	0	0	-1	0
V_3	1	0	0	1	0	0	-1	0	0
V_4	1	0	0	0	1	0	0	-1	0
V_5	2	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	1	0	1	0	0	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	-1	0	0
Y_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1Z_2	Z_3Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	2	2	2	1	1	1	1
U_1	1	2	0	1	-1	0	0	-1
U_2	1	0	2	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_1Z_2	Z_3Z_4	Z_5	R_1	R_2	R_3	R_4	Bipartite graph
	2	2	2	1	1	1	1	
v^1	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	1	0	0	0	2	0	$H_{3/2}^1$
v^6	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	1	0	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	0	2	$H_{3/2}^1$
v^9	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	2	0	0	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
S_1S_2	2	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1A_2	A_3A_4	A_5	R_1	R_2	R_3	R_4
	dim	2	2	2	1	1	1	1
T_1	1	2	0	1	-1	-1	0	0
T_2	1	0	2	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1A_2	A_3A_4	A_5	R_1	R_2	R_3	R_4	Bipartite graph
	2	2	2	1	1	1	1	
v^1	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	2	$H_{3/2}^1$
v^6	0	1	0	0	0	1	1	$H_{3/2}^1$
v^7	0	1	0	0	0	2	0	$H_{3/2}^1$
v^8	1	0	0	0	2	0	0	$H_{3/2}^1$
v^9	1	0	0	1	1	0	0	$H_{3/2}^1$
v^{10}	1	0	0	2	0	0	0	$H_{3/2}^1$

SG 106

Space group $P4_2bc$, No. 106, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	Bipartite graph
1	1	1	1	2	2	

v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
dim		1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	0	0	-1	0	0
Λ_2	1	0	1	0	0	0	0	-1	0
Λ_3	1	0	0	1	0	0	-1	0	0
Λ_4	1	0	0	0	1	0	0	-1	0
Λ_5	2	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^3	1	0	1	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
dim		1	1	1	1	2	2	2	2
Σ_1	1	1	0	0	1	1	0	-2	-1
Σ_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	Bipartite graph
	1	1	1	1	2	2	2	2	

v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5A_5
	dim	2	2	2	2	2	4
V_1	1	1	0	0	0	-1	0
V_2	1	1	0	0	-1	0	0
V_3	1	0	1	0	-1	0	0
V_4	1	0	1	0	0	-1	0
V_5	2	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5A_5	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	1	1	0	1	1	0	$H_{4/4}^2$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_1M_2	M_3M_4	M_5	X_1
dim	2	2	2	2
Y_1Y_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5	R_1
dim	2	2	2	2
U_1	1	2	0	1
U_2	1	0	2	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	Z_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	1	1	0	2	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5A_5
dim	2	2	2	2	2	4
S_1S_2	2	1	1	1	-1	-1
						-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5A_5	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	1	1	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	0	1	$H_{3/2}^1$
v^{11}	1	1	0	0	0	1	$H_{3/2}^1$
v^{12}	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3A_4	A_5A_5	R_1
dim	2	2	4	2
T_1T_2	2	1	1	2
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_4	A_5A_5	R_1	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/2}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	X_1
dim	2	2
W_1	2	1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 107

Space group $I4mm$, No. 107, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	1	0	1	-1	0	-1	0	-1
Σ_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	0	1	1	-1	-1	0	0
Δ_2	1	0	1	1	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_1	N_2	P_1	P_2P_4	P_3
	dim	1	1	1	2	1
Q_1	1	1	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	N_1	N_2	P_1	P_2P_4	P_3	Bipartite graph
	1	1	1	2	1	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	1	0	0	$H_{2/1}^1$
v^3	0	2	0	1	0	$H_{3/2}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$

v^5	1	0	1	0	0	$H_{2/1}^1$
v^6	1	1	0	1	0	$H_{3/2}^1$
v^7	2	0	0	1	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W : $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1	X_2	X_3	X_4	P_1	P_2P_4	P_3
	dim	1	1	1	1	1	2	1
W_1	1	1	0	0	0	-1	0	0
W_2	1	0	1	0	0	0	-1	0
W_3	1	0	0	1	0	0	0	-1
W_4	1	0	0	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	X_3	X_4	P_1	P_2P_4	P_3	
	1	1	1	1	1	2	1	Bipartite graph
v^1	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	1	0	$H_{3/2}^1$
v^3	1	0	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y : $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	1	1	1	1	2
Y_1	1	1	0	0	1	-1	0	0	-1	-1
Y_2	1	0	1	1	0	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$

v^6	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 108

Space group $I4cm$, No. 108, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	1	0	1	-1	0	-1	0	-1
Σ_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	0	1	1	-1	-1	0	0
Δ_2	1	0	1	1	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_1N_2	P_1P_2	P_3P_3	P_4P_4
	dim	2	2	2	2
Q_1Q_1	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	N_1N_2	P_1P_2	P_3P_3	P_4P_4	Bipartite graph
	2	2	2	2	
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1	X_2	X_3	X_4	P_1P_2	P_3P_3	P_4P_4
	dim	1	1	1	1	2	2	2
W_1	1	1	0	0	0	-1	0	0
W_2	1	0	1	0	0	0	0	-2
W_3	1	0	0	1	0	-1	0	0
W_4	1	0	0	0	1	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	X_3	X_4	P_1P_2	P_3P_3	P_4P_4	
	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	2	0	1	0	$H_{3/2}^1$
v^2	0	2	0	0	0	0	1	$H_{3/2}^1$
v^3	1	0	1	0	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	1	1	1	1	2
Y_1	1	1	0	0	1	-1	0	0	-1	-1
Y_2	1	0	1	1	0	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 109

Space group $I4_1md$, No. 109, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	1	0	1	-2	0	-1
Σ_2	1	0	1	0	1	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^{10}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	1	1	1	1	0	$H_{5/4}^3$
v^{13}	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	0	1	1	-1
Δ_2	1	0	1	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_1	N_2	P_1
	dim	1	1	2
Q_1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	N_1	N_2	P_1	
	1	1	2	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_1	P_1
	dim	2	2
W_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	P_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	Z_1Z_2	Z_3Z_4	Z_5
dim	2	2	2	2
Y_1Y_2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	Z_1Z_2	Z_3Z_4	Z_5	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

SG 110

Space group $I4_1cd$, No. 110, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	
dim	1	1	1	1	2	2	2	2	
Σ_1	1	1	0	1	0	1	-2	0	-1
Σ_2	1	0	1	0	1	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	
1	1	1	1	2	2	2	2	Bipartite graph

v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^{10}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	1	1	1	1	0	$H_{5/4}^3$
v^{13}	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	0	1	1	-1
Δ_2	1	0	1	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	N_1N_2	P_1P_1
dim	2	4
Q_1Q_1	2	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	N_1N_2	P_1P_1	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	X_1	P_1P_1
dim	2	4
W_1	2	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	P_1P_1	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	Z_1Z_2	Z_3Z_4	Z_5
dim	2	2	2	2
Y_1Y_2	2	1	-1 -1 -1	

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	Z_1Z_2	Z_3Z_4	Z_5	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

SG 111

Space group $P\bar{4}2m$, No. 111, without SOC Δ High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$ **A** matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	-1	0	0
Δ_2	1	0	1	0	1	1	0	0	-1	-1

 $\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

 Λ High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$ **A** matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	0	0	1	0	-1	0	0	-1	0
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	0	0	0	0	-1
Λ_3	1	0	1	1	0	0	0	-1	-1	0	0

 $\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
V_1	1	1	0	0	1	0	-1	0	0	-1	0
V_2V_4	2	0	0	0	0	1	0	0	0	0	-1
V_3	1	0	1	1	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	0	0	-1
Y_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
1	1	1	1	2	1	1	1	1	

v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U : $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
U_1	1	1	0	1	0	1	-1	-1	0	0
U_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z ₁	Z ₂	Z ₃	Z ₄	Z ₅	A ₁	A ₂	A ₃	A ₄	A ₅
	dim	1	1	1	1	2	1	1	1	1	2
S ₁	1	1	0	0	1	1	-1	0	0	-1	-1
S ₂	1	0	1	1	0	1	0	-1	-1	0	-1

ker $M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Z ₁	Z ₂	Z ₃	Z ₄	Z ₅	A ₁	A ₂	A ₃	A ₄	A ₅	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v ¹	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v ²	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v ³	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v ⁴	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v ⁵	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v ⁶	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v ⁷	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v ⁸	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v ⁹	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v ¹⁰	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v ¹¹	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v ¹²	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v ¹³	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v ¹⁴	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v ¹⁵	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v ¹⁶	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v ¹⁷	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A ₁	A ₂	A ₃	A ₄	A ₅	R ₁	R ₂	R ₃	R ₄
	dim	1	1	1	1	2	1	1	1	1
T ₁	1	1	0	1	0	1	-1	0	0	-1
T ₂	1	0	1	0	1	1	0	-1	-1	0

ker $M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 112

Space group $P\bar{4}2c$, No. 112, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	-1	0	0
Δ_2	1	0	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$

v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
dim	1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	1	0	0	-1
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	-1	0
Λ_3	1	0	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	1	1	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$

v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
V_1	1	1	0	0	1	0	0	0	-1
V_2V_4	2	0	0	0	0	1	-1	-1	0
V_3	1	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	1	1	0	0	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	1	0	1	-1	0	0	-1
Y_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1Z_2	Z_3Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	2	2	2	1	1	1	1
U_1	1	0	2	1	-1	-1	0	0
U_2	1	2	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_1Z_2	Z_3Z_4	Z_5	R_1	R_2	R_3	R_4	Bipartite graph
	2	2	2	1	1	1	1	
v^1	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	2	0	0	$H_{3/2}^1$
v^6	0	1	0	1	1	0	0	$H_{3/2}^1$
v^7	0	1	0	2	0	0	0	$H_{3/2}^1$

v^8	1	0	0	0	0	0	2	$H_{3/2}^1$
v^9	1	0	0	0	0	1	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	2	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5	
dim	2	2	2	2	2	2	
S_1S_2	2	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3A_4	A_5	R_1	R_2	R_3	R_4
dim	2	2	2	1	1	1	1
T_1	1	2	0	1	-1	0	-1
T_2	1	0	2	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 10.

A_1A_2	A_3A_4	A_5	R_1	R_2	R_3	R_4	Bipartite graph
2	2	2	1	1	1	1	

v^1	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	1	0	0	0	2	0	$H_{3/2}^1$
v^6	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	1	0	0	2	0	0	$H_{3/2}^1$
v^8	1	0	0	0	0	0	2	$H_{3/2}^1$
v^9	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	2	0	0	0	$H_{3/2}^1$

SG 113

Space group $P\bar{4}2_1m$, No. 113, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	0	0	1	0	-1	0	0	-1	0
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	0	0	0	0	-1
Λ_3	1	0	1	1	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	0	1	1	0	-2	-1
Σ_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$

v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
V_1V_3	2	0	0	1	0	0	-1
V_2V_2	2	0	1	0	0	-1	0
V_4V_4	2	1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1M_2	M_3M_4	M_5	X_1
	dim	2	2	2	2
Y_1Y_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1
	dim	1	1	1	1	2	2
U_1	1	1	0	1	0	1	-1
U_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	Bipartite graph
	1	1	1	1	2	2	
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
S_1	1	1	0	0	1	1	0	-2	-1
S_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$

v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3A_4	A_5	R_1
dim	2	2	2	2
T_1T_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_4	A_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	R_1	X_1
dim	2	2
W_1W_2	2	1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 114

Space group $P\bar{4}2_1c$, No. 114, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	1	0	1	-1
Δ_2	1	0	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	-1
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	-1	-1	0
Λ_3	1	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	1	1	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
dim	1	1	1	1	2	2	2	2	
Σ_1	1	1	0	0	1	1	0	-2	-1
Σ_2	1	0	1	1	0	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	1	1	0	$H_{5/4}^3$
v^9	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	1	0	1	1	1	0	$H_{5/4}^3$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5A_5	
dim	2	2	2	2	2	4	
V_1V_3	2	0	0	1	-1	-1	0
V_2V_2	2	0	1	0	0	0	-1
V_4V_4	2	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5A_5	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_1M_2	M_3M_4	M_5	X_1
dim	2	2	2	2
Y_1Y_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5	R_1
dim	2	2	2	2
U_1	1	0	2	1
U_2	1	2	0	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_4	Z_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	1	1	0	2	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5A_5	
dim	2	2	2	2	2	4	
S_1S_2	2	1	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5A_5	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	1	1	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	0	1	$H_{3/2}^1$
v^{11}	1	1	0	0	0	1	$H_{3/2}^1$
v^{12}	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3A_4	A_5A_5	R_1	
dim	2	2	4	2	
T_1T_2	2	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_4	A_5A_5	R_1	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/2}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	X_1
dim	2	2
W_1W_2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 115

Space group $P\bar{4}m2$, No. 115, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	0	1	1	-1	0	-1
Δ_2	1	0	1	1	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	0	0	1	0	-1	0	0	-1	0
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	0	0	0	0	-1
Λ_3	1	0	1	1	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	1	0	1	-1	0	-1	0	-1
Σ_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$

v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
V_1	1	1	0	0	1	0	-1	0	0	-1	0
V_2V_4	2	0	0	0	0	1	0	0	0	0	-1
V_3	1	0	1	1	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	M_1	M_2	M_3	M_4	M_5	A_1	A_2	A_3	A_4	A_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	0	1	1	-1	-1	0	0
Y_2	1	0	1	1	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
U_1	1	1	0	0	1	1	-1	0	0	-1
U_2	1	0	1	1	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$

v^8	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1	A_2	A_3	A_4	A_5
	dim	1	1	1	1	2	1	1	1	1	2
S_1	1	1	0	1	0	1	-1	0	-1	0	-1
S_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1	A_2	A_3	A_4	A_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	R_1	R_2	R_3	R_4
	dim	1	1	1	1	2	1	1	1	1
T_1	1	1	0	0	1	1	-1	-1	0	0
T_2	1	0	1	1	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	R_1	R_2	R_3	R_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 116

Space group $P\bar{4}c2$, No. 116, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	0	1	1	-1	0	0	-1
Δ_2	1	0	1	1	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
1	1	1	1	2	1	1	1	1	

v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	-1
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	-1	-1	0
Λ_3	1	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	1	1	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	2

Σ_1	1	1	0	1	0	1	-1	0	-1	0	-1
Σ_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
V_1	1	1	0	0	1	0	0	0	-1
V_2V_4	2	0	0	0	0	1	-1	-1	0
V_3	1	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1	M_2	M_3	M_4	M_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	0	1	$H_{3/2}^1$

v^4	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	1	1	0	0	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Y_1	1	1	0	0	1	1	-1	-1	0	0
Y_2	1	0	1	1	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	R_1	R_2	R_3	R_4
	dim	2	2	2			2	2		
U_1U_2	2	1	1	1			-1	-1		

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_1Z_2	Z_3Z_4	Z_5	R_1R_2	R_3R_4	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
S_1	1	2	0	1	-2	0	-1
S_2	1	0	2	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Z_1Z_2	Z_3Z_4	Z_5	A_1A_2	A_3A_4	A_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	1	0	0	1	0	$H_{2/2}^1$
v^4	1	0	0	1	0	0	$H_{2/2}^1$
v^5	1	1	0	0	0	2	$H_{4/4}^2$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3A_4	A_5	R_1R_2	R_3R_4
	dim	2	2	2	2	2
T_1T_2	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1A_2	A_3A_4	A_5	R_1R_2	R_3R_4	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1R_2	R_3R_4	X_1	X_2	X_3	X_4
	dim	2	2	1	1	1	1
W_1	1	1	0	-1	0	0	0
W_2	1	0	1	0	-1	0	0
W_3	1	1	0	0	0	-1	0
W_4	1	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1R_2	R_3R_4	X_1	X_2	X_3	X_4	
	2	2	1	1	1	1	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

SG 117

Space group $P4b2$, No. 117, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	0	1	1	-1
Δ_2	1	0	1	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Λ_1	1	1	0	0	1	0	-1	0	0	-1	0
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	0	0	0	0	-1
Λ_3	1	0	1	1	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	1	0	1	0	-2	-1
Σ_2	1	0	1	0	1	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	1	1	1	1	0	$H_{5/4}^3$
v^{13}	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5
	dim	2	2	2	2	2	2
V_1	1	0	0	1	0	0	-1
V_2	1	0	0	1	0	0	-1
V_3V_4	2	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1M_2	M_3M_4	M_5	A_1A_2	A_3A_4	A_5	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_1M_2	M_3M_4	M_5	X_1	
dim	2	2	2	2	
Y_1Y_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	
dim	1	1	1	1	2	2	
U_1	1	1	0	0	1	1	-1
U_2	1	0	1	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Z_1	Z_2	Z_3	Z_4	Z_5	R_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5
	dim	1	1	1	1	2	2	2	2
S_1	1	1	0	1	0	1	0	-2	-1
S_2	1	0	1	0	1	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1	Z_2	Z_3	Z_4	Z_5	A_1A_2	A_3A_4	A_5	
	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	1	1	1	1	0	$H_{5/4}^3$
v^{13}	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1A_2	A_3A_4	A_5	R_1
	dim	2	2	2	2
T_1T_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_4	A_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	X_1
dim		2	2
W_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

R_1		X_1	
2		2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 118

Space group $P\bar{4}n2$, No. 118, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
dim		1	1	1	1	2	2
Δ_1	1	1	0	0	1	1	-1
Δ_2	1	0	1	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

Γ_1		Γ_2	Γ_3	Γ_4	Γ_5	X_1	
1		1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	-1
$\Lambda_2\Lambda_4$	2	0	0	0	0	1	-1	-1	0
Λ_3	1	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	1	1	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	1	0	1	0	-2	-1
Σ_2	1	0	1	0	1	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	1	1	1	1	0	$H_{5/4}^3$

v^{13}	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1M_2	M_3M_4	M_5	A_1	A_2	A_3	A_4	A_5
	dim	2	2	2	1	1	1	1	2
V_1	1	0	0	1	0	-1	-1	0	0
V_2	1	0	0	1	-1	0	0	-1	0
V_3V_4	2	1	1	0	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1M_2	M_3M_4	M_5	A_1	A_2	A_3	A_4	A_5	Bipartite graph
	2	2	2	1	1	1	1	2	
v^1	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	0	1	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1M_2	M_3M_4	M_5	X_1
	dim	2	2	2	2
Y_1Y_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1M_2	M_3M_4	M_5	X_1	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Z_1 Z_2$	$Z_3 Z_4$	Z_5	R_1
dim	2	2	2	2
$U_1 U_2$	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$Z_1 Z_2$	$Z_3 Z_4$	Z_5	R_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Z_1 Z_2$	$Z_3 Z_4$	Z_5	A_1	A_2	A_3	A_4	A_5
dim	2	2	2	1	1	1	1	2
S_1	1	2	0	1	0	-1	0	-1
S_2	1	0	2	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	$Z_1 Z_2$	$Z_3 Z_4$	Z_5	A_1	A_2	A_3	A_4	A_5	
	2	2	2	1	1	1	1	2	Bipartite graph
v^1	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	1	0	2	0	0	0	0	$H_{3/2}^1$
v^9	1	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	2	$H_{4/4}^2$
v^{13}	1	1	0	0	0	1	1	1	$H_{5/4}^3$

v^{14}	1	1	0	0	1	1	0	1	$H_{5/4}^3$
v^{15}	1	1	0	1	0	0	1	1	$H_{5/4}^3$
v^{16}	1	1	0	1	1	0	0	1	$H_{5/4}^3$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	R_1
	dim	1	1	1	1	2	2
T_1	1	0	1	1	0	1	-1
T_2	1	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1	A_2	A_3	A_4	A_5	R_1	Bipartite graph
	1	1	1	1	2	2	
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	X_1
	dim	2	2
W_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1	X_1	Bipartite graph
	2	2	
v^1	1	1	$H_{2/1}^1$

SG 119

Space group $I\bar{4}m2$, No. 119, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	1	1	1	1
Δ_1	1	1	0	1	0	1	-1	0	0	-1
Δ_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	1	1	1	1	2
Y_1	1	1	1	0	0	-1	0	-1	0	-1
Y_2	1	0	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	1	1	1	1	2	
v^1	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^7	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

SG 120

Space group $I\bar{4}c2$, No. 120, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	0	1	1	-1	0	0	-1	-1
Σ_2	1	0	1	1	0	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
dim	1	1	1	1	2	1	1	1	1

Δ_1	1	1	0	1	0	1	-1	0	0	-1
Δ_2	1	0	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	N_1N_2	P_1P_3	P_2P_4
dim	2	2	2
Q_1Q_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	N_1N_2	P_1P_3	P_2P_4	Bipartite graph
	2	2	2	
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_1	X_2	X_3	X_4	P_1P_3	P_2P_4
dim		1	1	1	1	2	2
W_1	1	1	0	1	0	-2	0
W_2	1	0	1	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	X_1	X_2	X_3	X_4	P_1P_3	P_2P_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	2	0	1	0	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	0	2	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5
dim		1	1	1	1	1	1	1	1	2
Y_1	1	1	1	0	0	-1	0	-1	0	-1
Y_2	1	0	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^7	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

SG 121

Space group $I\bar{4}2m$, No. 121, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5
	dim	1	1	1	1	2	1	1	1	1	2
Σ_1	1	1	0	1	0	1	-1	0	-1	0	-1
Σ_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1	Z_2	Z_3	Z_4	Z_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4
dim	1	1	1	1	2	1	1	1	1

Δ_1	1	1	0	0	1	1	-1	-1	0	0
Δ_2	1	0	1	1	0	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	1	1	1	1	
v^1	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_1	N_2	P_1	P_2	P_3	P_4	P_5
	dim	1	1	1	1	1	1	2
Q_1	1	1	0	-1	0	-1	0	-1
Q_2	1	0	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	N_1	N_2	P_1	P_2	P_3	P_4	P_5	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	0	$H_{2/1}^1$
v^3	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	1	0	1	0	0	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	0	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1	X_2	X_3	X_4	P_1	P_2	P_3	P_4	P_5
dim		1	1	1	1	1	1	1	1	2
W_1	1	1	0	0	0	-1	0	0	-1	0
W_2	1	0	1	0	0	0	0	0	0	-1
W_3	1	0	0	1	0	0	-1	-1	0	0
W_4	1	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	X_1	X_2	X_3	X_4	P_1	P_2	P_3	P_4	P_5	Bipartite graph
	1	1	1	1	1	1	1	1	2	
v^1	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^4	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5
dim		1	1	1	1	1	1	1	1	2
Y_1	1	1	0	0	1	-1	0	0	-1	-1
Y_2	1	0	1	1	0	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	Z_1	Z_2	Z_3	Z_4	Z_5	Bipartite graph
	1	1	1	1	1	1	1	1	2	
v^1	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$

v^9	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 122

Space group $I\bar{4}2d$, No. 122, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5
	dim	1	1	1	1	2	2	2	2
Σ_1	1	1	0	1	0	1	0	-2	-1
Σ_2	1	0	1	0	1	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Z_1Z_2	Z_3Z_4	Z_5	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	2	1	1	0	$H_{4/4}^2$
v^3	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^{10}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	1	1	1	1	0	$H_{5/4}^3$
v^{13}	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{14}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^{16}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1
	dim	1	1	1	1	2	2
Δ_1	1	1	0	0	1	1	-1
Δ_2	1	0	1	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_1	N_2	P_1	P_2
	dim	1	1	2	2
Q_1	1	1	0	-1	-1
Q_2	1	0	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	N_1	N_2	P_1	P_2	
	1	1	2	2	Bipartite graph
v^1	1	1	0	1	$H_{3/2}^1$
v^2	1	1	1	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W : $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_1	P_1	P_2
	dim	2	2	2
W_1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	P_1	P_2	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	Z_1Z_2	Z_3Z_4	Z_5
dim	2	2	2	2
Y_1Y_2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	Z_1Z_2	Z_3Z_4	Z_5	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

SG 123

Space group $P4/mmm$, No. 123, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	
dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	-1	0	0	0	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	0	0	0	-1	-1	0	0
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	-1	0	0	0	0	0
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph	
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1		
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	
dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
Λ_1	1	1	0	0	1	0	0	0	0	0	-1	0	0	-1	0	0	0	0	0	0	
Λ_2	1	0	1	1	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0	0	
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	-1	0	0	-1	0	0	
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	-1	-1	0	0	0	
Λ_5	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph	
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2		
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-		
dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2		
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	0	0	0	-1	0	0	-1	
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0	-1	0	0	0	0	-1	
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	0	0	0	0	0	-1	-1	0	
Σ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1	0	-1	0	0	-1	0	

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	Bipartite graph	
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2		
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$

v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{24}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{26}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{30}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{33}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{34}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2
V_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0	0	0	0	0
V_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0	0
V_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0
V_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	-1	-1	0	0	0	0

v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U : $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
U_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	-1	0	0	0	0
U_2	1	0	0	0	1	0	0	0	1	1	0	0	0	0	0	0	-1	-1	0
U_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	-1	0	0	0	0	0
U_4	1	0	0	1	0	0	0	1	0	0	1	0	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	
1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	Bipartite graph

v^1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S : $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2
S_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	0	0	0	-1	0	0	-1
S_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0	-1	0	0	0	0	-1
S_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	0	0	0	0	0	-1	-1	0
S_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 34.

Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	
1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	Bipartite graph

T_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	0	0	0	0	-1
T_2	1	0	0	1	0	0	0	0	1	0	0	1	0	0	-1	-1	0	0	0
T_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	0	-1	0
T_4	1	0	0	0	1	0	0	0	1	1	0	0	0	-1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

SG 124

Space group $P4/mcc$, No. 124, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	-1	0	0	0	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	0	0	0	0	-1	-1	0
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	-1	0	0	0	0	0
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix



corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	4
Λ_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0
Λ_2	1	0	1	1	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4	
	1	1	1	1	1	1	1	1	2	2	2	2	4	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	0	0	0	-1	0	0	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0	-1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	0	0	0	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$

v^4	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^7	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{18}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{24}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{26}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{28}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{30}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{33}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{34}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	A_1	A_2	A_3A_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	4
V_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0
V_2	1	0	1	1	0	0	0	0	0	0	0	-1	0	0
V_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0
V_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0

$$\overline{V_5 \quad 2 \quad | \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 1 \quad | \quad 0 \quad 0 \quad -2}$$

ker $M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	A_1	A_2	A_3A_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	4	
v^1	0	0	0	0	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
Y_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	0	0	0	0	-1
Y_2	1	0	0	1	0	0	0	1	0	0	1	0	0	0	-1	-1	0	0	0
Y_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	0	-1	0
Y_4	1	0	0	0	1	0	0	0	1	1	0	0	0	-1	0	0	-1	0	0

ker $M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$

v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	Z_3Z_4	R_1	R_2
dim	2	2	4	2	2
U_1	2	1	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1	Z_2	Z_3Z_4	R_1	R_2	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3Z_4	A_1	A_2	A_3A_4
	dim	2	2	4	2	2	4
S_1	2	1	1	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Z_1	Z_2	Z_3Z_4	A_1	A_2	A_3A_4	Bipartite graph
	2	2	4	2	2	4	
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	2	0	$H_{3/2}^1$
v^3	0	0	1	1	1	0	$H_{3/2}^1$
v^4	0	0	1	2	0	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	1	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3A_4	R_1	R_2
	dim	2	2	4	2	2
T_1	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3A_4	R_1	R_2	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	R_2	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
dim	2	2	1	1	1	1	1	1	1	1
W_1	1	1	0	-1	0	0	0	0	-1	0
W_2	1	0	1	0	0	-1	0	0	0	-1
W_3	1	1	0	0	-1	0	0	-1	0	0
W_4	1	0	1	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	R_1	R_2	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	2	2	1	1	1	1	1	1	1	1	
v^1	0	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^3	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^5	1	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^6	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^8	1	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$

SG 125

Space group $P4/nbm$, No. 125, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	-1	0
Δ_4	1	0	0	1	0	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^8	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{12}	0	1	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{15}	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
Λ_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0	0	0	0	0	0
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$

v^6	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	0	0	-1	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	-1	-1	0	0
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	0	0	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$

V_1	1	0	0	0	1	0	0	0	-1
V_2	1	0	0	0	1	0	0	0	-1
V_3	1	0	1	0	0	0	-1	0	0
V_4	1	0	1	0	0	0	-1	0	0
V_5	2	1	0	1	0	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1	M_2	M_3	M_4	A_1	A_2	A_3	A_4	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	X_1	X_2
	dim	2	2	2	2	2	2
Y_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1	R_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
U_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
U_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
U_3	1	0	1	0	0	0	1	0	0	1	0	-1	0
U_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1	R_2	
	1	1	1	1	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^8	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{12}	0	1	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{15}	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	A_1	A_2	A_3	A_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
S_1	1	1	0	0	0	0	0	1	0	0	1	0	0	-1	-1
S_2	1	0	0	1	0	1	0	0	0	0	1	-1	-1	0	0
S_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
S_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	A_1	A_2	A_3	A_4	Bipartite graph
v^1	1	1	1	1	1	1	1	1	2	2	2	2	2	2	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	0	0	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	0	1	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{25}	0	0	1	0	1	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	1	0	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{30}	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{31}	0	1	0	0	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{39}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{40}	1	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$

v^{41}	1	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	R_1	R_2
dim	2	2	2	2	2	2	2
T_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_1	A_2	A_3	A_4	R_1	R_2	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	R_2	X_1	X_2
dim	2	2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

R_1	R_2	X_1	X_2	Bipartite graph
2	2	2	2	

v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 126

Space group $P4/nnc$, No. 126, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	0	-1	0
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^8	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{12}	0	1	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{15}	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	4
Λ_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0
Λ_2	1	0	1	1	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	4	
v^1	0	0	0	0	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	0	0	-1	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	-1	-1	0	0
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	0	0	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	0	1	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{25}	0	0	1	0	1	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	1	0	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{30}	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{31}	0	1	0	0	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{39}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{40}	1	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$

v^{41}	1	0	0	0	1	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	0	1	$H_{5/4}^3$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	A_1	A_2	A_3	A_4
	dim	2	2	2	2	2	2	2	2
V_1	1	0	0	0	1	0	-1	0	0
V_2	1	0	0	0	1	0	0	0	-1
V_3	1	0	1	0	0	0	-1	0	0
V_4	1	0	1	0	0	0	0	0	-1
V_5	2	1	0	1	0	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	M_3	M_4	A_1	A_2	A_3	A_4	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^3	0	1	0	1	0	1	0	1	$H_{4/4}^2$
v^4	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	X_1	X_2
	dim	2	2	2	2	2	2
Y_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3Z_4	R_1	R_2
	dim	2	2	4	2	2
U_1	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1	Z_2	Z_3Z_4	R_1	R_2	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3Z_4	A_1	A_2	A_3	A_4
	dim	2	2	4	2	2	2	2
S_1	2	1	1	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Z_1	Z_2	Z_3Z_4	A_1	A_2	A_3	A_4	
	2	2	4	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	1	$H_{3/2}^1$
v^5	0	0	1	0	1	1	0	$H_{3/2}^1$
v^6	0	0	1	0	2	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	1	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	1	2	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{17}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	1	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	A_4	R_1	R_2
	dim	2	2	2	2	2	2
T_1	1	0	1	1	0	-1	0
T_2	1	0	0	1	1	-1	0
T_3	1	1	1	0	0	0	-1
T_4	1	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	A_4	R_1	R_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/2}^1$
v^2	0	1	0	1	1	1	$H_{4/4}^2$
v^3	1	0	0	0	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	R_2	X_1	X_2
	dim	2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_1	R_2	X_1	X_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 127

Space group $P4/mbm$, No. 127, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$

v^8	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$	
v^9	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-		
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	1	2	2	
Λ_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0	0	0	0	0	0	0
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0	0	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0	0	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph	
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2		
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	
dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-2	0	0	0	0	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-2	0	0	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	0	0	-2	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	0	-2	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	2	1	0	1	0	0	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	2	0	0	1	0	1	0	0	$H_{4/4}^2$
v^5	0	0	0	0	0	0	0	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	2	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	1	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^9	0	0	0	0	0	2	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{12}	0	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	0	1	0	0	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	1	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{16}	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{19}	0	0	1	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	1	0	0	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{24}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	0	1	0	1	0	0	0	0	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{26}	0	2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{28}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

v^{29}	1	0	0	0	1	0	0	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{30}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{31}	1	0	1	0	0	0	0	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{32}	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-	
dim	2	2	2	2	2	2	2	2	2	2	2	2	
$V_1 V_2$	2	1	1	0	0	0	0	-1	-1	0	0	0	0
$V_3 V_4$	2	0	0	1	1	0	0	0	0	-1	-1	0	0
V_5	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-	Bipartite graph
	2	2	2	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	X_1	X_2	
dim	2	2	2	2	2	2	2	2	
$Y_1 Y_2$	2	1	0	1	0	0	1	-1	0
$Y_3 Y_4$	2	0	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	X_1	X_2	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1	R_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
U_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
U_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
U_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
U_4	1	0	0	1	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1	R_2	
	1	1	1	1	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	$A_1^+A_2^+$	$A_1^-A_2^-$	$A_3^+A_4^+$	$A_3^-A_4^-$	A_5^+	A_5^-
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2
S_1	1	1	0	0	0	0	0	1	0	0	1	-2	0	0	0	0	-1
S_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-2	0	0	-1
S_3	1	0	1	0	0	0	0	0	1	1	0	0	0	0	-2	-1	0
S_4	1	0	0	0	1	0	1	0	0	1	0	0	-2	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	$A_1^+A_2^+$	$A_1^-A_2^-$	$A_3^+A_4^+$	$A_3^-A_4^-$	A_5^+	A_5^-	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	2	1	0	1	0	0	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	2	0	0	1	0	1	0	0	$H_{4/4}^2$
v^5	0	0	0	0	0	0	0	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	2	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	1	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^9	0	0	0	0	0	2	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{12}	0	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	0	1	0	0	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	1	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{16}	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{19}	0	0	1	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	1	0	0	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{24}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	0	1	0	1	0	0	0	0	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{26}	0	2	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{28}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{29}	1	0	0	0	1	0	0	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{30}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{31}	1	0	1	0	0	0	0	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$

$$\overline{v^{32} \mid 2 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ \mid 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ \mid H_{3/2}^1}$$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-	R_1	R_2
	dim	2	2	2	2	2	2	2	2
$T_1 T_2$	2	1	0	1	0	0	1	-1	0
$T_3 T_4$	2	0	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-	R_1	R_2	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_1	R_2	X_1	X_2
	dim	2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_1	R_2	X_1	X_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 128

Space group $P4/mnc$, No. 128, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	4
Λ_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0
Λ_2	1	0	1	1	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	4	
v^1	0	0	0	0	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-
dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	1	0	0	1	-2	0	0	0	0	-1
Σ_2	1	0	0	1	0	1	0	0	0	1	0	0	-2	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	1	0	0	0	0	-2	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-2	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	2	1	0	1	0	0	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	2	0	0	1	0	1	0	0	$H_{4/4}^2$
v^5	0	0	0	0	0	0	0	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	2	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	1	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^9	0	0	0	0	0	2	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{12}	0	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$

v^{13}	0	0	0	1	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	0	1	0	0	0	1	1	0	0	1	0	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{16}	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	1	0	0	1	1	0	1	0	0	$H_{5/4}^3$
v^{19}	0	0	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{20}	0	0	2	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{22}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{23}	0	1	0	0	0	1	0	0	1	0	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	1	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	0	1	0	1	0	0	0	0	1	0	0	1	0	1	0	$H_{5/4}^3$
v^{26}	0	2	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{28}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{29}	1	0	0	0	1	0	0	0	0	1	1	0	1	0	0	$H_{5/4}^3$
v^{30}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{31}	1	0	1	0	0	0	0	0	0	1	1	0	1	0	0	$H_{5/4}^3$
v^{32}	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	A_1	A_2	$A_3 A_4$
dim	2	2	2	2	2	2	2	2	4
$V_1 V_2$	2	1	1	0	0	0	0	0	-1
$V_3 V_4$	2	0	0	1	1	0	0	0	-1
V_5	2	0	0	0	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	A_1	A_2	$A_3 A_4$	Bipartite graph
	2	2	2	2	2	2	2	2	4	
v^1	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y : $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	X_1	X_2
	dim	2	2	2	2	2	2	2	2
$Y_1 Y_2$	2	1	0	1	0	0	1	-1	0
$Y_3 Y_4$	2	0	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	X_1	X_2	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U : $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	$Z_3 Z_4$	R_1^+	R_1^-
	dim	2	2	4	2	2
U_1	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1	Z_2	$Z_3 Z_4$	R_1^+	R_1^-	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	Z_3Z_4	A_1	A_2	A_3A_4	
dim	2	2	4	2	2	4	
S_1	2	1	1	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Z_1	Z_2	Z_3Z_4	A_1	A_2	A_3A_4	Bipartite graph
	2	2	4	2	2	4	
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	2	0	$H_{3/2}^1$
v^3	0	0	1	1	1	0	$H_{3/2}^1$
v^4	0	0	1	2	0	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	1	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3A_4	R_1^+	R_1^-	
dim	2	2	4	2	2	
T_1	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3A_4	R_1^+	R_1^-	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1^+	R_1^-	X_1	X_2
dim	2	2	2	2
W_1	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_1^+	R_1^-	X_1	X_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 129

Space group $P4/nmm$, No. 129, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	-1	0
Δ_3	1	0	1	0	0	0	1	0	0	1	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^3	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^6	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^7	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$

v^8	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{11}	0	1	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{13}	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0	0	0	0	0
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	-1	-1	0	0	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
dim		1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	-1	0	0
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	1	1	0	0	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

v^{22}	0	0	1	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{25}	0	0	1	0	1	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	1	0	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{30}	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{31}	0	1	0	0	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{39}	1	0	0	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{40}	1	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{41}	1	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	A_1	A_2	A_3	A_4
	dim	2	2	2	2	2	2	2	2
V_1V_4	2	1	0	0	0	-1	0	0	0
V_2V_3	2	0	0	1	0	0	0	-1	0
V_5	2	0	1	0	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1	M_2	M_3	M_4	A_1	A_2	A_3	A_4	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	$H_{2/1}^1$

v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	X_1	X_2
	dim	2	2	2	2	2	2
Y_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1	R_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
U_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
U_2	1	0	0	0	1	0	0	0	1	1	0	-1	0
U_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
U_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	R_1	R_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^3	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^6	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^7	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{11}	0	1	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{13}	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S : $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	A_1	A_2	A_3	A_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
S_1	1	1	0	0	0	0	0	1	0	0	1	-1	-1	0	0
S_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	-1
S_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
S_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	A_1	A_2	A_3	A_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	1	1	0	0	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$

v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{25}	0	0	1	0	1	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	1	0	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{30}	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{31}	0	1	0	0	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{39}	1	0	0	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{40}	1	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{41}	1	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	A_4	R_1	R_2
dim	2	2	2	2	2	2
T_1	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_1	A_2	A_3	A_4	R_1	R_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	R_2	X_1	X_2
dim	2	2	2	2
W_1W_4	2	1	0	-1 0
W_2W_3	2	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1	R_2	X_1	X_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 130

Space group $P4/ncc$, No. 130, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	-1	0
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^3	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^6	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^7	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{11}	0	1	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{13}	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	4
Λ_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0
Λ_2	1	0	1	1	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0

Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	-1	0
Λ_5	2	0	0	0	0	0	0	0	0	0	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	4	
v^1	0	0	0	0	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
dim		1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	-1	0	0
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	1	1	0	0	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$

v ⁹	0	0	0	0	0	2	0	0	0	1	1	0	0	1	H _{5/4} ³
v ¹⁰	0	0	0	0	1	0	0	1	0	0	0	0	1	0	H _{3/2} ¹
v ¹¹	0	0	0	0	1	0	1	0	1	0	0	1	0	1	H _{5/4} ³
v ¹²	0	0	0	0	1	0	1	0	1	0	1	0	1	0	H _{5/4} ³
v ¹³	0	0	0	0	1	1	0	0	0	0	0	0	0	1	H _{3/2} ¹
v ¹⁴	0	0	0	0	2	0	0	0	1	0	0	0	1	1	H _{5/4} ³
v ¹⁵	0	0	0	1	0	0	0	1	0	1	0	1	0	1	H _{5/4} ³
v ¹⁶	0	0	0	1	0	0	0	1	0	1	1	0	1	0	H _{5/4} ³
v ¹⁷	0	0	0	1	0	0	1	0	0	0	1	0	0	0	H _{3/2} ¹
v ¹⁸	0	0	0	1	0	1	0	0	0	1	1	0	0	1	H _{5/4} ³
v ¹⁹	0	0	0	1	1	0	0	0	0	0	0	0	0	1	H _{3/2} ¹
v ²⁰	0	0	0	2	0	0	0	0	0	1	1	0	0	1	H _{5/4} ³
v ²¹	0	0	1	0	0	0	0	1	0	0	0	0	1	0	H _{3/2} ¹
v ²²	0	0	1	0	0	0	1	0	1	0	0	1	0	1	H _{5/4} ³
v ²³	0	0	1	0	0	0	1	0	1	0	1	0	1	0	H _{5/4} ³
v ²⁴	0	0	1	0	0	1	0	0	0	0	0	0	0	1	H _{3/2} ¹
v ²⁵	0	0	1	0	1	0	0	0	1	0	0	0	1	1	H _{5/4} ³
v ²⁶	0	0	1	1	0	0	0	0	0	0	0	0	0	1	H _{3/2} ¹
v ²⁷	0	0	2	0	0	0	0	0	1	0	0	0	1	1	H _{5/4} ³
v ²⁸	0	1	0	0	0	0	0	1	0	1	0	1	1	0	H _{5/4} ³
v ²⁹	0	1	0	0	0	0	1	0	0	0	0	1	0	0	H _{3/2} ¹
v ³⁰	0	1	0	0	0	1	0	0	0	1	0	1	0	1	H _{5/4} ³
v ³¹	0	1	0	0	0	1	0	0	0	1	1	0	1	0	H _{5/4} ³
v ³²	0	1	0	0	1	0	0	0	0	0	0	0	1	0	H _{3/2} ¹
v ³³	0	1	0	1	0	0	0	0	0	1	0	1	0	1	H _{5/4} ³
v ³⁴	0	1	0	1	0	0	0	0	0	1	1	0	1	0	H _{5/4} ³
v ³⁵	0	1	1	0	0	0	0	0	0	0	0	0	1	0	H _{3/2} ¹
v ³⁶	0	2	0	0	0	0	0	0	0	1	0	1	1	0	H _{5/4} ³
v ³⁷	1	0	0	0	0	0	0	1	0	0	0	1	0	0	H _{3/2} ¹
v ³⁸	1	0	0	0	0	0	1	0	1	0	1	1	0	0	H _{5/4} ³
v ³⁹	1	0	0	0	0	1	0	0	0	0	1	0	0	0	H _{3/2} ¹
v ⁴⁰	1	0	0	0	1	0	0	0	1	0	0	1	0	1	H _{5/4} ³
v ⁴¹	1	0	0	0	1	0	0	0	1	0	1	0	1	0	H _{5/4} ³
v ⁴²	1	0	0	1	0	0	0	0	0	0	1	0	0	0	H _{3/2} ¹
v ⁴³	1	0	1	0	0	0	0	0	1	0	0	1	0	1	H _{5/4} ³
v ⁴⁴	1	0	1	0	0	0	0	0	1	0	1	0	1	0	H _{5/4} ³
v ⁴⁵	1	1	0	0	0	0	0	0	0	0	0	1	0	0	H _{3/2} ¹
v ⁴⁶	2	0	0	0	0	0	0	0	1	0	1	1	0	0	H _{5/4} ³

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_1	M_2	M_3	M_4	A_1A_2	A_3A_4	
dim	2	2	2	2	4	4	
V_1V_4	2	1	0	0	0	-1	0
V_2V_3	2	0	0	1	0	-1	0
V_5	2	0	1	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	A_1A_2	A_3A_4	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	2	0	0	0	1	$H_{3/2}^1$
v^4	1	0	1	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_1	M_2	M_3	M_4	X_1	X_2	
dim	2	2	2	2	2	2	
Y_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3Z_4	R_1R_2
	dim	2	2	4	4
U_1	2	1	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_1	Z_2	Z_3Z_4	R_1R_2	
	2	2	4	4	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	1	0	1	$H_{3/2}^1$
v^4	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3Z_4	A_1A_2	A_3A_4
	dim	2	2	4	4	4
S_1	2	1	1	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1	Z_2	Z_3Z_4	A_1A_2	A_3A_4	
	2	2	4	4	4	Bipartite graph
v^1	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	$H_{3/2}^1$
v^6	1	1	0	1	0	$H_{3/2}^1$
v^7	2	0	0	0	1	$H_{3/2}^1$
v^8	2	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1A_2	A_3A_4	R_1R_2
	dim	4	4	4
T_1T_4	2	1	1	-1
T_2T_3	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_2	A_3A_4	R_1R_2	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_1R_2	X_1	X_2
	dim	4	2	2
W_1W_4	2	1	-1	0
W_2W_3	2	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_1R_2	X_1	X_2	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 131

Space group $P4_2/mmc$, No. 131, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	-1	0	0	0	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	0	0	0	0	-1	-1	0
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	-1	0	0	0	0	0
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph	
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1		
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	-1
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	
dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	0	0	0	-1	0	0	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0	-1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	0	0	0	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$

v ¹⁰	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	H _{3/2} ¹
v ¹¹	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	H _{2/1} ¹
v ¹²	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	H _{2/1} ¹
v ¹³	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ¹⁴	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	H _{2/1} ¹
v ¹⁵	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	H _{2/1} ¹
v ¹⁶	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	H _{2/1} ¹
v ¹⁷	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	1	0	H _{3/2} ¹
v ¹⁸	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	H _{2/1} ¹
v ¹⁹	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	H _{2/1} ¹
v ²⁰	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	H _{3/2} ¹
v ²¹	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	H _{2/1} ¹
v ²²	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	H _{2/1} ¹
v ²³	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	H _{3/2} ¹
v ²⁴	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	H _{2/1} ¹
v ²⁵	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	H _{2/1} ¹
v ²⁶	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	H _{3/2} ¹
v ²⁷	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	H _{2/1} ¹
v ²⁸	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	H _{2/1} ¹
v ²⁹	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	H _{3/2} ¹
v ³⁰	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	H _{3/2} ¹
v ³¹	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	H _{2/1} ¹
v ³²	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	H _{2/1} ¹
v ³³	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	H _{3/2} ¹
v ³⁴	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	H _{3/2} ¹

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M ₁ ⁺	M ₁ ⁻	M ₂ ⁺	M ₂ ⁻	M ₃ ⁺	M ₃ ⁻	M ₄ ⁺	M ₄ ⁻	M ₅ ⁺	M ₅ ⁻	A ₁	A ₂	A ₃	A ₄
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
V ₁	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
V ₂	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
V ₃	1	0	0	0	0	1	0	0	1	0	0	0	0	0	-1
V ₄	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	0
V ₅	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

ker M ∩ Z_{≥0}¹⁴ and the corresponding bipartite graph, number of Hilbert basis: 12.

M ₁ ⁺	M ₁ ⁻	M ₂ ⁺	M ₂ ⁻	M ₃ ⁺	M ₃ ⁻	M ₄ ⁺	M ₄ ⁻	M ₅ ⁺	M ₅ ⁻	A ₁	A ₂	A ₃	A ₄	
1	1	1	1	1	1	1	1	2	2	2	2	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0			$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0		$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0		$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0		$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	0	0	0	0	0	0	1		$H_{3/2}^1$
v^6	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1		$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	0	0	0	0	1	0	0		$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	0	1	0	0		$H_{3/2}^1$
v^9	0	1	0	0	0	0	1	0	0	0	0	0	1	0	0		$H_{3/2}^1$
v^{10}	0	1	0	0	0	1	0	0	0	0	0	0	1	0	0		$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	0	0	1		$H_{3/2}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1		$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
Y_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	0	0	0	0	-1
Y_2	1	0	0	1	0	0	0	1	0	0	1	0	0	0	-1	-1	0	0	0
Y_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	0	-1	0
Y_4	1	0	0	0	1	0	0	0	1	1	0	0	0	-1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$

v^{14}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$	
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-
dim		2	2	2	2	1	1	1	1	1	1	1	1
U_1	1	1	0	0	1	-1	0	0	-1	0	0	0	0
U_2	1	0	0	1	1	0	0	0	0	0	-1	-1	0
U_3	1	1	1	0	0	0	-1	-1	0	0	0	0	0
U_4	1	0	1	1	0	0	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1	Z_2	Z_3	Z_4	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	Bipartite graph
	2	2	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^2	0	0	0	1	0	0	0	1	0	1	0	0	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^4	0	0	0	1	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	1	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{13}	1	0	0	0	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{14}	1	0	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^{15}	1	0	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{16}	1	0	0	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	A_1	A_2	A_3	A_4
	dim	2	2	2	2	2	2	2	2
S_1	2	1	1	1	1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1	Z_2	Z_3	Z_4	A_1	A_2	A_3	A_4	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	A_4	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-
	dim	2	2	2	2	1	1	1	1	1	1	1	1
T_1	1	0	0	1	1	-1	0	0	0	0	0	0	-1
T_2	1	1	1	0	0	0	0	0	-1	-1	0	0	0
T_3	1	0	1	1	0	0	-1	0	0	0	0	-1	0
T_4	1	1	0	0	1	0	0	-1	0	0	-1	0	0

v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	-1
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	
1	1	1	1	1	1	1	1	2	2	2	2	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	0	0	0	-1	0	0	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0	-1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	0	0	0	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

v^{14}	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{24}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{26}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{30}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{33}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{34}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	A_1	A_2	A_3	A_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
V_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
V_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
V_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0	0
V_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	-1
V_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	A_1	A_2	A_3	A_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$

v^5	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
Y_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	0	0	0	0	-1
Y_2	1	0	0	1	0	0	0	1	0	0	1	0	0	0	-1	-1	0	0	0
Y_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	0	-1	0
Y_4	1	0	0	0	1	0	0	0	1	1	0	0	0	-1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	Z_3	Z_4	R_1	R_2
dim	2	2	2	2	2	2
U_1	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1	Z_2	Z_3	Z_4	R_1	R_2	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	Z_3	Z_4	A_1	A_2	A_3	A_4
dim	2	2	2	2	2	2	2	2
S_1	1	1	0	0	1	-1	0	0 -1
S_2	1	0	0	1	1	0	0	-1 -1
S_3	1	1	1	0	0	-1	-1	0 0
S_4	1	0	1	1	0	0	-1	-1 0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z ₁	Z ₂	Z ₃	Z ₄	A ₁	A ₂	A ₃	A ₄	
	2	2	2	2	2	2	2	2	Bipartite graph
v ¹	0	0	0	1	0	0	0	1	H _{2/2} ¹
v ²	0	0	1	0	0	0	1	0	H _{2/2} ¹
v ³	0	1	0	0	0	1	0	0	H _{2/2} ¹
v ⁴	0	1	0	1	1	0	1	0	H _{4/4} ²
v ⁵	1	0	0	0	1	0	0	0	H _{2/2} ¹
v ⁶	1	0	1	0	0	1	0	1	H _{4/4} ²

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A ₁	A ₂	A ₃	A ₄	R ₁	R ₂
	dim	2	2	2	2	2	2
T ₁	2	1	1	1	1	-1	-1

ker $M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A ₁	A ₂	A ₃	A ₄	R ₁	R ₂	
	2	2	2	2	2	2	Bipartite graph
v ¹	0	0	0	1	0	1	H _{2/1} ¹
v ²	0	0	0	1	1	0	H _{2/1} ¹
v ³	0	0	1	0	0	1	H _{2/1} ¹
v ⁴	0	0	1	0	1	0	H _{2/1} ¹
v ⁵	0	1	0	0	0	1	H _{2/1} ¹
v ⁶	0	1	0	0	1	0	H _{2/1} ¹
v ⁷	1	0	0	0	0	1	H _{2/1} ¹
v ⁸	1	0	0	0	1	0	H _{2/1} ¹

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R ₁	R ₂	X ₁ ⁺	X ₁ ⁻	X ₂ ⁺	X ₂ ⁻	X ₃ ⁺	X ₃ ⁻	X ₄ ⁺	X ₄ ⁻
	dim	2	2	1	1	1	1	1	1	1	1
W ₁	1	1	0	-1	0	0	0	0	-1	0	0
W ₂	1	0	1	0	0	-1	0	0	0	0	-1
W ₃	1	1	0	0	-1	0	0	-1	0	0	0
W ₄	1	0	1	0	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	R_1	R_2	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	2	2	1	1	1	1	1	1	1	1	
v^1	0	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^3	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^5	1	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^6	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^8	1	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$

SG 133

Space group $P4_2/nbc$, No. 133, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	0	-1	0
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^8	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$

v^{10}	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	1	1	1	1	$H_{5/4}^3$
v^{12}	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^{15}	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
dim		1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	-1
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	0	0	-1	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	-1	-1	0	0
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	0	0	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	0	1	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{25}	0	0	1	0	1	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	1	0	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{30}	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{31}	0	1	0	0	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$

v^{32}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{39}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{40}	1	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{41}	1	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	A_1	A_2	A_3A_4
	dim	2	2	2	2	2	2	4
V_1	1	0	0	0	1	0	-1	0
V_2	1	0	0	0	1	-1	0	0
V_3	1	0	1	0	0	-1	0	0
V_4	1	0	1	0	0	0	-1	0
V_5	2	1	0	1	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	A_1	A_2	A_3A_4	
	2	2	2	2	2	2	4	Bipartite graph
v^1	0	0	2	0	0	0	1	$H_{3/2}^1$
v^2	0	1	0	1	1	1	0	$H_{4/4}^2$
v^3	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	2	0	0	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	X_1	X_2
	dim	2	2	2	2	2	2
Y_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_1	Z_2	Z_3	Z_4	R_1	R_2
	dim	2	2	2	2	2	2
U_1	1	1	0	0	1	-1	0
U_2	1	0	0	1	1	0	-1
U_3	1	1	1	0	0	-1	0
U_4	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	Z_3	Z_4	R_1	R_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	1	1	1	$H_{4/4}^2$
v^3	1	0	0	0	1	0	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z ₁	Z ₂	Z ₃	Z ₄	A ₁	A ₂	A ₃ A ₄
	dim	2	2	2	2	2	2	4
S ₁	2	1	1	1	1	-1	-1	-2

ker $M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Z ₁	Z ₂	Z ₃	Z ₄	A ₁	A ₂	A ₃ A ₄	
	2	2	2	2	2	2	4	Bipartite graph
v ¹	0	0	0	1	0	1	0	H _{2/1} ¹
v ²	0	0	0	1	1	0	0	H _{2/1} ¹
v ³	0	0	0	2	0	0	1	H _{3/2} ¹
v ⁴	0	0	1	0	0	1	0	H _{2/1} ¹
v ⁵	0	0	1	0	1	0	0	H _{2/1} ¹
v ⁶	0	0	1	1	0	0	1	H _{3/2} ¹
v ⁷	0	0	2	0	0	0	1	H _{3/2} ¹
v ⁸	0	1	0	0	0	1	0	H _{2/1} ¹
v ⁹	0	1	0	0	1	0	0	H _{2/1} ¹
v ¹⁰	0	1	0	1	0	0	1	H _{3/2} ¹
v ¹¹	0	1	1	0	0	0	1	H _{3/2} ¹
v ¹²	0	2	0	0	0	0	1	H _{3/2} ¹
v ¹³	1	0	0	0	0	1	0	H _{2/1} ¹
v ¹⁴	1	0	0	0	1	0	0	H _{2/1} ¹
v ¹⁵	1	0	0	1	0	0	1	H _{3/2} ¹
v ¹⁶	1	0	1	0	0	0	1	H _{3/2} ¹
v ¹⁷	1	1	0	0	0	0	1	H _{3/2} ¹
v ¹⁸	2	0	0	0	0	0	1	H _{3/2} ¹

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A ₁	A ₂	A ₃ A ₄	R ₁	R ₂
	dim	2	2	4	2	2
T ₁	2	1	1	2	-1	-1

ker $M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A ₁	A ₂	A ₃ A ₄	R ₁	R ₂	
	2	2	4	2	2	Bipartite graph
v ¹	0	0	1	0	2	H _{3/2} ¹
v ²	0	0	1	1	1	H _{3/2} ¹
v ³	0	0	1	2	0	H _{3/2} ¹

v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	R_2	X_1	X_2
dim		2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_1	R_2	X_1	X_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 134

Space group $P4_2/nmm$, No. 134, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
dim		1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	0	-1	0
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^8	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{12}	0	1	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{15}	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	0	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	-1	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	-1
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

v^6	0	0	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	0	0	-1	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	-1	-1	0	0
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	0	0	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$

v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	0	1	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{25}	0	0	1	0	1	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	1	0	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{30}	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{31}	0	1	0	0	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{39}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{40}	1	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{41}	1	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-
	dim	2	2	2	2	1	1	1	1	1	1	1	1	2	2
V_1	1	0	0	0	1	0	0	0	0	0	-1	-1	0	0	0
V_2	1	0	0	0	1	-1	0	0	-1	0	0	0	0	0	0
V_3	1	0	1	0	0	0	-1	-1	0	0	0	0	0	0	0
V_4	1	0	1	0	0	0	0	0	0	-1	0	0	-1	0	0
V_5	2	1	0	1	0	0	0	0	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	Bipartite graph
	2	2	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^2	0	0	0	1	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	1	1	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	1	0	0	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $X(0, \frac{1}{2}, 0)$
A matrix

corep	M_1	M_2	M_3	M_4	X_1	X_2
dim	2	2	2	2	2	2
Y_1	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	R_1	R_2
	dim	2	2	2	2	2	2
U_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1	Z_2	Z_3	Z_4	R_1	R_2	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S : $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-
	dim	2	2	2	2	1	1	1	1	1	1	1	1	2	2
S_1	1	1	0	0	1	0	0	0	-1	0	-1	0	0	-1	0
S_2	1	0	0	1	1	-1	0	0	0	0	0	-1	0	0	-1
S_3	1	1	1	0	0	0	0	-1	0	-1	0	0	0	0	-1
S_4	1	0	1	1	0	0	-1	0	0	0	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Z_1	Z_2	Z_3	Z_4	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	Bipartite graph
	2	2	2	2	1	1	1	1	1	1	1	2	2		
v^1	0	0	0	1	0	0	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^2	0	0	0	1	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	1	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^6	0	0	1	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^7	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	1	1	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	1	1	0	0	0	0	0	0	2	0	1	0	$H_{5/4}^3$
v^{10}	0	0	1	1	1	0	0	0	0	0	1	0	1	0	$H_{5/4}^3$

v ¹¹	0	0	1	1	2	0	0	0	0	0	0	0	0	1	0	H _{5/4} ³
v ¹²	0	1	0	0	0	0	0	0	1	0	0	1	0	0	0	H _{3/2} ¹
v ¹³	0	1	0	0	0	0	1	0	0	0	0	0	1	0	0	H _{3/2} ¹
v ¹⁴	0	1	0	0	0	1	0	0	1	0	0	0	0	0	0	H _{3/2} ¹
v ¹⁵	0	1	0	0	0	1	1	0	0	0	0	0	0	0	0	H _{3/2} ¹
v ¹⁶	0	1	0	1	0	0	0	0	0	0	0	0	0	1	1	H _{4/4} ²
v ¹⁷	0	1	0	1	0	0	0	0	0	1	0	1	0	1	0	H _{5/4} ³
v ¹⁸	0	1	0	1	0	0	0	0	1	0	1	0	1	0	0	H _{5/4} ³
v ¹⁹	0	1	0	1	0	0	0	1	0	0	0	1	0	1	0	H _{5/4} ³
v ²⁰	0	1	0	1	0	0	1	0	0	0	1	0	1	0	0	H _{5/4} ³
v ²¹	0	1	0	1	0	1	0	0	0	1	0	0	0	0	1	H _{5/4} ³
v ²²	0	1	0	1	0	1	0	1	0	0	0	0	0	0	1	H _{5/4} ³
v ²³	0	1	0	1	1	0	0	0	1	0	0	0	0	1	0	H _{5/4} ³
v ²⁴	0	1	0	1	1	0	1	0	0	0	0	0	0	1	0	H _{5/4} ³
v ²⁵	0	1	1	0	0	0	0	0	0	0	0	0	2	0	1	H _{5/4} ³
v ²⁶	0	1	1	0	0	1	0	0	0	0	0	0	1	0	1	H _{5/4} ³
v ²⁷	0	1	1	0	0	2	0	0	0	0	0	0	0	0	1	H _{5/4} ³
v ²⁸	1	0	0	0	0	0	0	0	1	1	0	0	0	0	0	H _{3/2} ¹
v ²⁹	1	0	0	0	0	0	0	1	1	0	0	0	0	0	0	H _{3/2} ¹
v ³⁰	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	H _{3/2} ¹
v ³¹	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	H _{3/2} ¹
v ³²	1	0	0	1	0	0	0	0	0	2	0	0	0	0	1	H _{5/4} ³
v ³³	1	0	0	1	0	0	0	1	0	1	0	0	0	0	1	H _{5/4} ³
v ³⁴	1	0	0	1	0	0	0	2	0	0	0	0	0	0	1	H _{5/4} ³
v ³⁵	1	0	1	0	0	0	0	0	0	0	0	0	0	1	1	H _{4/4} ²
v ³⁶	1	0	1	0	0	0	0	0	0	1	0	1	0	1	0	H _{5/4} ³
v ³⁷	1	0	1	0	0	0	0	0	1	0	1	0	1	0	0	H _{5/4} ³
v ³⁸	1	0	1	0	0	0	0	1	0	0	0	0	1	0	1	H _{5/4} ³
v ³⁹	1	0	1	0	0	0	1	0	0	0	1	0	1	0	0	H _{5/4} ³
v ⁴⁰	1	0	1	0	0	1	0	0	0	1	0	0	0	0	1	H _{5/4} ³
v ⁴¹	1	0	1	0	0	1	0	1	0	0	0	0	0	0	1	H _{5/4} ³
v ⁴²	1	0	1	0	1	0	0	0	1	0	0	0	0	1	0	H _{5/4} ³
v ⁴³	1	0	1	0	1	0	1	0	0	0	0	0	0	1	0	H _{5/4} ³
v ⁴⁴	1	1	0	0	0	0	0	0	2	0	0	0	0	1	0	H _{5/4} ³
v ⁴⁵	1	1	0	0	0	0	1	0	1	0	0	0	0	1	0	H _{5/4} ³
v ⁴⁶	1	1	0	0	0	0	2	0	0	0	0	0	0	1	0	H _{5/4} ³

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	R_1	R_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
T_1	1	0	1	0	0	0	1	0	0	1	0	-1	0
T_2	1	1	0	0	0	1	0	0	0	0	1	-1	0
T_3	1	0	0	1	0	0	0	1	0	0	1	0	-1
T_4	1	0	0	0	1	0	0	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	R_1	R_2	
	1	1	1	1	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^8	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{12}	0	1	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{15}	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_1	R_2	X_1	X_2
	dim	2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

R_1	R_2	X_1	X_2	
2	2	2	2	Bipartite graph

v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 135

Space group $P4_2/mbc$, No. 135, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix



corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	-1
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-2	0	0	0	0	-1
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-2	0	0	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	0	0	-2	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	0	-2	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	2	1	0	1	0	0	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	2	0	0	1	0	1	0	0	$H_{4/4}^2$
v^5	0	0	0	0	0	0	0	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	2	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	1	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^9	0	0	0	0	0	2	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{12}	0	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	0	1	0	0	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	1	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{16}	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{19}	0	0	1	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	1	0	0	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{24}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	0	1	0	1	0	0	0	0	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{26}	0	2	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{28}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{29}	1	0	0	0	1	0	0	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{30}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{31}	1	0	1	0	0	0	0	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$
v^{32}	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	$A_1 A_2$	$A_3 A_4$
	dim	2	2	2	2	2	2	4	4
$V_1 V_2$	2	1	1	0	0	0	0	-1	0
$V_3 V_4$	2	0	0	1	1	0	0	-1	0

V_5	2	0	0	0	0	1	1	0	-2
-------	---	---	---	---	---	---	---	---	----

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	A_1A_2	A_3A_4	Bipartite graph
	2	2	2	2	2	2	4	4	
v^1	0	0	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^6	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $X(0, \frac{1}{2}, 0)$
A matrix

corep		$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	X_1	X_2
	dim	2	2	2	2	2	2	2	2
Y_1Y_2	2	1	0	1	0	0	1	-1	0
Y_3Y_4	2	0	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	X_1	X_2	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_1	Z_2	Z_3	Z_4	R_1	R_2
dim	2	2	2	2	2	2

U_1	1	1	0	0	1	-1	0
U_2	1	0	0	1	1	0	-1
U_3	1	1	1	0	0	0	-1
U_4	1	0	1	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	Z_2	Z_3	Z_4	R_1	R_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	$H_{4/4}^2$
v^2	1	0	1	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S : $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	A_1A_2	A_3A_4
	dim	2	2	2	2	4	4
S_1	2	1	1	1	1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Z_1	Z_2	Z_3	Z_4	A_1A_2	A_3A_4	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	0	2	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	1	$H_{3/2}^1$
v^4	0	0	1	1	1	0	$H_{3/2}^1$
v^5	0	0	2	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	0	$H_{3/2}^1$
v^9	0	1	1	0	0	1	$H_{3/2}^1$
v^{10}	0	1	1	0	1	0	$H_{3/2}^1$
v^{11}	0	2	0	0	0	1	$H_{3/2}^1$
v^{12}	0	2	0	0	1	0	$H_{3/2}^1$
v^{13}	1	0	0	1	0	1	$H_{3/2}^1$
v^{14}	1	0	0	1	1	0	$H_{3/2}^1$
v^{15}	1	0	1	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	1	0	$H_{3/2}^1$
v^{17}	1	1	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	1	0	$H_{3/2}^1$

v^{19}	2	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3A_4	R_1	R_2
	dim	4	4	2	2
T_1T_2	2	1	1	-1	0
T_3T_4	2	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_2	A_3A_4	R_1	R_2	
	4	4	2	2	Bipartite graph
v^1	0	1	1	1	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_1	R_2	X_1	X_2
	dim	2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_1	R_2	X_1	X_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 136

Space group $P4_2/mnm$, No. 136, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	0	-1
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	
	1	1	1	1	1	1	1	1	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	-1
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-
dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	1	0	0	1	-2	0	0	0	0	-1
Σ_2	1	0	0	1	0	1	0	0	0	1	0	0	-2	0	0	-1
Σ_3	1	0	1	0	0	0	0	1	1	0	0	0	0	-2	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	0	-2	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$M_1^+M_2^+$	$M_1^-M_2^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	M_5^+	M_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	2	1	0	1	0	0	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	2	0	0	1	0	1	0	0	$H_{4/4}^2$
v^5	0	0	0	0	0	0	0	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	2	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	1	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^9	0	0	0	0	0	2	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	1	0	1	0	0	0	$H_{5/4}^3$

v^{12}	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	0	1	0	0	0	1	1	0	0	1	0	0	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{16}	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	1	0	0	1	1	0	0	0	0	$H_{5/4}^3$
v^{19}	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{20}	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{22}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	1	0	0	1	0	0	1	0	0	0	$H_{5/4}^3$
v^{24}	0	1	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	0	1	0	1	0	0	0	0	1	0	0	1	0	0	0	$H_{5/4}^3$
v^{26}	0	2	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{28}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{29}	1	0	0	0	1	0	0	0	0	1	1	0	0	0	0	$H_{5/4}^3$
v^{30}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{31}	1	0	1	0	0	0	0	0	0	1	1	0	0	0	0	$H_{5/4}^3$
v^{32}	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	A_1	A_2	A_3	A_4
dim	2	2	2	2	2	2	2	2	2	2
$V_1 V_2$	2	1	1	0	0	0	-1	0	0	0
$V_3 V_4$	2	0	0	1	1	0	0	0	-1	0
V_5	2	0	0	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	A_1	A_2	A_3	A_4	Bipartite graph
	2	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$

v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	X_1	X_2	
dim	2	2	2	2	2	2	2	2	
$Y_1 Y_2$	2	1	0	1	0	0	1	-1	0
$Y_3 Y_4$	2	0	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$M_1^+ M_2^+$	$M_1^- M_2^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	M_5^+	M_5^-	X_1	X_2	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^2	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_1	Z_2	Z_3	Z_4	R_1^+	R_1^-	
dim	2	2	2	2	2	2	
U_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_1	Z_2	Z_3	Z_4	R_1^+	R_1^-	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	A_1	A_2	A_3	A_4
dim		2	2	2	2	2	2	2	2
S_1	1	1	0	0	1	-1	0	0	-1
S_2	1	0	0	1	1	-1	-1	0	0
S_3	1	1	1	0	0	0	0	-1	-1
S_4	1	0	1	1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_1	Z_2	Z_3	Z_4	A_1	A_2	A_3	A_4	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	1	0	0	0	$H_{2/2}^1$
v^2	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	1	0	1	0	1	0	1	$H_{4/4}^2$
v^5	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^6	1	0	1	0	1	0	1	0	$H_{4/4}^2$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	R_1^+	R_1^-
dim		2	2	2	2	2	2
T_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_1	A_2	A_3	A_4	R_1^+	R_1^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$

v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_1^+	R_1^-	X_1	X_2
dim		2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_1^+	R_1^-	X_1	X_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 137

Space group $P4_2/nmc$, No. 137, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
dim		1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	-1	0
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	
1	1	1	1	1	1	1	1	2	2	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^3	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^6	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^7	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{11}	0	1	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{13}	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	0	0	-1
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	0
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2		
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$

v^8	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
dim		1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	-1	0	0
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	1	1	0	0	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

v^{22}	0	0	1	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{24}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{25}	0	0	1	0	1	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	1	0	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{30}	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{31}	0	1	0	0	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{39}	1	0	0	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{40}	1	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{41}	1	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_1	M_2	M_3	M_4	A_1	A_2	A_3A_4
dim	2	2	2	2	2	2	4
V_1V_4	2	1	0	0	0	0	-1
V_2V_3	2	0	0	1	0	0	-1
V_5	2	0	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	M_3	M_4	A_1	A_2	A_3A_4	
	2	2	2	2	2	2	4	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	$H_{2/1}^1$

v^3	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	0	1	0	0	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	X_1	X_2
	dim	2	2	2	2	2	2
Y_1	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	R_1	R_2
	dim	2	2	2	2	2	2
U_1	1	1	0	0	1	-1	0
U_2	1	0	0	1	1	-1	0
U_3	1	1	1	0	0	0	-1
U_4	1	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

Z_1	Z_2	Z_3	Z_4	R_1	R_2	
2	2	2	2	2	2	Bipartite graph

v^1	0	0	0	1	1	0	$H_{2/2}^1$
v^2	0	1	0	0	0	1	$H_{2/2}^1$
v^3	1	0	1	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	A_1	A_2	A_3A_4
	dim	2	2	2	2	2	2	4
S_1	2	1	1	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Z_1	Z_2	Z_3	Z_4	A_1	A_2	A_3A_4	
	2	2	2	2	2	2	4	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	$H_{2/1}^1$
v^5	0	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	1	0	0	1	$H_{3/2}^1$
v^7	0	0	2	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	1	0	1	0	0	1	$H_{3/2}^1$
v^{11}	0	1	1	0	0	0	1	$H_{3/2}^1$
v^{12}	0	2	0	0	0	0	1	$H_{3/2}^1$
v^{13}	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{18}	2	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix



corep	A_1	A_2	A_3A_4	R_1	R_2
dim	2	2	4	2	2
T_1	2	1	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3A_4	R_1	R_2	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_1	R_2	X_1	X_2
dim	2	2	2	2
W_1W_4	2	1	0	-1 0
W_2W_3	2	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_1	R_2	X_1	X_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 138

Space group $P4_2/nm$, No. 138, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0
Δ_2	1	0	0	0	1	0	0	0	1	1	0	-1	0
Δ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1
Δ_4	1	0	0	1	0	0	0	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^3	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^6	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^7	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{10}	0	1	0	0	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{11}	0	1	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{13}	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Λ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0
Λ_3	1	0	0	0	0	1	0	0	1	0	0	0	-1	0	0
Λ_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	-1
Λ_5	2	0	0	0	0	0	0	0	0	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	0	0	1	0	0	1	-1	-1	0	0
Σ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	-1
Σ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	0	1	1	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	1	1	0	0	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$

V_1V_4	2	1	0	0	0	-1	-1	0	0	0	0
V_2V_3	2	0	0	1	0	0	0	-1	-1	0	0
V_5	2	0	1	0	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	$A_1^+A_2^+$	$A_1^-A_2^-$	$A_3^+A_4^+$	$A_3^-A_4^-$	A_5^+	A_5^-	Bipartite graph
	2	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_1	M_2	M_3	M_4	X_1	X_2
dim	2	2	2	2	2	2
Y_1	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	X_1	X_2	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	$R_1 R_2$
dim		2	2	2	2	4
U_1	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_1	Z_2	Z_3	Z_4	$R_1 R_2$	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S : $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_1	Z_2	Z_3	Z_4	$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-
dim		2	2	2	2	2	2	2	2	2	2
S_1	1	1	0	0	1	0	-2	0	0	-1	0
S_2	1	0	0	1	1	-2	0	0	0	0	-1
S_3	1	1	1	0	0	0	0	-2	0	0	-1
S_4	1	0	1	1	0	0	0	0	-2	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 15.

	Z_1	Z_2	Z_3	Z_4	$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-	
	2	2	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	2	1	1	0	0	0	0	$H_{4/4}^2$
v^2	0	0	1	1	1	0	0	0	1	0	$H_{4/4}^2$
v^3	0	0	2	0	1	0	0	1	0	0	$H_{4/4}^2$
v^4	0	1	0	1	0	0	0	0	1	1	$H_{4/4}^2$
v^5	0	1	1	0	0	0	0	1	0	1	$H_{4/4}^2$
v^6	0	2	0	0	0	0	1	1	0	0	$H_{4/4}^2$
v^7	0	2	0	2	0	1	0	1	0	2	$H_{8/8}^1$
v^8	0	2	0	2	1	0	1	0	2	0	$H_{8/8}^1$

v^9	1	0	0	1	0	1	0	0	0	1	$H_{4/4}^2$
v^{10}	1	0	1	0	0	0	0	0	1	1	$H_{4/4}^2$
v^{11}	1	1	0	0	0	0	1	0	1	0	$H_{4/4}^2$
v^{12}	1	1	1	1	1	1	1	1	0	0	$H_{8/8}^1$
v^{13}	2	0	0	0	0	1	1	0	0	0	$H_{4/4}^2$
v^{14}	2	0	2	0	0	1	0	1	0	2	$H_{8/8}^1$
v^{15}	2	0	2	0	1	0	1	0	2	0	$H_{8/8}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-	$R_1 R_2$
	dim	2	2	2	2	2	2	4
$T_1 T_4$	2	1	0	1	0	0	1	-1
$T_2 T_3$	2	0	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	$A_1^+ A_2^+$	$A_1^- A_2^-$	$A_3^+ A_4^+$	$A_3^- A_4^-$	A_5^+	A_5^-	$R_1 R_2$	
	2	2	2	2	2	2	4	Bipartite graph
v^1	0	0	0	0	1	1	1	$H_{3/2}^1$
v^2	0	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	1	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	1	$H_{3/2}^1$
v^8	1	0	0	1	0	0	1	$H_{3/2}^1$
v^9	1	1	0	0	0	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		$R_1 R_2$	X_1	X_2
	dim	4	2	2
$W_1 W_4$	2	1	-1	0
$W_2 W_3$	2	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$R_1 R_2$	X_1	X_2	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 139

Space group $I4/mmm$, No. 139, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
Σ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	0	-1	0	0	0	0	0	-1
Σ_2	1	0	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	-1	0	0	0	-1
Σ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	-1	0	0	-1	0	0
Σ_4	1	0	0	0	1	0	0	0	1	1	0	0	0	0	-1	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

v^{20}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{24}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{26}	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{28}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{30}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{33}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{34}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
Δ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	-1	0	0	0	0
Δ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	0	0	-1	0	0	-1
Δ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0	0	0	0	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

v^{12}	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_1^+	N_1^-	N_2^+	N_2^-	P_1	P_2	P_3	P_4	P_5
	dim	1	1	1	1	1	1	1	1	2
Q_1	1	1	1	0	0	-1	0	-1	0	-1
Q_2	1	0	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	N_1^+	N_1^-	N_2^+	N_2^-	P_1	P_2	P_3	P_4	P_5	Bipartite graph
	1	1	1	1	1	1	1	1	2	
v^1	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^7	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	P_1	P_2	P_3	P_4	P_5
	dim	1	1	1	1	1	1	1	1	1	1	1	1	2
W_1	1	1	0	0	0	0	1	0	0	-1	0	0	-1	0
W_2	1	0	0	0	1	0	0	1	0	0	0	0	0	-1
W_3	1	0	1	0	0	1	0	0	0	0	-1	-1	0	0
W_4	1	0	0	1	0	0	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	P_1	P_2	P_3	P_4	P_5	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	1	1	2	
v^1	0	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-
	dim	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2
Y_1	1	1	0	0	0	0	0	0	1	-1	0	0	0	0	0	-1	0	0	-1
Y_2	1	0	0	1	0	0	1	0	0	0	0	0	-1	0	-1	0	0	-1	0
Y_3	1	0	1	0	0	0	0	1	0	0	-1	0	0	0	0	0	-1	-1	0
Y_4	1	0	0	0	1	1	0	0	0	0	0	-1	0	-1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{20}	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{22}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{24}	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 140

Space group $I4/mcm$, No. 140, without SOC

$$\Sigma$$

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
Σ_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	0	-1	0	0	0	0	0	-1
Σ_2	1	0	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	-1	0	0	0	-1
Σ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	-1	0	0	-1	0	0
Σ_4	1	0	0	0	1	0	0	0	1	1	0	0	0	0	-1	0	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{18}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{24}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{26}	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{28}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{30}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{33}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{34}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1
Δ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	-1	0	0	0	0
Δ_2	1	0	0	1	0	1	0	0	0	0	1	0	0	0	0	-1	0	0	-1
Δ_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	-1	0	0	0	0	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	N_1	P_1P_4	P_2P_3	P_5
dim	2	2	2	2
Q_1Q_2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	N_1	P_1P_4	P_2P_3	P_5	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	P_1P_4	P_2P_3	P_5
dim		1	1	1	1	1	1	1	1	2	2	2
W_1	1	1	0	0	0	0	1	0	0	-2	0	0
W_2	1	0	0	0	1	0	0	1	0	0	0	-1
W_3	1	0	1	0	0	1	0	0	0	0	-2	0
W_4	1	0	0	1	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{11}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	P_1P_4	P_2P_3	P_5	
	1	1	1	1	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^4	0	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	2	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	2	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-
dim		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2
Y_1	1	1	0	0	0	0	0	0	1	-1	0	0	0	0	0	-1	0	0	-1
Y_2	1	0	0	1	0	0	1	0	0	0	0	0	-1	0	-1	0	0	-1	0
Y_3	1	0	1	0	0	0	0	1	0	0	-1	0	0	0	0	0	-1	-1	0
Y_4	1	0	0	0	1	1	0	0	0	0	0	-1	0	-1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	Z_4^+	Z_4^-	Z_5^+	Z_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{20}	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{22}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{24}	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 141

Space group $I4_1/amd$, No. 141, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
dim		1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	1	0	0	0	0	1	0	-1	-1	0
Σ_2	1	0	0	1	0	0	0	1	0	0	1	-1	0	0	-1
Σ_3	1	0	1	0	0	0	1	0	0	1	0	0	0	-1	-1
Σ_4	1	0	0	0	1	0	0	0	1	1	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	1	1	0	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	1	0	0	1	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	1	1	0	0	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	1	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	1	0	0	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	1	0	1	0	1	0	0	1	$H_{5/4}^3$
v^{23}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{24}	0	0	1	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{25}	0	0	1	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	1	0	0	1	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$

v^{29}	0	1	0	0	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{30}	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{31}	0	1	0	0	0	1	0	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{39}	1	0	0	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{40}	1	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{41}	1	0	0	0	1	0	0	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	0	1	1	0	$H_{5/4}^3$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0
Δ_2	1	0	0	1	0	1	0	0	0	0	1	0	-1
Δ_3	1	0	1	0	0	0	0	0	1	1	0	-1	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^7	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$

v^9	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_1^+	N_1^-	N_2^+	N_2^-	P_1	P_2
	dim	1	1	1	1	2	2
Q_1	1	1	1	0	0	-1	-1
Q_2	1	0	0	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	N_1^+	N_1^-	N_2^+	N_2^-	P_1	P_2	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	1	0	$H_{3/2}^1$
v^5	1	0	0	1	0	1	$H_{3/2}^1$
v^6	1	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	1	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_1	X_2	P_1	P_2
	dim	2	2	2	2
W_1	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	P_1	P_2	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1	X_2	Z_1	Z_2	Z_3	Z_4
dim	2	2	2	2	2	2
Y_1	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_1	X_2	Z_1	Z_2	Z_3	Z_4	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	1	0	0	0	$H_{2/1}^1$

SG 142

Space group $I4_1/acd$, No. 142, without SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4
dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2
Σ_1	1	1	0	0	0	1	0	0	0	0	1	0	0	-1
Σ_2	1	0	0	1	0	0	0	1	0	0	1	-1	-1	0
Σ_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	-1
Σ_4	1	0	0	0	1	0	0	0	1	1	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Z_1	Z_2	Z_3	Z_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	2	0	1	1	0	0	1	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	1	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	2	0	1	0	1	1	0	0	$H_{5/4}^3$
v^6	0	0	0	0	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	2	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{10}	0	0	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	0	0	2	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{15}	0	0	0	1	0	0	0	1	0	1	1	0	0	1	$H_{5/4}^3$
v^{16}	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	1	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{18}	0	0	0	1	0	1	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{19}	0	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	0	0	2	0	0	0	0	0	1	1	0	0	1	$H_{5/4}^3$
v^{21}	0	0	1	0	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	1	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{23}	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{24}	0	0	1	0	1	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{25}	0	0	1	0	1	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{26}	0	0	1	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	0	0	2	0	0	0	0	0	1	0	1	1	0	0	$H_{5/4}^3$
v^{28}	0	1	0	0	0	0	0	1	0	1	0	1	0	1	$H_{5/4}^3$
v^{29}	0	1	0	0	0	0	0	1	0	1	1	0	1	0	$H_{5/4}^3$
v^{30}	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{31}	0	1	0	0	0	1	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{32}	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{33}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{34}	0	1	0	1	0	0	0	0	0	1	1	0	1	0	$H_{5/4}^3$
v^{35}	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{36}	0	2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{37}	1	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{38}	1	0	0	0	0	0	1	0	1	0	0	1	0	1	$H_{5/4}^3$

v^{39}	1	0	0	0	0	0	1	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{40}	1	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{41}	1	0	0	0	1	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{42}	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{43}	1	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{5/4}^3$
v^{44}	1	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{45}	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{46}	2	0	0	0	0	0	0	0	1	0	0	0	1	1	$H_{5/4}^3$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2
	dim	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	0	0	0	1	0	0	1	-1	0
Δ_2	1	0	0	1	0	1	0	0	0	0	1	0	-1
Δ_3	1	0	1	0	0	0	0	0	1	1	0	-1	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^5	0	0	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	1	0	1	1	1	$H_{5/4}^3$
v^7	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	0	0	0	1	0	1	0	1	1	$H_{5/4}^3$
v^9	0	0	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^{13}	0	1	0	1	0	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	0	1	0	0	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	N_1	$P_1 P_2$
dim	2	4
$Q_1 Q_2$	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

N_1	$P_1 P_2$	
2	4	Bipartite graph
v^1	2	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	X_1	X_2	$P_1 P_2$
dim	2	2	4
W_1	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_1	X_2	$P_1 P_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	Z_1	Z_2	Z_3	Z_4
dim	2	2	2	2	2	2
Y_1	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_1	X_2	Z_1	Z_2	Z_3	Z_4	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	1	0	0	0	$H_{2/1}^1$

SG 143

Space group $P3$, No. 143, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	A_1	A_2A_3
	dim	1	2	1	2
Δ_1	1	1	0	-1	0
Δ_2	1	0	1	0	-1
Δ_3	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	A_1	A_2A_3	
	1	2	1	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	K_1	K_2	K_3
	dim	1	2	1	1	1
T_1	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	K_1	K_2	K_3	
	1	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	M_1
dim	1	2	1
Σ_1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	M_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1	A_2A_3	H_1	H_2	H_3
dim	1	2	1	1	1
S_1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

A_1	A_2A_3	H_1	H_2	H_3	
1	2	1	1	1	Bipartite graph

v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R : $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2A_3	L_1
dim	1	2	1
R_1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1	A_2A_3	L_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 144

Space group $P3_1$, No. 144, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	A_1	A_2A_3
dim	1	2	1	2
Δ_1	1	0	0	-1
Δ_2	0	1	0	-1
Δ_3	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_1	$\Gamma_2\Gamma_3$	A_1	A_2A_3	
	1	2	1	2	Bipartite graph
v^1	1	1	1	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	K_1	K_2	K_3
dim	1	2	1	1	1
T_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	K_1	K_2	K_3	
	1	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	M_1
dim	1	2	1
Σ_1	1	1	2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	M_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1	A_2A_3	H_1	H_2	H_3
dim	1	2	1	1	1
S_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1	A_2A_3	H_1	H_2	H_3	
	1	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2A_3	L_1
dim	1	2	1
R_1	1	1	2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1	A_2A_3	L_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 145

Space group $P3_2$, No. 145, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	A_1	A_2A_3
dim		1	2	1	2
Δ_1	1	1	0	0	-1
Δ_2	1	0	1	-1	0
Δ_3	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		Γ_1	$\Gamma_2\Gamma_3$	A_1	A_2A_3	
		1	2	1	2	Bipartite graph
v^1	1	1	1	1	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	K_1	K_2	K_3
dim		1	2	1	1	1
T_1	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

		Γ_1	$\Gamma_2\Gamma_3$	K_1	K_2	K_3	
		1	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2		$H_{3/2}^1$
v^2	0	1	0	1	1		$H_{3/2}^1$
v^3	0	1	0	2	0		$H_{3/2}^1$
v^4	0	1	1	0	1		$H_{3/2}^1$
v^5	0	1	1	1	0		$H_{3/2}^1$
v^6	0	1	2	0	0		$H_{3/2}^1$
v^7	1	0	0	0	1		$H_{2/1}^1$
v^8	1	0	0	1	0		$H_{2/1}^1$
v^9	1	0	1	0	0		$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	M_1
dim	1	2	1
Σ_1	1	1	2
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	M_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1	A_2A_3	H_1	H_2	H_3
dim	1	2	1	1	1
S_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1	A_2A_3	H_1	H_2	H_3	
	1	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2A_3	L_1
dim	1	2	1
R_1	1	1	2
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1	A_2A_3	L_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 146

Space group $R3$, No. 146, without SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	F_1
dim	1	2	1
Σ_1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	F_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y : $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep	Z_1	Z_2Z_3	L_1
dim	1	2	1
Y_1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1	Z_2Z_3	L_1	
	1	2	1	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 147

Space group $P\bar{3}$, No. 147, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$
	dim	1	1	2	2	1	1	2	2
Δ_1	1	1	1	0	0	-1	-1	0	0
$\Delta_2\Delta_3$	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$	Bipartite graph
	1	1	2	2	1	1	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	K_1	K_2K_3
	dim	1	1	2	2	1	2
T_1	1	1	1	2	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	K_1	K_2K_3	Bipartite graph
	1	1	2	2	1	2	
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	0	1	$H_{2/2}^1$
v^4	0	0	1	0	2	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$

v^6	0	2	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	1	0	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	M_1^+	M_1^-
dim	1	1	2	2	1	1
Σ_1	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	M_1^+	M_1^-	Bipartite graph
	1	1	2	2	1	1	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_1^+	M_1^-	K_1	K_2K_3
dim	1	1	1	2
T'_1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

M_1^+	M_1^-	K_1	K_2K_3	Bipartite graph
1	1	1	2	

v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1^+	A_1^-	$A_2^+ A_3^+$	$A_2^- A_3^-$	H_1	$H_2 H_3$
dim	1	1	2	2	1	2
S_1	1	1	2	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1^+	A_1^-	$A_2^+ A_3^+$	$A_2^- A_3^-$	H_1	$H_2 H_3$	
	1	1	2	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	0	1	$H_{2/2}^1$
v^4	0	0	1	0	2	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	2	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	1	0	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1^+	A_1^-	$A_2^+ A_3^+$	$A_2^- A_3^-$	L_1^+	L_1^-
dim	1	1	2	2	1	1
R_1	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1^+	A_1^-	$A_2^+ A_3^+$	$A_2^- A_3^-$	L_1^+	L_1^-	
	1	1	2	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_1^+	L_1^-	H_1	$H_2 H_3$
	dim	1	1	1	2
S'_1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1^+	L_1^-	H_1	$H_2 H_3$	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_1	$K_2 K_3$	H_1	$H_2 H_3$
	dim	1	2	1	2
P_1	1	1	0	-1	0
$P_2 P_3$	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_1	K_2K_3	H_1	H_2H_3	
	1	2	1	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 148

Space group $R\bar{3}$, No. 148, without SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	F_1^+	F_1^-
	dim	1	1	2	2	1	1
Σ_1	1	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	F_1^+	F_1^-	
	1	1	2	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y : $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep		Z_1^+	Z_1^-	$Z_2^+Z_3^+$	$Z_2^-Z_3^-$	L_1^+	L_1^-
	dim	1	1	2	2	1	1
Y_1	1	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_1^+	Z_1^-	Z_2^+	Z_3^+	Z_2^-	Z_3^-	L_1^+	L_1^-	
	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	1	0	2	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	1	0	$H_{2/1}^1$

SG 149

Space group $P312$, No. 149, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
	dim	1	1	2	1	1	2
Δ_1	1	1	1	0	-1	-1	0
Δ_2	1	0	0	1	0	0	-1
Δ_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
dim	1	1	2	1	1	1
T_1	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	Bipartite graph
	1	1	2	1	1	1	
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	M_1	M_2
dim	1	1	2	1	1
Σ_1	1	1	0	1	-1
Σ_2	1	0	1	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	M_1	M_2	Bipartite graph
	1	1	2	1	1	
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	H_1	H_2	H_3
dim	1	1	2	1	1	1
S_1	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	L_1	L_2
dim	1	1	2	1	1
R_1	1	0	1	-1	0
R_2	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

SG 150

Space group $P321$, No. 150, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
dim		1	1	2	1	1	2
Δ_1	1	1	1	0	-1	-1	0
Δ_2	1	0	0	1	0	0	-1
Δ_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
dim		1	1	2	1	1	2
T_1	1	1	0	1	-1	0	-1
T_2	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	M_1	M_2
dim	1	1	2	1	1
Σ_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	M_1	M_2	K_1	K_2	K_3
dim	1	1	1	1	2
T'_1	1	1	0	-1	0
T'_2	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	K_1	K_2	K_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	H_1	H_2	H_3
dim	1	1	2	1	1	2
S_1	1	1	0	1	-1	0
S_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	L_1	L_2
dim	1	1	2	1	1
R_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	L_2	H_1	H_2	H_3
dim	1	1	1	1	2
S'_1	1	1	0	-1	0
S'_2	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	L_2	H_1	H_2	H_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	1	0	-1	-1	0
P_2	1	0	0	1	0	0	-1
P_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

SG 151

Space group $P3_112$, No. 151, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
	dim	1	1	2	1	1	2
Δ_1	1	1	1	0	0	0	-1
Δ_2	1	0	0	1	0	0	-1
Δ_3	1	0	0	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
dim	1	1	2	1	1	1
T_1	1			1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
	1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	M_1	M_2
dim	1	1	2	1	1
Σ_1	1			1	0
Σ_2	1			0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	H_1	H_2	H_3
dim	1	1	2	1	1	1
S_1	1			1	1	2
				-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	L_1	L_2
	dim	1	1	2	1	1
R_1	1	1	0	1	-1	0
R_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

SG 152

Space group $P3_121$, No. 152, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
	dim	1	1	2	1	1	2
Δ_1	1	1	1	0	0	0	-1
Δ_2	1	0	0	1	0	0	-1
Δ_3	1	0	0	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
	dim	1	1	2	1	1	2
T_1	1	1	0	1	-1	0	-1
T_2	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	M_1	M_2
	dim	1	1	2	1	1
Σ_1	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1	M_2	K_1	K_2	K_3
	dim	1	1	1	1	2
T'_1	1	1	0	-1	0	-1
T'_2	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	K_1	K_2	K_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
S_1	1	0	1	1	0	-1	-1
S_2	1	1	0	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	L_1	L_2
	dim	1	1	2	1	1
R_1	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	H_1	H_2	H_3
	dim	1	1	1	1	2
S'_1	1	1	0	-1	0	-1
S'_2	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	L_2	H_1	H_2	H_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	1	0	0	0	-1
P_2	1	0	0	1	0	0	-1
P_3	1	0	0	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

SG 153

Space group $P3_212$, No. 153, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
	dim	1	1	2	1	1	2
Δ_1	1	1	1	0	0	0	-1
Δ_2	1	0	0	1	-1	-1	0
Δ_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
	dim	1	1	2	1	1	1
T_1	1	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
	1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	M_1	M_2
	dim	1	1	2	1	1
Σ_1	1	1	0	1	-1	0
Σ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2})$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	H_1	H_2	H_3
	dim	1	1	2	1	1	1
S_1	1	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R : $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	L_1	L_2
	dim	1	1	2	1	1
R_1	1	1	0	1	-1	0
R_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

SG 154

Space group $P3_221$, No. 154, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
	dim	1	1	2	1	1	2
Δ_1	1	1	1	0	0	0	-1
Δ_2	1	0	0	1	-1	-1	0
Δ_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	Bipartite graph
	1	1	2	1	1	2	
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
	dim	1	1	2	1	1	2
T_1	1	1	0	1	-1	0	-1
T_2	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	Bipartite graph
	1	1	2	1	1	2	
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	Γ_2	Γ_3	M_1	M_2
dim	1	1	2	1	1
Σ_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_1	M_2	K_1	K_2	K_3
dim	1	1	1	1	2
T'_1	1	1	0	-1	0 -1
T'_2	1	0	1	0	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	K_1	K_2	K_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	H_1	H_2	H_3
dim	1	1	2	1	1	2
S_1	1	0	1	1	0	-1 -1
S_2	1	1	0	1	-1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	L_1	L_2
dim	1	1	2	1	1
R_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	L_2	H_1	H_2	H_3
dim	1	1	1	1	2
S'_1	1	1	0	-1	0
S'_2	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	L_2	H_1	H_2	H_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	1	0	0	0	-1
P_2	1	0	0	1	-1	-1	0
P_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

SG 155

Space group $R32$, No. 155, without SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	F_1	F_2
	dim	1	1	2	1	1
Σ_1	1	1	0	1	-1	0
Σ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	F_1	F_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y: $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep		Z_1	Z_2	Z_3	L_1	L_2
	dim	1	1	2	1	1
Y_1	1	1	0	1	-1	0
Y_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_1	Z_2	Z_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

SG 156

Space group $P3m1$, No. 156, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
	dim	1	1	2	1	1	2
Δ_1	1	1	0	0	-1	0	0
Δ_2	1	0	1	0	0	-1	0
Δ_3	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
dim	1	1	2	1	1	1
T_1	1			1	1	2
				-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
	1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	M_1	M_2
dim	1	1	2	1	1
Σ_1	1			1	0
Σ_2	1			0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	H_1	H_2	H_3
dim	1	1	2	1	1	1
S_1	1			1	1	2
				-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	L_1	L_2
dim	1	1	2	1	1
R_1	1			1	0
R_2	1			0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

SG 157

Space group $P31m$, No. 157, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1	A_2	A_3
	dim	1	1	2	1	1	2
Δ_1	1	1	0	0	-1	0	0
Δ_2	1	0	1	0	0	-1	0
Δ_3	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	A_1	A_2	A_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
	dim	1	1	2	1	1	2
T_1	1	1	0	1	-1	0	-1
T_2	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	M_1	M_2
dim	1	1	2	1	1
Σ_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	M_1	M_2	K_1	K_2	K_3
dim	1	1	1	1	2
T'_1	1	0	-1	0	-1
T'_2	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	K_1	K_2	K_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	H_1	H_2	H_3
dim	1	1	2	1	1	2
S_1	1	0	1	-1	0	-1
S_2	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1	A_2	A_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	L_1	L_2
dim	1	1	2	1	1
R_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$

v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	H_1	H_2	H_3
	dim	1	1	1	1	2
S'_1	1	1	0	-1	0	-1
S'_2	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	L_2	H_1	H_2	H_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	0	0	-1	0	0
P_2	1	0	1	0	0	-1	0
P_3	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

SG 158

Space group $P3c1$, No. 158, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	A_1A_2	A_3A_3
dim		1	1	2	2	4
Δ_1	1	1	0	0	-1	0
Δ_2	1	0	1	0	-1	0
Δ_3	2	0	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

		Γ_1	Γ_2	Γ_3	A_1A_2	A_3A_3	
		1	1	2	2	4	Bipartite graph
v^1	0	0	2	0	1		$H_{3/2}^1$
v^2	1	1	0	1	0		$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
dim		1	1	2	1	1	1
T_1	1	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
		1	1	2	1	1	1	Bipartite graph
v^1	0	0	1	0	0	2		$H_{3/2}^1$
v^2	0	0	1	0	1	1		$H_{3/2}^1$
v^3	0	0	1	0	2	0		$H_{3/2}^1$
v^4	0	0	1	1	0	1		$H_{3/2}^1$
v^5	0	0	1	1	1	0		$H_{3/2}^1$
v^6	0	0	1	2	0	0		$H_{3/2}^1$
v^7	0	1	0	0	0	1		$H_{2/1}^1$
v^8	0	1	0	0	1	0		$H_{2/1}^1$
v^9	0	1	0	1	0	0		$H_{2/1}^1$
v^{10}	1	0	0	0	0	1		$H_{2/1}^1$
v^{11}	1	0	0	0	1	0		$H_{2/1}^1$
v^{12}	1	0	0	1	0	0		$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	M_1	M_2
	dim	1	1	2	1	1
Σ_1	1	1	0	1	-1	0
Σ_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

 U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	L_1L_2
	dim	1	1	2
U_1	1	1	0	-1
U_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_1	M_2	L_1L_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

 S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3A_3	H_1H_1	H_2H_2	H_3H_3
	dim	2	4	2	2	2
S_1S_1	2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1A_2	A_3A_3	H_1H_1	H_2H_2	H_3H_3	
	2	4	2	2	2	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3A_3	L_1L_2
	dim	2	4	2
R_1	1	1	2	-1
R_2	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_2	A_3A_3	L_1L_2	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_1L_2	H_1H_1	H_2H_2	H_3H_3
	dim	2	2	2	2
$S'_1S'_1$	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1L_2	H_1H_1	H_2H_2	H_3H_3	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1H_1	H_2H_2	H_3H_3
	dim	1	1	1	2	2	2
P_1	1	1	0	0	-2	0	0
P_2	1	0	1	0	0	-2	0
P_3	1	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1H_1	H_2H_2	H_3H_3	
	1	1	1	2	2	2	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	2	0	0	1	0	$H_{3/2}^1$
v^3	2	0	0	1	0	0	$H_{3/2}^1$

SG 159

Space group $P31c$, No. 159, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0)$, $A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	A_1A_2	A_3A_3
	dim	1	1	2	2	4
Δ_1	1	1	0	0	-1	0
Δ_2	1	0	1	0	-1	0
Δ_3	2	0	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	Γ_2	Γ_3	A_1A_2	A_3A_3	
	1	1	2	2	4	Bipartite graph
v^1	0	0	2	0	1	$H_{3/2}^1$
v^2	1	1	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	K_1	K_2	K_3
	dim	1	1	2	1	1	2
T_1	1	1	0	1	-1	0	-1
T_2	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	K_1	K_2	K_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	M_1	M_2
	dim	1	1	2	1	1
Σ_1	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	M_1	M_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$

v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	L_1L_2
	dim	1	1	2
U_1	1	1	0	-1
U_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_1	M_2	L_1L_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1	M_2	K_1	K_2	K_3
	dim	1	1	1	1	2
T'_1	1	1	0	-1	0	-1
T'_2	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_1	M_2	K_1	K_2	K_3	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3A_3	H_1	H_2	H_3
	dim	2	4	1	1	2
S_1	1	1	2	0	-1	-1
S_2	1	1	2	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1A_2	A_3A_3	H_1	H_2	H_3	
	2	4	1	1	2	Bipartite graph
v^1	0	1	0	0	2	$H_{3/4}^1$
v^2	0	1	1	1	1	$H_{4/4}^1$
v^3	0	1	2	2	0	$H_{5/4}^1$
v^4	1	0	0	0	1	$H_{2/2}^1$
v^5	1	0	1	1	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3A_3	L_1L_2
	dim	2	4	2
R_1R_1	2	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_2	A_3A_3	L_1L_2	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_1L_2	H_1	H_2	H_3
	dim	2	1	1	2
S'_1	1	1	0	-1	-1
S'_2	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$L_1 L_2$	H_1	H_2	H_3	
	2	1	1	2	Bipartite graph
v^1	1	0	0	1	$H_{2/2}^1$
v^2	1	1	1	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	0	0	0	-1	0
P_2	1	0	1	0	-1	0	0
P_3	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	1	0	$H_{2/1}^1$

SG 160

Space group $R3m$, No. 160, without SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	F_1	F_2
	dim	1	1	2	1	1
Σ_1	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	F_1	F_2	
	1	1	2	1	1	Bipartite graph

v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y: $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$
A matrix

corep		Z_1	Z_2	Z_3	L_1	L_2
	dim	1	1	2	1	1
Y_1	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_1	Z_2	Z_3	L_1	L_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

SG 161

Space group $R3c$, No. 161, without SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	F_1	F_2
	dim	1	1	2	1	1
Σ_1	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	F_1	F_2	
	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y: $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep	Z_1Z_2	Z_3Z_3	L_1L_2
dim	2	4	2
Y_1Y_1	2	1	2
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_1Z_2	Z_3Z_3	L_1L_2	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 162

Space group $P\bar{3}1m$, No. 162, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-
dim	1	1	1	1	2	2	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	-1	0	0	-1	0
Δ_2	1	0	1	1	0	0	0	0	-1	-1	0	0
Δ_3	2	0	0	0	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	1	0	0	1	1	-1	-1	0	0
Σ_2	1	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	0	0	1	-1	0	-1
T'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	H_1	H_2	H_3
	dim	1	1	1	1	2	2	1	1	2
S_1	1	1	0	0	1	1	1	-1	0	-1
S_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	H_1	H_2	H_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	L_1^+	L_1^-	L_2^+	L_2^-
dim		1	1	1	1	2	2	1	1	1	1
R_1	1	1	1	0	0	1	1	-1	-1	0	0
R_2	1	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	L_1^+	L_1^-	L_2^+	L_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_1^+	L_1^-	L_2^+	L_2^-	H_1	H_2	H_3
dim		1	1	1	1	1	1	2
S'_1	1	1	0	0	1	-1	0	-1
S'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	L_1^+	L_1^-	L_2^+	L_2^-	H_1	H_2	H_3	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$

v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	0	0	-1	0	0
P_2	1	0	1	0	0	-1	0
P_3	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

SG 163

Space group $P\bar{3}1c$, No. 163, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0)$, $A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	A_1A_2	A_3
	dim	1	1	1	1	2	2	4	2
Δ_1	1	1	0	0	1	0	0	0	-1
Δ_2	1	0	1	1	0	0	0	0	-1
Δ_3	2	0	0	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	$A_1 A_2$	A_3	
	1	1	1	1	2	2	4	2	Bipartite graph
v^1	0	0	0	0	0	2	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	1	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	1	0	0	1	1	-1	-1	0	0
Σ_2	1	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	L_1
	dim	1	1	1	1	2
U_1	1	1	0	0	1	-1
U_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1^+	M_1^-	M_2^+	M_2^-	L_1	Bipartite graph
	1	1	1	1	2	
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3
dim		1	1	1	1	1	1	2
T'_1	1	1	0	0	1	-1	0	-1
T'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1A_2	A_3	H_1	H_2	H_3
dim		4	2	1	1	2
S_1	1	2	1	0	-1	-1
S_2	1	2	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1A_2	A_3	H_1	H_2	H_3	Bipartite graph
	4	2	1	1	2	
v^1	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	1	1	0	$H_{3/2}^1$
v^3	1	0	0	0	2	$H_{3/4}^1$
v^4	1	0	1	1	1	$H_{4/4}^1$
v^5	1	0	2	2	0	$H_{5/4}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$A_1 A_2$	A_3	L_1
dim	4	2	2
$R_1 R_2$	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$A_1 A_2$	A_3	L_1	
	4	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	2	$H_{3/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	H_1	H_2	H_3
dim	2	1	1	2
S'_1	1	0	-1	-1
S'_2	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1	H_1	H_2	H_3	
	2	1	1	2	Bipartite graph
v^1	1	0	0	1	$H_{2/2}^1$
v^2	1	1	1	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	K_1	K_2	K_3	H_1	H_2	H_3
dim	1	1	2	1	1	2
P_1	1	0	0	0	-1	0
P_2	1	0	1	-1	0	0
P_3	2	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	1	0	$H_{2/1}^1$

SG 164

Space group $P\bar{3}m1$, No. 164, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-
dim		1	1	1	1	2	2	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	-1	0	0	-1	0	0
Δ_2	1	0	1	1	0	0	0	0	-1	-1	0	0	0
Δ_3	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	
	1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	1	0	0	1	1	-1	0	-1
T_2	1	0	0	1	1	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	0	1	1	1	-1	0	0	-1
Σ_2	1	0	1	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-	
	1	1	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$

v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	1	0	0	-1	0	-1
T'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	H_1	H_2	H_3
	dim	1	1	1	1	2	2	1	1	2
S_1	1	1	1	0	0	1	1	-1	0	-1
S_2	1	0	0	1	1	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	H_1	H_2	H_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	L_1^+	L_1^-	L_2^+	L_2^-
dim		1	1	1	1	2	2	1	1	1	1
R_1	1	1	0	0	1	1	1	-1	0	0	-1
R_2	1	0	1	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	L_1^+	L_1^-	L_2^+	L_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$

v^{14}	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1^+	L_1^-	L_2^+	L_2^-	H_1	H_2	H_3
	dim	1	1	1	1	1	1	2
S'_1	1	1	1	0	0	-1	0	-1
S'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	L_1^+	L_1^-	L_2^+	L_2^-	H_1	H_2	H_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	1	0	-1	-1	0
P_2P_3	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

K_1	K_2	K_3	H_1	H_2	H_3	
1	1	2	1	1	2	Bipartite graph

v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

SG 165

Space group $P\bar{3}c1$, No. 165, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	A_1A_2	A_3
	dim	1	1	1	1	2	2	4	2
Δ_1	1	1	0	0	1	0	0	0	-1
Δ_2	1	0	1	1	0	0	0	0	-1
Δ_3	2	0	0	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	A_1A_2	A_3	Bipartite graph
	1	1	1	1	2	2	4	2	
v^1	0	0	0	0	0	2	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	1	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	1	0	0	1	1	-1	0	-1
T_2	1	0	0	1	1	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-
dim		1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	0	1	1	1	-1	0	0	-1
Σ_2	1	0	1	1	0	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	M_1^+	M_1^-	M_2^+	M_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$

v^{14}	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	L_1
	dim	1	1	1	1	2
U_1	1	1	0	0	1	-1
U_2	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1^+	M_1^-	M_2^+	M_2^-	L_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	1	0	0	-1	0	-1
T'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3	H_1H_2	H_3H_3
dim	4	2	2	4
S_1S_2	2	2	1	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_1A_2	A_3	H_1H_2	H_3H_3	Bipartite graph
	4	2	2	4	
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	2	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3	L_1
dim	4	2	2
R_1	1	2	1 -1
R_2	1	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_2	A_3	L_1	Bipartite graph
	4	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	2	$H_{3/4}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	H_1H_2	H_3H_3
dim	2	2	4
$S'_1S'_2$	2	1	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1	H_1H_2	H_3H_3	
	2	2	4	Bipartite graph
v^1	1	1	0	$H_{2/1}^1$
v^2	2	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1H_2	H_3H_3
dim		1	1	2	2	4
P_1	1	1	1	0	-2	0
P_2P_3	2	0	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_1	K_2	K_3	H_1H_2	H_3H_3	
	1	1	2	2	4	Bipartite graph
v^1	0	0	2	0	1	$H_{3/2}^1$
v^2	0	2	0	1	0	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	2	0	0	1	0	$H_{3/2}^1$

SG 166

Space group $R\bar{3}m$, No. 166, without SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	F_1^+	F_1^-	F_2^+	F_2^-
dim		1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	1	0	0	1	1	-1	-1	0	0
Σ_2	1	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	F_1^+	F_1^-	F_2^+	F_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y: $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep		Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	L_1^+	L_1^-	L_2^+	L_2^-
dim		1	1	1	1	2	2	1	1	1	1
Y_1	1	1	1	0	0	1	1	-1	-1	0	0
Y_2	1	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_1^+	Z_1^-	Z_2^+	Z_2^-	Z_3^+	Z_3^-	L_1^+	L_1^-	L_2^+	L_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$

v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 167

Space group $R\bar{3}c$, No. 167, without SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	F_1^+	F_1^-	F_2^+	F_2^-
dim		1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	1	0	0	1	1	-1	-1	0	0
Σ_2	1	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	F_1^+	F_1^-	F_2^+	F_2^-	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y : $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$
A matrix

corep		$Z_1 Z_2$	Z_3	L_1
dim	2	4	2	2
$Y_1 Y_2$	2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$Z_1 Z_2$	Z_3	L_1	Bipartite graph
	4	2	2	
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	2	$H_{3/2}^1$

SG 168

Space group $P6$, No. 168, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	$\Gamma_2 \Gamma_3$	Γ_4	$\Gamma_5 \Gamma_6$	A_1	$A_2 A_3$	A_4	$A_5 A_6$
dim	1	1	2	1	2	1	2	1	2
Δ_1	1	1	0	0	0	-1	0	0	0
Δ_2	1	0	1	0	0	0	-1	0	0
Δ_3	1	0	1	0	0	0	-1	0	0
Δ_4	1	0	0	1	0	0	0	-1	0
Δ_5	1	0	0	0	1	0	0	0	-1
Δ_6	1	0	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	$\Gamma_2 \Gamma_3$	Γ_4	$\Gamma_5 \Gamma_6$	A_1	$A_2 A_3$	A_4	$A_5 A_6$	Bipartite graph
	1	2	1	2	1	2	1	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	0	0	$H_{2/2}^1$
v^4	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3
dim	1	2	1	2	1	2
T_1	1	2	1	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3	Bipartite graph
	1	2	1	2	1	2	
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2
dim	1	2	1	2	1	1
Σ_1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2	Bipartite graph
	1	2	1	2	1	1	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_1	M_2	K_1	K_2K_3
dim	1	1	1	2
T'_1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	K_1	K_2K_3	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1	A_2A_3	A_4	A_5A_6	H_1	H_2H_3
dim	1	2	1	2	1	2
S_1	1	1	2	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1	A_2A_3	A_4	A_5A_6	H_1	H_2H_3	
	1	2	1	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_3	A_4	A_5A_6	L_1	L_2
	dim	1	2	1	2	1	1
R_1	1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1	A_2A_3	A_4	A_5A_6	L_1	L_2	
	1	2	1	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	H_1	H_2H_3
	dim	1	1	1	2
S'_1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1	L_2	H_1	H_2H_3	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_1	K_2K_3	H_1	H_2H_3
	dim	1	2	1	2
P_1	1	1	0	-1	0
P_2	1	0	1	0	-1
P_3	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_1	K_2K_3	H_1	H_2H_3	
	1	2	1	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 169

Space group $P6_1$, No. 169, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1A_2	A_3A_6	A_4A_5
	dim	1	2	1	2	2	2	2
Δ_1	1	1	0	0	0	0	-1	0
Δ_2	1	0	1	0	0	-1	0	0
Δ_3	1	0	1	0	0	0	-1	0
Δ_4	1	0	0	1	0	0	0	-1
Δ_5	1	0	0	0	1	-1	0	0
Δ_6	1	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1A_2	A_3A_6	A_4A_5	
	1	2	1	2	2	2	2	Bipartite graph
v^1	1	1	1	1	1	1	1	$H_{7/6}^3$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3
dim	1	2	1	2	1	2
T_1	1	2	1	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3	Bipartite graph
	1	2	1	2	1	2	
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2
dim	1	2	1	2	1	1
Σ_1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2	Bipartite graph
	1	2	1	2	1	1	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	$L_1 L_2$
dim		1	1	2
U_1	1	1	0	-1
U_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_1	M_2	$L_1 L_2$	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1	M_2	K_1	$K_2 K_3$
dim		1	1	1	2
T'_1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	K_1	$K_2 K_3$	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3A_6	A_4A_5	H_1H_1	H_2H_3
dim	2	2	2	2	2
S_1S_1	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1A_2	A_3A_6	A_4A_5	H_1H_1	H_2H_3	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3A_6	A_4A_5	L_1L_2
dim	2	2	2	2
R_1R_1	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_6	A_4A_5	L_1L_2	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1L_2	H_1H_1	H_2H_3
dim	2	2	2
$S'_1S'_1$	2	1	-1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1L_2	H_1H_1	H_2H_3	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2K_3	H_1H_1	H_2H_3
	dim	1	2	2	2
P_1	1	1	0	0	-1
P_2	1	0	1	0	-1
P_3	1	0	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	K_1	K_2K_3	H_1H_1	H_2H_3	
	1	2	2	2	Bipartite graph
v^1	2	2	1	2	$H_{7/6}^3$

SG 170

Space group $P6_5$, No. 170, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1A_2	A_3A_6	A_4A_5
	dim	1	2	1	2	2	2	2
Δ_1	1	1	0	0	0	0	0	-1
Δ_2	1	0	1	0	0	0	0	-1
Δ_3	1	0	1	0	0	-1	0	0
Δ_4	1	0	0	1	0	0	-1	0
Δ_5	1	0	0	0	1	0	-1	0
Δ_6	1	0	0	0	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1A_2	A_3A_6	A_4A_5	
	1	2	1	2	2	2	2	Bipartite graph
v^1	1	1	1	1	1	1	1	$H_{7/6}^3$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3
dim	1	2	1	2	1	2
T_1	1	1	2	1	2	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3	
	1	2	1	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2
dim	1	2	1	2	1	1
Σ_1	1	1	2	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2	
	1	2	1	2	1	1	Bipartite graph

v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	L_1L_2
	dim	1	1	2
U_1	1	1	0	-1
U_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_1	M_2	L_1L_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	K_1	K_2K_3
	dim	1	1	1	2
T'_1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	K_1	K_2K_3	
	1	1	1	2	Bipartite graph

v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3A_6	A_4A_5	H_1H_1	H_2H_3
dim	2	2	2	2	2
S_1S_1	2	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1A_2	A_3A_6	A_4A_5	H_1H_1	H_2H_3	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3A_6	A_4A_5	L_1L_2
dim	2	2	2	2
R_1R_1	2	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_6	A_4A_5	L_1L_2	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1L_2	H_1H_1	H_2H_3
dim		2	2	2
$S'_1S'_1$	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1L_2	H_1H_1	H_2H_3	Bipartite graph
	2	2	2	
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2K_3	H_1H_1	H_2H_3
dim		1	2	2	2
P_1	1	1	0	0	-1
P_2	1	0	1	-2	0
P_3	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	K_1	K_2K_3	H_1H_1	H_2H_3	Bipartite graph
	1	2	2	2	
v^1	2	2	1	2	$H_{7/6}^3$

SG 171

Space group $P6_2$, No. 171, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1	A_2A_3	A_4	A_5A_6
dim	1	2	1	2	1	2	1	2
Δ_1	1	1	0	0	0	-1	0	0
Δ_2	1	0	1	0	0	0	0	-1
Δ_3	1	0	1	0	0	-1	0	0
Δ_4	1	0	0	1	0	0	0	-1
Δ_5	1	0	0	0	1	0	-1	0
Δ_6	1	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1	A_2A_3	A_4	A_5A_6	
	1	2	1	2	1	2	1	2	Bipartite graph
v^1	0	1	1	0	1	0	0	1	$H_{4/3}^2$
v^2	1	0	0	1	0	1	1	0	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3
dim	1	2	1	2	1	2
T_1	1	1	2	1	2	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3	
	1	2	1	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2
	dim	1	2	1	2	1	1
Σ_1	1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2	
	1	2	1	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	K_1	K_2K_3
	dim	1	1	1	2
T'_1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	K_1	K_2K_3	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_3	A_4	A_5A_6	H_1	H_2H_3
	dim	1	2	1	2	1	2
S_1	1	1	2	1	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1	A_2A_3	A_4	A_5A_6	H_1	H_2H_3	
	1	2	1	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_3	A_4	A_5A_6	L_1	L_2
	dim	1	2	1	2	1	1
R_1	1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1	A_2A_3	A_4	A_5A_6	L_1	L_2	
	1	2	1	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	H_1	H_2H_3
dim		1	1	1	2
S'_1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1	L_2	H_1	H_2H_3	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2K_3	H_1	H_2H_3
dim		1	2	1	2
P_1	1	1	0	0	-1
P_2	1	0	1	-1	0
P_3	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	K_1	K_2K_3	H_1	H_2H_3	
	1	2	1	2	Bipartite graph
v^1	1	1	1	1	$H_{4/3}^2$

SG 172

Space group $P6_4$, No. 172, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1	A_2A_3	A_4	A_5A_6
dim	1	2	1	2	1	2	1	2
Δ_1	1	1	0	0	0	0	0	-1
Δ_2	1	0	1	0	0	0	-1	0
Δ_3	1	0	1	0	0	-1	0	0
Δ_4	1	0	0	1	0	-1	0	0
Δ_5	1	0	0	0	1	-1	0	0
Δ_6	1	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1	A_2A_3	A_4	A_5A_6	
	1	2	1	2	1	2	1	2	Bipartite graph
v^1	0	1	1	0	0	1	1	0	$H_{4/3}^2$
v^2	1	0	0	1	1	0	0	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3
dim	1	2	1	2	1	2
T_1	1	1	2	1	2	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3	
	1	2	1	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2
	dim	1	2	1	2	1	1
Σ_1	1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2	
	1	2	1	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	K_1	K_2K_3
	dim	1	1	1	2
T'_1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	K_1	K_2K_3	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_3	A_4	A_5A_6	H_1	H_2H_3
	dim	1	2	1	2	1	2
S_1	1	1	2	1	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1	A_2A_3	A_4	A_5A_6	H_1	H_2H_3	
	1	2	1	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2A_3	A_4	A_5A_6	L_1	L_2
	dim	1	2	1	2	1	1
R_1	1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1	A_2A_3	A_4	A_5A_6	L_1	L_2	
	1	2	1	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_1	L_2	H_1	H_2H_3
dim	1	1	1	2
S'_1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1	L_2	H_1	H_2H_3	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_1	K_2K_3	H_1	H_2H_3
dim	1	2	1	2
P_1	1	0	0	-1
P_2	1	0	0	-1
P_3	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	K_1	K_2K_3	H_1	H_2H_3	
	1	2	1	2	Bipartite graph
v^1	1	1	1	1	$H_{4/3}^2$

SG 173

Space group $P6_3$, No. 173, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1A_2	A_3A_6	A_4A_5
dim	1	2	1	2	2	2	2
Δ_1	1	1	0	0	0	-1	0
Δ_2	1	0	1	0	0	0	-1
Δ_3	1	0	1	0	0	0	-1
Δ_4	1	0	0	1	0	-1	0
Δ_5	1	0	0	0	1	0	-1
Δ_6	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1A_2	A_3A_6	A_4A_5	
	1	2	1	2	2	2	2	Bipartite graph
v^1	0	1	0	1	0	1	1	$H_{4/4}^2$
v^2	1	0	1	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3
dim	1	2	1	2	1	2
T_1	1	1	2	1	2	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2K_3	
	1	2	1	2	1	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ: $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2
	dim	1	2	1	2	1	1
Σ_1	1	1	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2	
	1	2	1	2	1	1	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	L_1L_2
	dim	1	1	2
U_1	1	1	0	-1
U_2	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_1	M_2	L_1L_2	
	1	1	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_1	M_2	K_1	K_2K_3
dim	1	1	1	2
T'_1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_1	M_2	K_1	K_2K_3	
	1	1	1	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3A_6	A_4A_5	H_1H_1	H_2H_3
dim	2	2	2	2	2
S_1S_1	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1A_2	A_3A_6	A_4A_5	H_1H_1	H_2H_3	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3A_6	A_4A_5	L_1L_2
dim	2	2	2	2
R_1R_1	2	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_2	A_3A_6	A_4A_5	L_1L_2	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1L_2	H_1H_1	H_2H_3
dim	2	2	2
$S'_1S'_1$	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1L_2	H_1H_1	H_2H_3	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	K_1	K_2K_3	H_1H_1	H_2H_3
dim	1	2	2	2
P_1	1	1	0	-2 0
P_2	1	0	1	0 -1
P_3	1	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_1	K_2K_3	H_1H_1	H_2H_3	
	1	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	2	0	1	0	$H_{3/2}^1$

SG 174

Space group $P\bar{6}$, No. 174, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1	A_2A_3	A_4	A_5A_6
	dim	1	2	1	2	1	2	1	2
Δ_1	1	1	0	1	0	-1	0	-1	0
$\Delta_2\Delta_3$	2	0	1	0	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	A_1	A_2A_3	A_4	A_5A_6	Bipartite graph
	1	2	1	2	1	2	1	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	2	1	2	1	1	1	1	1	1
T_1	1	1	2	0	0	-1	-1	-1	0	0	0
T_2	1	0	0	1	2	0	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	2	1	2	1	1	1	1	1	1	
v^1	0	0	0	1	0	0	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$

v^8	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	2	0	0	0	0	$H_{3/2}^1$
v^{13}	0	1	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{14}	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{15}	0	1	0	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{18}	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2
	dim	1	2	1	2	1	1
Σ_1	1	1	2	0	0	-1	0
Σ_2	1	0	0	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	$\Gamma_5\Gamma_6$	M_1	M_2	Bipartite graph
	1	2	1	2	1	1	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	1	0	0	2	0	$H_{3/2}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1	A_2A_3	A_4	A_5A_6	H_1	H_2	H_3	H_4	H_5	H_6
	dim	1	2	1	2	1	1	1	1	1	1
S_1	1	1	2	0	0	-1	-1	-1	0	0	0
S_2	1	0	0	1	2	0	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	A_1	A_2A_3	A_4	A_5A_6	H_1	H_2	H_3	H_4	H_5	H_6	Bipartite graph
	1	2	1	2	1	1	1	1	1	1	
v^1	0	0	0	1	0	0	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	2	0	0	0	0	$H_{3/2}^1$
v^{13}	0	1	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{14}	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{15}	0	1	0	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{18}	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2A_3	A_4	A_5A_6	L_1	L_2
	dim	1	2	1	2	1	1
R_1	1	1	2	0	0	-1	0
R_2	1	0	0	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_1	A_2A_3	A_4	A_5A_6	L_1	L_2	Bipartite graph
	1	2	1	2	1	1	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	1	0	0	2	0	$H_{3/2}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$

SG 175

Space group $P6/m$, No. 175, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$
dim	1	1	2	2	1	1	2	2	1	1	2	2	1	1	2	2
Δ_1	1	1	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0
$\Delta_2\Delta_3$	2	0	0	1	1	0	0	0	0	0	-1	-1	0	0	0	0
Δ_4	1	0	0	0	0	1	1	0	0	0	0	0	-1	-1	0	0
$\Delta_5\Delta_6$	2	0	0	0	0	0	0	1	1	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	Bipartite graph
	1	1	2	2	1	1	2	2	1	1	2	2	1	1	2	2	
v^1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_1	K_2K_3	K_4	K_5K_6
dim	1	1	2	2	1	1	2	2	1	2	1	2
T_1	1	1	0	0	2	0	1	2	0	-1	-2	0
T_2	1	0	1	2	0	1	0	0	2	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_1	K_2K_3	K_4	K_5K_6	Bipartite graph
	1	1	2	2	1	1	2	2	1	2	1	2	
v^1	0	0	0	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	0	2	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^{10}	0	0	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{11}	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{12}	0	0	1	0	0	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{13}	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	2	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{18}	2	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	M_1^+	M_1^-	M_2^+	M_2^-
	dim	1	1	2	2	1	1	2	2	1	1	1	1
Σ_1	1	1	0	0	2	0	1	2	0	-1	0	0	-1
Σ_2	1	0	1	2	0	1	0	0	2	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	M_1^+	M_1^-	M_2^+	M_2^-	Bipartite graph
	1	1	2	2	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$

v^8	0	0	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	1	0	0	0	0	0	0	0	2	$H_{3/2}^1$
v^{12}	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{15}	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{16}	0	0	1	0	0	0	0	0	0	2	0	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2K_3	K_4	K_5K_6
	dim	1	1	1	1	1	2	1	2
T'_1	1	1	0	0	1	-1	-2	0	0
T'_2	1	0	1	1	0	0	0	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2K_3	K_4	K_5K_6	Bipartite graph
	1	1	1	1	1	2	1	2	
v^1	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^2	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{10}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	H_1	H_2H_3	H_4	H_5H_6
	dim	1	1	2	2	1	1	2	2	1	2	1	2
S_1	1	1	0	0	2	0	1	2	0	-1	-2	0	0
S_2	1	0	1	2	0	1	0	0	2	0	0	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	H_1	H_2H_3	H_4	H_5H_6	Bipartite graph
	1	1	2	2	1	1	2	2	1	2	1	2	
v^1	0	0	0	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	0	2	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^{10}	0	0	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{11}	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{12}	0	0	1	0	0	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{13}	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	2	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{18}	2	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	L_1^+	L_1^-	L_2^+	L_2^-
	dim	1	1	2	2	1	1	2	2	1	1	1	1
R_1	1	1	0	0	2	0	1	2	0	-1	0	0	-1
R_2	1	0	1	2	0	1	0	0	2	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	A_1^+	A_1^-	$A_2^+A_3^+$	$A_2^-A_3^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	L_1^+	L_1^-	L_2^+	L_2^-	Bipartite graph
	1	1	2	2	1	1	2	2	1	1	1	1	

v^1	0	0	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	1	0	0	0	0	0	0	0	2	$H_{3/2}^1$
v^{12}	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{15}	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{16}	0	0	1	0	0	0	0	0	0	2	0	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1^+	L_1^-	L_2^+	L_2^-	H_1	H_2H_3	H_4	H_5H_6
	dim	1	1	1	1	1	2	1	2
S'_1	1	1	0	0	1	-1	-2	0	0
S'_2	1	0	1	1	0	0	0	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	L_1^+	L_1^-	L_2^+	L_2^-	H_1	H_2H_3	H_4	H_5H_6	
	1	1	1	1	1	2	1	2	Bipartite graph
v^1	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^2	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	0	0	1	$H_{3/2}^1$

v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{10}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2K_3	K_4	K_5K_6	H_1	H_2H_3	H_4	H_5H_6
	dim	1	2	1	2	1	2	1	2
P_1	1	1	0	1	0	-1	0	-1	0
P_2P_3	2	0	1	0	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	K_1	K_2K_3	K_4	K_5K_6	H_1	H_2H_3	H_4	H_5H_6	Bipartite graph
	1	2	1	2	1	2	1	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 176

Space group $P6_3/m$, No. 176, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_1A_3	A_2
	dim	1	1	2	2	1	1	2	2	4	2
Δ_1	1	1	1	0	0	0	0	0	0	0	-1
$\Delta_2\Delta_3$	2	0	0	1	1	0	0	0	0	-1	0
Δ_4	1	0	0	0	0	1	1	0	0	0	-1
$\Delta_5\Delta_6$	2	0	0	0	0	0	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_1A_3	A_2	
	1	1	2	2	1	1	2	2	4	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_1	K_2K_3	K_4	K_5K_6
dim	1	1	2	2	1	1	2	2	1	2	1	2
T_1	1	0	0	2	0	1	2	0	-1	-2	0	0
T_2	1	0	1	2	0	1	0	2	0	0	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_1	K_2K_3	K_4	K_5K_6	
	1	1	2	2	1	1	2	2	1	2	1	2	Bipartite graph
v^1	0	0	0	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	0	2	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^{10}	0	0	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{11}	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{12}	0	0	1	0	0	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{13}	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	2	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{18}	2	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	M_1^+	M_1^-	M_2^+	M_2^-
dim		1	1	2	2	1	1	2	2	1	1	1	1
Σ_1	1	1	0	0	2	0	1	2	0	-1	0	0	-1
Σ_2	1	0	1	2	0	1	0	0	2	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	M_1^+	M_1^-	M_2^+	M_2^-	Bipartite graph
	1	1	2	2	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	1	0	0	0	0	0	0	0	2	$H_{3/2}^1$
v^{12}	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{15}	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{16}	0	0	1	0	0	0	0	0	0	2	0	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	L_1
	dim	1	1	1	1	2
U_1	1	1	1	0	0	-1
U_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1^+	M_1^-	M_2^+	M_2^-	L_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2K_3	K_4	K_5K_6
	dim	1	1	1	1	1	2	1	2
T'_1	1	1	0	0	1	-1	-2	0	0
T'_2	1	0	1	1	0	0	0	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	M_1^+	M_1^-	M_2^+	M_2^-	K_1	K_2K_3	K_4	K_5K_6	
	1	1	1	1	1	2	1	2	Bipartite graph
v^1	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^2	0	0	0	2	0	1	0	0	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{10}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1A_3	A_2	H_1H_4	H_2H_6	H_3H_5
dim	4	2	2	2	2
S_1S_2	2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1A_3	A_2	H_1H_4	H_2H_6	H_3H_5	Bipartite graph
	4	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	2	$H_{3/2}^1$
v^5	1	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	1	0	1	$H_{3/2}^1$
v^8	1	0	1	1	0	$H_{3/2}^1$
v^9	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_3	A_2	L_1
dim	4	2	2
R_1R_2	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_3	A_2	L_1	Bipartite graph
	4	2	2	
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	2	$H_{3/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	H_1H_4	H_2H_6	H_3H_5
dim	2	2	2	2
$S'_1S'_2$	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	H_1H_4	H_2H_6	H_3H_5	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_1	K_2K_3	K_4	K_5K_6	H_1H_4	H_2H_6	H_3H_5
	dim	1	2	1	2	2	2	2
P_1	1	1	0	1	0	-2	0	0
P_2P_3	2	0	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	K_1	K_2K_3	K_4	K_5K_6	H_1H_4	H_2H_6	H_3H_5	
	1	2	1	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	1	0	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	1	0	0	$H_{3/2}^1$
v^7	2	0	0	0	1	0	0	$H_{3/2}^1$

SG 177

Space group *P622*, No. 177, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
Δ_1	1	1	1	0	0	0	0	-1	-1	0	0	0	0
Δ_2	1	0	0	0	0	0	1	0	0	0	0	0	-1
Δ_3	1	0	0	0	0	0	1	0	0	0	0	0	-1
Δ_4	1	0	0	1	1	0	0	0	0	-1	-1	0	0

Δ_5	1	0	0	0	0	1	0	0	0	0	0	-1	0
Δ_6	1	0	0	0	0	1	0	0	0	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	0	0	-1
Σ_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	1	0	0	-1	0	-1
T'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3
	dim	1	1	1	1	2	2	1	1	2
S_1	1	1	0	0	1	1	1	-1	0	-1
S_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4
	dim	1	1	1	1	2	2	1	1	1	1
R_1	1	1	0	1	0	1	1	-1	0	0	-1
R_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_1	L_2	L_3	L_4	H_1	H_2	H_3
	dim	1	1	1	1	1	1	2
S'_1	1	1	1	0	0	-1	0	-1
S'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	L_1	L_2	L_3	L_4	H_1	H_2	H_3	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$

v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	1	0	-1	-1	0
P_2	1	0	0	1	0	0	-1
P_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

SG 178

Space group $P6_122$, No. 178, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3
	dim	1	1	1	1	2	2	2	2	2
Δ_1	1	1	1	0	0	0	0	-1	0	0
Δ_2	1	0	0	0	0	0	1	0	0	-1
Δ_3	1	0	0	0	0	0	1	-1	0	0
Δ_4	1	0	0	1	1	0	0	0	-1	0
Δ_5	1	0	0	0	0	1	0	0	0	-1
Δ_6	1	0	0	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	
	1	1	1	1	2	2	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	1	1	1	$H_{7/6}^3$
v^2	0	1	1	0	1	1	1	1	1	$H_{7/6}^3$
v^3	1	0	0	1	1	1	1	1	1	$H_{7/6}^3$
v^4	1	0	1	0	1	1	1	1	1	$H_{7/6}^3$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	0	0	-1
Σ_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	L_1
	dim	1	1	1	1	2
U_1	1	1	0	1	0	-1
U_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	L_1	Bipartite graph
	1	1	1	1	2	
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
dim		1	1	1	1	1	1	2
T'_1	1	1	1	0	0	-1	0	-1
T'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	$H_1 H_2$	H_3
dim		2	2	2	2	2
$S_1 S_2$	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1	A_2	A_3	$H_1 H_2$	H_3	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	L_1
dim	2	2	2	2
$R_1 R_2$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_1	$H_1 H_2$	H_3
dim	2	2	2
$S'_1 S'_2$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1	$H_1 H_2$	H_3	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_1	K_2	K_3	$H_1 H_2$	H_3
dim	1	1	2	2	2
P_1	1	1	0	0	-1
P_2	1	0	0	0	-1
P_3	1	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1H_2	H_3	
	1	1	2	2	2	Bipartite graph
v^1	0	2	2	1	2	$H_{7/6}^3$
v^2	1	1	2	1	2	$H_{7/6}^3$
v^3	2	0	2	1	2	$H_{7/6}^3$

SG 179

Space group $P6_522$, No. 179, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3
	dim	1	1	1	1	2	2	2	2	2
Δ_1	1	1	1	0	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	0	-1	0
Δ_3	1	0	0	0	0	0	1	0	0	-1
Δ_4	1	0	0	1	1	0	0	-1	0	0
Δ_5	1	0	0	0	0	1	0	-1	0	0
Δ_6	1	0	0	0	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	
	1	1	1	1	2	2	2	2	2	Bipartite graph
v^1	0	1	0	1	1	1	1	1	1	$H_{7/6}^3$
v^2	0	1	1	0	1	1	1	1	1	$H_{7/6}^3$
v^3	1	0	0	1	1	1	1	1	1	$H_{7/6}^3$
v^4	1	0	1	0	1	1	1	1	1	$H_{7/6}^3$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
dim		1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	0	0	-1
Σ_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	
	1	1	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$

v^{14}	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_1	M_2	M_3	M_4	L_1
dim	1	1	1	1	2
U_1	1	0	1	0	-1
U_2	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	L_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_1	M_2	M_3	M_4	K_1	K_2	K_3
dim	1	1	1	1	1	1	2
T'_1	1	1	1	0	0	-1	-1
T'_2	1	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	$H_1 H_2$	H_3
dim	2	2	2	2	2
$S_1 S_2$	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1	A_2	A_3	$H_1 H_2$	H_3	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	L_1
dim	2	2	2	2
$R_1 R_2$	2	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	$H_1 H_2$	H_3
dim	2	2	2
$S'_1 S'_2$	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1	$H_1 H_2$	H_3	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	$H_1 H_2$	H_3
dim		1	1	2	2	2
P_1	1	1	1	0	0	-1
P_2	1	0	0	1	-2	0
P_3	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	$H_1 H_2$	H_3	
	1	1	2	2	2	Bipartite graph
v^1	0	2	2	1	2	$H_{7/6}^3$
v^2	1	1	2	1	2	$H_{7/6}^3$
v^3	2	0	2	1	2	$H_{7/6}^3$

SG 180

Space group $P6_222$, No. 180, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6
dim	1	1	1	1	2	2	1	1	1	1	2	2

Δ_1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	-1	
Δ_2	1	0	0	0	0	0	0	0	1	0	0	0	0	0	-1	0
Δ_3	1	0	0	0	0	0	0	1	-1	-1	0	0	0	0	0	
Δ_4	1	0	0	1	1	0	0	0	0	0	0	0	0	-1	0	
Δ_5	1	0	0	0	0	1	0	0	0	0	0	0	0	0	-1	
Δ_6	1	0	0	0	0	1	0	0	0	0	-1	-1	0	0	0	

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6	
	1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	1	0	1	0	0	1	0	$H_{4/3}^2$
v^2	0	0	0	1	0	1	1	0	0	0	1	0	$H_{4/3}^2$
v^3	0	0	1	0	0	1	0	1	0	0	1	0	$H_{4/3}^2$
v^4	0	0	1	0	0	1	1	0	0	0	1	0	$H_{4/3}^2$
v^5	0	1	0	0	1	0	0	0	0	1	0	1	$H_{4/3}^2$
v^6	0	1	0	0	1	0	0	0	1	0	0	1	$H_{4/3}^2$
v^7	1	0	0	0	1	0	0	0	0	1	0	1	$H_{4/3}^2$
v^8	1	0	0	0	1	0	0	0	1	0	0	1	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$

v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	0	0	-1
Σ_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	
	1	1	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	M_1	M_2	M_3	M_4	K_1	K_2	K_3
dim	1	1	1	1	1	1	2

T'_1	1	1	1	0	0	-1	0	-1
T'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3
dim		1	1	1	1	2	2	1	1	2
S_1	1	1	0	0	1	1	1	0	-1	-1
S_2	1	0	1	1	0	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4
	dim	1	1	1	1	2	2	1	1	1	1
R_1	1	1	0	1	0	1	1	-1	0	0	-1
R_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	L_3	L_4	H_1	H_2	H_3
	dim	1	1	1	1	1	1	2
S'_1	1	1	1	0	0	-1	0	-1
S'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	L_1	L_2	L_3	L_4	H_1	H_2	H_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	1	0	0	0	-1
P_2	1	0	0	1	-1	-1	0
P_3	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

SG 181

Space group $P6_422$, No. 181, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6
dim		1	1	1	1	2	2	1	1	1	1	2	2
Δ_1	1	1	1	0	0	0	0	0	0	0	0	-1	0
Δ_2	1	0	0	0	0	0	1	0	0	-1	-1	0	0
Δ_3	1	0	0	0	0	0	1	0	0	0	0	0	-1
Δ_4	1	0	0	1	1	0	0	0	0	0	0	0	-1
Δ_5	1	0	0	0	0	1	0	-1	-1	0	0	0	0
Δ_6	1	0	0	0	0	1	0	0	0	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	1	0	1	0	0	0	1	0	1	$H_{4/3}^2$
v^2	0	0	0	1	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^3	0	0	1	0	0	1	0	0	0	1	0	1	$H_{4/3}^2$
v^4	0	0	1	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^5	0	1	0	0	1	0	0	1	0	0	1	0	$H_{4/3}^2$
v^6	0	1	0	0	1	0	1	0	0	0	1	0	$H_{4/3}^2$
v^7	1	0	0	0	1	0	0	1	0	0	1	0	$H_{4/3}^2$
v^8	1	0	0	0	1	0	1	0	0	0	1	0	$H_{4/3}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
dim		1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$

v^8	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	0	0	-1
Σ_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	1	0	0	-1	0	-1
T'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3
	dim	1	1	1	1	2	2	1	1	2
S_1	1	1	0	0	1	1	1	-1	0	-1
S_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4
	dim	1	1	1	1	2	2	1	1	1	1
R_1	1	1	0	1	0	1	1	-1	0	0	-1
R_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_1	L_2	L_3	L_4	H_1	H_2	H_3
	dim	1	1	1	1	1	1	2
S'_1	1	1	1	0	0	-1	0	-1
S'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	L_1	L_2	L_3	L_4	H_1	H_2	H_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
	dim	1	1	2	1	1	2
P_1	1	1	1	0	0	0	-1
P_2	1	0	0	1	0	0	-1
P_3	1	0	0	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	1	1	0	1	1	$H_{4/3}^2$
v^2	0	1	1	1	0	1	$H_{4/3}^2$
v^3	1	0	1	0	1	1	$H_{4/3}^2$
v^4	1	0	1	1	0	1	$H_{4/3}^2$

SG 182

Space group $P6_322$, No. 182, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0)$, $A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3
	dim	1	1	1	1	2	2	2	2	2
Δ_1	1	1	1	0	0	0	0	0	0	-1
Δ_2	1	0	0	0	0	0	1	0	-1	0
Δ_3	1	0	0	0	0	0	1	-1	0	0
Δ_4	1	0	0	1	1	0	0	0	0	-1
Δ_5	1	0	0	0	0	1	0	-1	0	0
Δ_6	1	0	0	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	
	1	1	1	1	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	1	1	1	1	0	$H_{4/4}^2$
v^2	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

v^{11}	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	0	0	-1
Σ_2	1	0	1	0	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	L_1
	dim	1	1	1	1	2
U_1	1	1	0	1	0	-1
U_2	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	L_1	
	1	1	1	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	1	0	0	-1	0	-1
T'_2	1	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	H_1	H_2	H_3
	dim	2	2	2	2	2	
$S_1 S_2$	2	1	1	1	-1	-1	

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1	A_2	A_3	H_1H_2	H_3	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	L_1
dim	2	2	2	2
R_1R_2	2	1	1	1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_1	H_1H_2	H_3
dim	2	2	2
$S'_1S'_2$	2	1	-1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1	H_1H_2	H_3	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1H_2	H_3
	dim	1	1	2	2	2
P_1	1	1	1	0	-2	0
P_2	1	0	0	1	0	-1
P_3	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_1	K_2	K_3	H_1H_2	H_3	
	1	1	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/2}^1$
v^2	0	2	0	1	0	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	2	0	0	1	0	$H_{3/2}^1$

SG 183

Space group $P6mm$, No. 183, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
Δ_1	1	1	0	0	0	0	0	-1	0	0	0	0	0
Δ_2	1	0	1	0	0	0	0	0	-1	0	0	0	0
Δ_3	1	0	0	1	0	0	0	0	0	-1	0	0	0
Δ_4	1	0	0	0	1	0	0	0	0	0	-1	0	0
Δ_5	2	0	0	0	0	1	0	0	0	0	0	-1	0
Δ_6	2	0	0	0	0	0	1	0	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 6.

Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6	
1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph

v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	-1	0	0
Σ_2	1	0	1	0	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	0	0	1	-1	0	-1
T'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3
	dim	1	1	1	1	2	2	1	1	2
S_1	1	1	0	0	1	1	1	-1	0	-1
S_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4
	dim	1	1	1	1	2	2	1	1	1	1
R_1	1	1	0	1	0	1	1	-1	-1	0	0
R_2	1	0	1	0	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	L_3	L_4	H_1	H_2	H_3
	dim	1	1	1	1	1	1	2
S'_1	1	1	0	0	1	-1	0	-1
S'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	L_1	L_2	L_3	L_4	H_1	H_2	H_3	Bipartite graph
	1	1	1	1	1	1	2	
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1	H_2	H_3
dim		1	1	2	1	1	2
P_1	1	1	0	0	-1	0	0
P_2	1	0	1	0	0	-1	0
P_3	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1	H_2	H_3	
	1	1	2	1	1	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

SG 184

Space group *P6cc*, No. 184, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_2	A_3A_3	A_4A_5	A_6A_6
dim		1	1	1	1	2	2	2	4	2	4
Δ_1	1	1	0	0	0	0	0	-1	0	0	0
Δ_2	1	0	1	0	0	0	0	-1	0	0	0
Δ_3	1	0	0	1	0	0	0	0	0	-1	0
Δ_4	1	0	0	0	1	0	0	0	0	-1	0
Δ_5	2	0	0	0	0	1	0	0	-2	0	0
Δ_6	2	0	0	0	0	0	1	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_2	A_3A_3	A_4A_5	A_6A_6	
	1	1	1	1	2	2	2	4	2	4	Bipartite graph
v^1	0	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^3	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^4	1	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
dim		1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
dim		1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	-1	0	0
Σ_2	1	0	1	0	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$

v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	L_1L_2	L_3L_4
	dim	1	1	1	1	2	2
U_1	1	1	0	0	0	-1	0
U_2	1	0	1	0	0	0	-1
U_3	1	0	0	1	0	-1	0
U_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1	M_2	M_3	M_4	L_1L_2	L_3L_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	0	1	0	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	0	0	1	-1	0	-1
T'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3A_3	A_4A_5	A_6A_6	H_1H_2	H_3H_3
dim	2	4	2	4	2	4
S_1S_2	2	1	2	1	2	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1A_2	A_3A_3	A_4A_5	A_6A_6	H_1H_2	H_3H_3	
	2	4	2	4	2	4	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	2	0	$H_{3/2}^1$
v^3	0	0	1	0	1	0	$H_{2/1}^1$
v^4	0	0	2	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	1	0	0	1	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3A_3	A_4A_5	A_6A_6	L_1L_2	L_3L_4
dim	2	4	2	4	2	2
R_1R_2	2	1	2	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1A_2	A_3A_3	A_4A_5	A_6A_6	L_1L_2	L_3L_4	Bipartite graph
	2	4	2	4	2	2	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	2	$H_{3/2}^1$
v^7	0	1	0	0	1	1	$H_{3/2}^1$
v^8	0	1	0	0	2	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1L_2	L_3L_4	H_1H_2	H_3H_3
dim	2	2	2	4
$S'_1S'_2$	2	1	1	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1L_2	L_3L_4	H_1H_2	H_3H_3	Bipartite graph
	2	2	2	4	
v^1	0	1	1	0	$H_{2/1}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	0	1	0	$H_{2/1}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_1	K_2	K_3	H_1H_2	H_3H_3
	dim	1	1	2	2	4
P_1	1	1	0	0	-1	0
P_2	1	0	1	0	-1	0
P_3	2	0	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_1	K_2	K_3	H_1H_2	H_3H_3	
	1	1	2	2	4	Bipartite graph
v^1	0	0	2	0	1	$H_{3/2}^1$
v^2	1	1	0	1	0	$H_{3/2}^1$

SG 185

Space group $P6_3cm$, No. 185, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_4	A_2A_5	A_3A_6
	dim	1	1	1	1	2	2	2	2	4
Δ_1	1	1	0	0	0	0	0	-1	0	0
Δ_2	1	0	1	0	0	0	0	0	-1	0
Δ_3	1	0	0	1	0	0	0	0	-1	0
Δ_4	1	0	0	0	1	0	0	-1	0	0
Δ_5	2	0	0	0	0	1	0	0	0	-1
Δ_6	2	0	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_4	A_2A_5	A_3A_6	
	1	1	1	1	2	2	2	2	4	Bipartite graph
v^1	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	Bipartite graph
	1	1	1	1	2	2	1	1	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	-1	0	0
Σ_2	1	0	1	0	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$

v^8	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	L_1L_2	L_3L_4
	dim	1	1	1	1	2	2
U_1	1	1	0	0	0	-1	0
U_2	1	0	1	0	0	0	-1
U_3	1	0	0	1	0	0	-1
U_4	1	0	0	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1	M_2	M_3	M_4	L_1L_2	L_3L_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	0	0	1	-1	0	-1
T'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1A_4	A_2A_5	A_3A_6	H_1H_1	H_2H_2	H_3H_3	
dim	2	2	4	2	2	4	
S_1S_1	2	1	0	1	-1	0	-1
S_2S_2	2	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1A_4	A_2A_5	A_3A_6	H_1H_1	H_2H_2	H_3H_3	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_4	A_2A_5	A_3A_6	L_1L_2	L_3L_4	
dim	2	2	4	2	2	
R_1R_2	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1A_4	A_2A_5	A_3A_6	L_1L_2	L_3L_4	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1L_2	L_3L_4	H_1H_1	H_2H_2	H_3H_3
	dim	2	2	2	2	4
$S'_1S'_1$	2	1	0	-1	0	-1
$S'_2S'_2$	2	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1L_2	L_3L_4	H_1H_1	H_2H_2	H_3H_3	Bipartite graph
	2	2	2	2	4	
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	H_1H_1	H_2H_2	H_3H_3
	dim	1	1	2	2	2	4
P_1	1	1	0	0	-2	0	0
P_2	1	0	1	0	0	-2	0
P_3	2	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_1	K_2	K_3	H_1H_1	H_2H_2	H_3H_3	
	1	1	2	2	2	4	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	2	0	0	1	0	$H_{3/2}^1$
v^3	2	0	0	1	0	0	$H_{3/2}^1$

SG 186

Space group $P6_3mc$, No. 186, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_4	A_2A_5	A_3A_6
	dim	1	1	1	1	2	2	2	2	4
Δ_1	1	1	0	0	0	0	0	-1	0	0
Δ_2	1	0	1	0	0	0	0	0	-1	0
Δ_3	1	0	0	1	0	0	0	-1	0	0
Δ_4	1	0	0	0	1	0	0	0	-1	0
Δ_5	2	0	0	0	0	1	0	0	0	-1
Δ_6	2	0	0	0	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_4	A_2A_5	A_3A_6	
	1	1	1	1	2	2	2	2	4	Bipartite graph
v^1	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^2	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^3	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3
	dim	1	1	1	1	2	2	1	1	2
T_1	1	1	0	0	1	1	1	-1	0	-1
T_2	1	0	1	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	
	1	1	1	1	2	2	1	1	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
dim		1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	1	0	1	1	-1	-1	0	0
Σ_2	1	0	1	0	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	
	1	1	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$

v^{14}	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	L_1L_2	L_3L_4
	dim	1	1	1	1	2	2
U_1	1	1	0	0	0	-1	0
U_2	1	0	1	0	0	-1	0
U_3	1	0	0	1	0	0	-1
U_4	1	0	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1	M_2	M_3	M_4	L_1L_2	L_3L_4	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	1	1	0	1	$H_{3/2}^1$
v^2	1	1	0	0	1	0	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3
	dim	1	1	1	1	1	1	2
T'_1	1	1	0	0	1	-1	0	-1
T'_2	1	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	
	1	1	1	1	1	1	2	Bipartite graph
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1A_4	A_2A_5	A_3A_6	H_1H_2	H_3
dim	2	2	4	2	2
S_1S_2	2	1	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1A_4	A_2A_5	A_3A_6	H_1H_2	H_3	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_4	A_2A_5	A_3A_6	L_1L_2	L_3L_4
dim	2	2	4	2	2
R_1R_1	2	1	0	1	-1 0
R_2R_2	2	0	1	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1A_4	A_2A_5	A_3A_6	L_1L_2	L_3L_4	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_1L_2	L_3L_4	H_1H_2	H_3
dim	2	2	2	2
$S'_1S'_2$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_1L_2	L_3L_4	H_1H_2	H_3	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_1	K_2	K_3	H_1H_2	H_3
dim	1	1	2	2	2
P_1	1	1	0	0	-1 0
P_2	1	0	1	0	-1 0
P_3	2	0	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_1	K_2	K_3	H_1H_2	H_3	
	1	1	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	1	1	0	1	0	$H_{3/2}^1$

SG 187

Space group $P\bar{6}m2$, No. 187, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	-1	0	0	-1	0	0
Δ_2	1	0	1	1	0	0	0	0	-1	-1	0	0	0
Δ_3	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	2	2	1	1	1	1	1	1
T_1	1	1	1	0	0	2	0	-1	-1	-1	0	0	0
T_2	1	0	0	1	1	0	2	0	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	0	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	0	0	0	2	0	0	$H_{3/2}^1$

v^7	0	0	0	0	1	0	0	0	2	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	1	0	0	2	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	0	0	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	0	0	1	0	-1	0	0	0
Σ_2	1	0	0	0	1	0	1	0	-1	0	0
Σ_3	1	0	0	1	0	0	1	0	0	-1	0
Σ_4	1	0	1	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	H_4	H_5	H_6
	dim	1	1	1	1	2	2	1	1	1	1	1	1
S_1	1	1	1	0	0	2	0	-1	-1	-1	0	0	0
S_2	1	0	0	1	1	0	2	0	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	H_4	H_5	H_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	0	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	0	2	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	1	0	0	2	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	0	0	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4
	dim	1	1	1	1	2	2	1	1	1	1
R_1	1	1	0	0	0	1	0	-1	0	0	0
R_2	1	0	0	0	1	0	1	0	-1	0	0
R_3	1	0	0	1	0	0	1	0	0	-1	0
R_4	1	0	1	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4	
	1	1	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 188

Space group $P\bar{6}c2$, No. 188, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	$A_1 A_2$	A_3
	dim	1	1	1	1	2	2	4	2
Δ_1	1	1	0	0	1	0	0	0	-1
Δ_2	1	0	1	1	0	0	0	0	-1
Δ_3	2	0	0	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	$A_1 A_2$	A_3	
	1	1	1	1	2	2	4	2	Bipartite graph
v^1	0	0	0	0	0	2	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	1	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	2	2	1	1	1	1	1	1
T_1	1	1	1	0	0	2	0	-1	-1	-1	0	0	0
T_2	1	0	0	1	1	0	2	0	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	0	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	0	2	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	1	0	0	2	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	0	0	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{22}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	0	0	0	1	0	-1	0	0	0
Σ_2	1	0	0	0	1	0	1	0	-1	0	0
Σ_3	1	0	0	1	0	0	1	0	0	-1	0
Σ_4	1	0	1	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	
	1	1	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	L_1
	dim	1	1	1	1	2
U_1	1	1	1	0	0	-1
U_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	L_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3	H_1H_4	H_2H_5	H_3H_6
dim	4	2	2	2	2
S_1S_2	2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_1A_2	A_3	H_1H_4	H_2H_5	H_3H_6	Bipartite graph
	4	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	2	$H_{3/2}^1$
v^5	1	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	1	0	1	$H_{3/2}^1$
v^8	1	0	1	1	0	$H_{3/2}^1$
v^9	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1A_2	A_3	L_1
dim	4	2	2
R_1	2	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_1A_2	A_3	L_1	Bipartite graph
	4	2	2	
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	2	$H_{3/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	H_1H_4	H_2H_5	H_3H_6
dim	2	2	2	2
$S'_1S'_2$	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	H_1H_4	H_2H_5	H_3H_6	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_1	K_2	K_3	K_4	K_5	K_6	H_1H_4	H_2H_5	H_3H_6
dim	1	1	1	1	1	1	2	2	2
P_1	1	1	0	0	1	0	0	-2	0
P_2	1	0	0	1	0	0	1	0	-2
P_3	1	0	1	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	K_1	K_2	K_3	K_4	K_5	K_6	H_1H_4	H_2H_5	H_3H_6	
	1	1	1	1	1	1	2	2	2	Bipartite graph
v^1	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^9	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$

SG 189

Space group $P\bar{6}2m$, No. 189, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
Δ_1	1	1	0	0	1	0	0	-1	0	0	-1	0	0
Δ_2	1	0	1	1	0	0	0	0	-1	-1	0	0	0
Δ_3	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1	A_2	A_3	A_4	A_5	A_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
T_1	1	1	0	0	0	1	0	-1	0	0	0	-1	0
T_2	1	0	0	0	1	0	1	0	0	0	-1	0	-1
T_3	1	0	0	1	0	0	1	0	0	-1	0	0	-1
T_4	1	0	1	0	0	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6	
	1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	1	0	0	2	0	-1	0	0	-1
Σ_2	1	0	0	1	1	0	2	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	
	1	1	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	1	1	1	1	2	2
T'_1	1	1	0	0	0	-1	0	0	0	-1	0
T'_2	1	0	1	0	0	0	0	0	-1	0	-1
T'_3	1	0	0	1	0	0	0	-1	0	0	-1
T'_4	1	0	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^3	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

 S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	H_4	H_5	H_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
S_1	1	1	0	0	0	1	0	-1	0	0	0	-1	0
S_2	1	0	0	0	1	0	1	0	0	0	-1	0	-1
S_3	1	0	0	1	0	0	1	0	0	-1	0	0	-1
S_4	1	0	1	0	0	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1	A_2	A_3	A_4	A_5	A_6	H_1	H_2	H_3	H_4	H_5	H_6	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

v^6	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4
	dim	1	1	1	1	2	2	1	1	1	1
R_1	1	1	1	0	0	2	0	-1	0	0	-1
R_2	1	0	0	1	1	0	2	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	A_1	A_2	A_3	A_4	A_5	A_6	L_1	L_2	L_3	L_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	L_3	L_4	H_1	H_2	H_3	H_4	H_5	H_6
	dim	1	1	1	1	1	1	1	1	2	2

S'_1	1	1	0	0	0	-1	0	0	0	-1	0
S'_2	1	0	1	0	0	0	0	0	-1	0	-1
S'_3	1	0	0	1	0	0	0	-1	0	0	-1
S'_4	1	0	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_1	L_2	L_3	L_4	H_1	H_2	H_3	H_4	H_5	H_6	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^3	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_1	K_2	K_3	K_4	K_5	K_6	H_1	H_2	H_3	H_4	H_5	H_6
dim		1	1	1	1	2	2	1	1	1	1	2	2
P_1	1	1	0	0	1	0	0	-1	0	0	-1	0	0
P_2	1	0	1	1	0	0	0	0	-1	-1	0	0	0
P_3	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	K_1	K_2	K_3	K_4	K_5	K_6	H_1	H_2	H_3	H_4	H_5	H_6	
	1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

SG 190

Space group $P\bar{6}2c$, No. 190, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_2	A_3
	dim	1	1	1	1	2	2	4	2
Δ_1	1	1	0	0	1	0	0	0	-1
Δ_2	1	0	1	1	0	0	0	0	-1
Δ_3	2	0	0	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	A_1A_2	A_3	
	1	1	1	1	2	2	4	2	Bipartite graph
v^1	0	0	0	0	0	2	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	1	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
T_1	1	1	0	0	0	1	0	-1	0	0	0	-1	0
T_2	1	0	0	0	1	0	1	0	0	0	-1	0	-1
T_3	1	0	0	1	0	0	1	0	0	-1	0	0	-1
T_4	1	0	1	0	0	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	K_1	K_2	K_3	K_4	K_5	K_6	
	1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph

v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	1	1	1	1
Σ_1	1	1	1	0	0	2	0	-1	0	0	-1
Σ_2	1	0	0	1	1	0	2	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	Γ_6	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	L_1
dim		1	1	1	1	2
U_1	1	1	1	0	0	-1
U_2	1	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1	M_2	M_3	M_4	L_1	
	1	1	1	1	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1	M_2	M_3	M_4	K_1	K_2	K_3	K_4	K_5	K_6
dim		1	1	1	1	1	1	1	1	2	2
T'_1	1	1	0	0	0	-1	0	0	0	-1	0
T'_2	1	0	1	0	0	0	0	0	-1	0	-1
T'_3	1	0	0	1	0	0	0	-1	0	0	-1
T'_4	1	0	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_1	M_2	M_3	M_4	K_1	K_2	K_3	K_4	K_5	K_6	
	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^3	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	$A_1 A_2 A_3$	$H_1 H_2 H_3$
dim	4 2	2 2 2
S_1	2 1	-1 -1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	$A_1 A_2 A_3$	$H_1 H_2 H_3$	Bipartite graph
	4 2	2 2 2	
v^1	0 1	0 0 1	$H_{2/1}^1$
v^2	0 1	0 1 0	$H_{2/1}^1$
v^3	0 1	1 0 0	$H_{2/1}^1$
v^4	1 0	0 0 2	$H_{3/2}^1$
v^5	1 0	0 1 1	$H_{3/2}^1$
v^6	1 0	0 2 0	$H_{3/2}^1$
v^7	1 0	1 0 1	$H_{3/2}^1$
v^8	1 0	1 1 0	$H_{3/2}^1$
v^9	1 0	2 0 0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$A_1 A_2 A_3$	L_1
dim	4 2	2
$R_1 R_2$	2 1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$A_1 A_2 A_3$	L_1	Bipartite graph
	4 2	2	
v^1	0 1	1	$H_{2/1}^1$
v^2	1 0	2	$H_{3/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_1	H_1	H_2	H_3
	dim	2	2	2	2
S'_1	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	H_1	H_2	H_3	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0)$, $H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	K_4	K_5	K_6	H_1	H_2	H_3
	dim	1	1	1	1	2	2	2	2	2
P_1	1	1	0	0	1	0	0	0	0	-1
P_2	1	0	1	1	0	0	0	0	0	-1
P_3	2	0	0	0	0	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	K_1	K_2	K_3	K_4	K_5	K_6	H_1	H_2	H_3	
	1	1	1	1	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 191

Space group $P6/mmm$, No. 191, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	A_6^+	A_6^-
dim		1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2
Δ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	-1	0	0	-1	0	0	0	0	0	0	0	0
Δ_2	1	0	1	1	0	0	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0	0	0	0
Δ_3	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	-1	-1	0	0	0	0	0	0
Δ_4	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	-1	0	0	-1	0	0	0	0
Δ_5	2	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	-1	-1
Δ_6	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{24}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	A_6^+	A_6^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{15}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{17}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{18}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6
dim		1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2
T_1	1	1	0	0	0	0	0	0	1	0	1	1	0	-1	0	0	0	-1	0
T_2	1	0	0	0	1	1	0	0	0	1	0	0	1	0	0	0	-1	0	-1
T_3	1	0	1	0	0	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1
T_4	1	0	0	1	0	0	1	0	0	0	1	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{16}	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{18}	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{19}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{24}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	
dim		1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	1
Σ_1	1	1	0	0	0	0	1	0	0	0	1	1	0	-1	0	0	0	0	0	0	-1	
Σ_2	1	0	0	0	1	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1	0	0	
Σ_3	1	0	1	0	0	1	0	0	0	1	0	0	1	0	-1	0	0	0	0	-1	0	
Σ_4	1	0	0	1	0	0	0	0	1	0	1	1	0	0	0	0	-1	-1	0	0	0	

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{24}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{26}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$

$$\overline{v^{32} \mid 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \mid 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \mid H_{2/1}^1}$$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	1	1	1	1	1	1	1	1	2	2
T'_1	1	1	0	0	1	0	0	0	0	-1	0	0	0	-1	0
T'_2	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	-1
T'_3	1	0	1	1	0	0	0	0	0	0	0	-1	0	0	-1
T'_4	1	0	0	0	0	1	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
v^1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	A_6^+	A_6^-	H_1	H_2	H_3	H_4	H_5	H_6
	dim	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2

R_1	1	1	0	0	0	0	0	1	0	0	0	1	1	0	-1	0	0	0	0	0	0	-1
R_2	1	0	0	0	1	0	0	1	0	1	0	0	0	1	0	0	-1	0	0	-1	0	0
R_3	1	0	1	0	0	1	0	0	0	1	0	0	0	1	0	-1	0	0	0	0	-1	0
R_4	1	0	0	1	0	0	0	0	1	0	1	1	0	0	0	0	-1	-1	0	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	A_1^+	A_1^-	A_2^+	A_2^-	A_3^+	A_3^-	A_4^+	A_4^-	A_5^+	A_5^-	A_6^+	A_6^-	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	L_4^+	L_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{24}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{26}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of *S'*: $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	L_4^+	L_4^-	H_1	H_2	H_3	H_4	H_5	H_6
	dim	1	1	1	1	1	1	1	1	1	1	1	1	2	2
S'_1	1	1	0	0	1	0	0	0	0	-1	0	0	0	-1	0
S'_2	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	-1
S'_3	1	0	1	1	0	0	0	0	0	0	0	-1	0	0	-1
S'_4	1	0	0	0	0	1	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	L_4^+	L_4^-	H_1	H_2	H_3	H_4	H_5	H_6	
	1	1	1	1	1	1	1	1	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	K_4	K_5	K_6	H_1	H_2	H_3	H_4	H_5	H_6
	dim	1	1	1	1	2	2	1	1	1	1	2	2
P_1	1	1	0	0	1	0	0	-1	0	0	-1	0	0
P_2	1	0	1	1	0	0	0	0	-1	-1	0	0	0
P_3	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	K_1	K_2	K_3	K_4	K_5	K_6	H_1	H_2	H_3	H_4	H_5	H_6	
	1	1	1	1	2	2	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

SG 192

Space group $P6/mcc$, No. 192, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1A_2	A_3	A_4A_5	A_6
dim		1	1	1	1	1	1	1	1	2	2	2	2	4	2	4	2
Δ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	-1	0	0
Δ_2	1	0	1	1	0	0	0	0	0	0	0	0	0	0	-1	0	0
Δ_3	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	-1
Δ_4	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	-1
Δ_5	2	0	0	0	0	0	0	0	0	0	0	1	1	-2	0	0	0
Δ_6	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1A_2	A_3	A_4A_5	A_6	
	1	1	1	1	1	1	1	1	2	2	2	2	4	2	4	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	0	0	2	1	0	0	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$

v^7	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	1	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{12}	0	1	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{13}	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{14}	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6
dim		1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2
T_1	1	1	0	0	0	0	0	0	1	0	1	1	0	-1	0	0	0	-1	0
T_2	1	0	0	0	1	1	0	0	0	1	0	0	1	0	0	0	-1	0	-1
T_3	1	0	1	0	0	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1
T_4	1	0	0	1	0	0	1	0	0	0	1	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{18}	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

v^{19}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	
	dim	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	1
Σ_1	1	1	0	0	0	0	1	0	0	0	1	1	0	-1	0	0	0	0	0	0	0	-1
Σ_2	1	0	0	0	1	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1	0	0	0
Σ_3	1	0	1	0	0	1	0	0	0	1	0	0	1	0	-1	0	0	0	0	-1	0	0
Σ_4	1	0	0	1	0	0	0	0	1	0	1	1	0	0	0	0	-1	-1	0	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

v^{21}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{24}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{26}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	L_1	L_2
dim		1	1	1	1	1	1	1	1	2	2
U_1	1	1	0	0	0	0	1	0	0	-1	0
U_2	1	0	0	1	0	0	0	0	1	0	-1
U_3	1	0	1	0	0	1	0	0	0	-1	0
U_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	L_1	L_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6	
dim	1	1	1	1	1	1	1	1	1	1	1	1	2	2	
T'_1	1	1	0	0	1	0	0	0	0	-1	0	0	0	-1	0
T'_2	1	0	0	0	0	0	1	1	0	0	0	-1	0	-1	
T'_3	1	0	1	1	0	0	0	0	0	0	0	-1	0	0	-1
T'_4	1	0	0	0	0	1	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1A_2	A_3	A_4A_5	A_6	H_1H_2	H_3	
dim	4	2	4	2	4	2	
S_1	2	2	1	2	1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

A_1A_2	A_3	A_4A_5	A_6	H_1H_2	H_3	Bipartite graph
4	2	4	2	4	2	

v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	2	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	2	$H_{3/2}^1$
v^4	0	0	1	0	1	0	$H_{2/2}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	1	1	0	$H_{3/2}^1$
v^7	0	2	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	0	0	2	$H_{3/2}^1$
v^9	1	0	0	0	1	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_1A_2	A_3	A_4A_5	A_6	L_1	L_2
	dim	4	2	4	2	2	2
R_1	2	2	1	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_1A_2	A_3	A_4A_5	A_6	L_1	L_2	Bipartite graph
	4	2	4	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	2	$H_{3/2}^1$
v^4	0	0	1	0	1	1	$H_{3/2}^1$
v^5	0	0	1	0	2	0	$H_{3/2}^1$
v^6	0	1	0	0	0	1	$H_{2/1}^1$
v^7	0	1	0	0	1	0	$H_{2/1}^1$
v^8	1	0	0	0	0	2	$H_{3/2}^1$
v^9	1	0	0	0	1	1	$H_{3/2}^1$
v^{10}	1	0	0	0	2	0	$H_{3/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_1	L_2	H_1H_2	H_3
	dim	2	2	4	2
S'_1	2	1	1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1	L_2	H_1H_2	H_3	
	2	2	4	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	2	1	0	$H_{3/2}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	1	1	0	$H_{3/2}^1$
v^5	2	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_1	K_2	K_3	K_4	K_5	K_6	H_1H_2	H_3
	dim	1	1	1	1	2	2	4	2
P_1	1	1	0	0	1	0	0	0	-1
P_2	1	0	1	1	0	0	0	0	-1
P_3	2	0	0	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	K_1	K_2	K_3	K_4	K_5	K_6	H_1H_2	H_3	
	1	1	1	1	2	2	4	2	Bipartite graph
v^1	0	0	0	0	0	2	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^7	1	1	0	0	0	0	0	1	$H_{3/2}^1$

SG 193

Space group $P6_3/mcm$, No. 193, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1	A_2	A_3
	dim	1	1	1	1	1	1	1	1	2	2	2	2	2	2	4
Δ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	-1	0	0
Δ_2	1	0	1	1	0	0	0	0	0	0	0	0	0	0	-1	0
Δ_3	1	0	0	0	0	0	1	1	0	0	0	0	0	0	-1	0
Δ_4	1	0	0	0	0	1	0	0	1	0	0	0	0	-1	0	0
Δ_5	2	0	0	0	0	0	0	0	0	0	0	1	1	0	0	-1
Δ_6	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{15}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1	A_2	A_3	
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	4	Bipartite graph
v^1	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	1	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2
T_1	1	1	0	0	0	0	0	0	1	0	1	1	0	-1	0	0	0	-1	0
T_2	1	0	0	0	1	1	0	0	0	1	0	0	1	0	0	0	-1	0	-1
T_3	1	0	1	0	0	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1
T_4	1	0	0	1	0	0	1	0	0	0	1	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6	
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$	
v^2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/2}^1$
v^6	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{15}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{18}	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{19}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{20}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{24}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	
	dim	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	1
Σ_1	1	1	0	0	0	0	1	0	0	0	1	1	0	-1	0	0	0	0	0	0	0	-1
Σ_2	1	0	0	0	1	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1	0	0	0
Σ_3	1	0	1	0	0	1	0	0	0	1	0	0	1	0	-1	0	0	0	0	-1	0	0
Σ_4	1	0	0	1	0	0	0	0	1	0	1	1	0	0	0	0	-1	-1	0	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	
1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	Bipartite graph

v ¹	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	H _{3/2} ¹
v ²	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	H _{3/2} ¹
v ³	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	H _{3/2} ¹
v ⁴	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	H _{3/2} ¹
v ⁵	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	H _{3/2} ¹
v ⁶	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	H _{3/2} ¹
v ⁷	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	H _{3/2} ¹
v ⁸	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	H _{3/2} ¹
v ⁹	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	H _{3/2} ¹
v ¹⁰	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	H _{3/2} ¹
v ¹¹	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	H _{3/2} ¹
v ¹²	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	H _{3/2} ¹
v ¹³	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	0	H _{3/2} ¹
v ¹⁴	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	H _{3/2} ¹
v ¹⁵	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	H _{3/2} ¹
v ¹⁶	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	H _{3/2} ¹
v ¹⁷	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ¹⁸	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ¹⁹	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	H _{2/1} ¹
v ²⁰	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ²¹	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	H _{2/1} ¹
v ²²	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	H _{2/1} ¹
v ²³	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	H _{2/1} ¹
v ²⁴	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ²⁵	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	H _{2/1} ¹
v ²⁶	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ²⁷	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ²⁸	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ²⁹	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	H _{2/1} ¹
v ³⁰	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	H _{2/1} ¹
v ³¹	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	H _{2/1} ¹
v ³²	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	H _{2/1} ¹

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M ₁ ⁺	M ₁ ⁻	M ₂ ⁺	M ₂ ⁻	M ₃ ⁺	M ₃ ⁻	M ₄ ⁺	M ₄ ⁻	L ₁	L ₂
	dim	1	1	1	1	1	1	1	1	2	2
U ₁	1	1	0	0	0	0	1	0	0	-1	0
U ₂	1	0	0	1	0	0	0	0	1	0	-1

U_3	1	0	1	0	0	1	0	0	0	0	-1
U_4	1	0	0	0	1	0	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	L_1	L_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	1	1	1	1	1	1	1	1	2	2
T'_1	1	1	0	0	1	0	0	0	0	-1	0	0	0	-1	0
T'_2	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	-1
T'_3	1	0	1	1	0	0	0	0	0	0	0	-1	0	0	-1
T'_4	1	0	0	0	0	1	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

v^{12}	0	1	0	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{16}	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	H_1H_4	H_2H_3	H_5H_6	
dim	2	2	4	2	2	4	
S_1S_2	2	1	0	1	-1	0	-1
S_3S_4	2	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	A_1	A_2	A_3	H_1H_4	H_2H_3	H_5H_6	Bipartite graph
	2	2	4	2	2	4	
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	L_1	L_2	
dim	2	2	4	2	2	
R_1	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_1	A_2	A_3	L_1	L_2	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$

v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	L_2	H_1H_4	H_2H_3	H_5H_6	
dim	2	2	2	2	4	
$S'_1S'_2$	2	1	0	-1	0	-1
$S'_3S'_4$	2	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_1	L_2	H_1H_4	H_2H_3	H_5H_6	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	1	0	1	0	0	$H_{2/1}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	K_1	K_2	K_3	K_4	K_5	K_6	H_1H_4	H_2H_3	H_5H_6
dim	1	1	1	1	2	2	2	2	4
P_1	1	0	0	1	0	0	-2	0	0
P_2	1	0	1	1	0	0	0	-2	0
P_3	2	0	0	0	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	K_1	K_2	K_3	K_4	K_5	K_6	H_1H_4	H_2H_3	H_5H_6	
	1	1	1	1	2	2	2	2	4	Bipartite graph
v^1	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	2	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$

v^7	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^9	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$

SG 194

Space group $P6_3/mmc$, No. 194, without SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1	A_2	A_3
dim		1	1	1	1	1	1	1	1	2	2	2	2	2	2	4
Δ_1	1	1	0	0	1	0	0	0	0	0	0	0	0	-1	0	0
Δ_2	1	0	1	1	0	0	0	0	0	0	0	0	0	0	-1	0
Δ_3	1	0	0	0	0	0	1	1	0	0	0	0	0	-1	0	0
Δ_4	1	0	0	0	0	1	0	0	1	0	0	0	0	0	-1	0
Δ_5	2	0	0	0	0	0	0	0	0	0	0	1	1	0	0	-1
Δ_6	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{15}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	A_1	A_2	A_3	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	2	4	
v^1	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	1	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6
dim		1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2
T_1	1	1	0	0	0	0	0	0	1	0	1	1	0	-1	0	0	0	-1	0
T_2	1	0	0	0	1	1	0	0	0	1	0	0	1	0	0	-1	0	0	-1
T_3	1	0	1	0	0	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1
T_4	1	0	0	1	0	0	1	0	0	0	1	1	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{18}	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{19}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{21}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	
dim		1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	1
Σ_1	1	1	0	0	0	0	1	0	0	0	1	1	0	-1	0	0	0	0	0	0	0	-1
Σ_2	1	0	0	0	1	0	0	1	0	1	0	0	1	0	0	-1	0	0	-1	0	0	0
Σ_3	1	0	1	0	0	1	0	0	0	1	0	0	1	0	-1	0	0	0	0	0	-1	0
Σ_4	1	0	0	1	0	0	0	0	1	0	1	1	0	0	0	0	-1	-1	0	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	Γ_6^+	Γ_6^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{21}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{24}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{26}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$

$$\underline{v^{32}} \mid 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \mid 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \mid H_{2/1}^1$$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	L_1	L_2
	dim	1	1	1	1	1	1	1	1	2	2
U_1	1	1	0	0	0	0	1	0	0	-1	0
U_2	1	0	0	1	0	0	0	0	1	-1	0
U_3	1	0	1	0	0	1	0	0	0	0	-1
U_4	1	0	0	0	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	L_1	L_2	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^7	1	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6
	dim	1	1	1	1	1	1	1	1	1	1	1	1	2	2
T'_1	1	1	0	0	1	0	0	0	0	-1	0	0	0	-1	0
T'_2	1	0	0	0	0	0	1	1	0	0	0	0	-1	0	-1
T'_3	1	0	1	1	0	0	0	0	0	0	0	-1	0	0	-1
T'_4	1	0	0	0	0	1	0	0	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	K_1	K_2	K_3	K_4	K_5	K_6	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_1	A_2	A_3	H_1	H_2	H_3
dim	2	2	4	2	2	2
S_1	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_1	A_2	A_3	H_1	H_2	H_3	Bipartite graph
	2	2	4	2	2	2	
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$

v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_1	A_2	A_3	L_1	L_2	
dim	2	2	4	2	2	
$R_1 R_2$	2	1	0	1	-1	0
$R_3 R_4$	2	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_1	A_2	A_3	L_1	L_2	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	0	1	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_1	L_2	H_1	H_2	H_3	
dim	2	2	2	2	2	
S'_1	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_1	L_2	H_1	H_2	H_3	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

v^1	0	0	1	0	1	0	2	$H_{4/3}^1$
v^2	0	0	1	0	1	1	1	$H_{4/3}^1$
v^3	0	0	1	0	1	2	0	$H_{4/3}^1$
v^4	0	0	1	1	0	0	2	$H_{4/3}^1$
v^5	0	0	1	1	0	1	1	$H_{4/3}^1$
v^6	0	0	1	1	0	2	0	$H_{4/3}^1$
v^7	0	1	0	0	2	0	0	$H_{3/2}^1$
v^8	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{10}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	M_1	M_2	M_3	M_4
dim	1	2	3	1	1	1	1
Σ_1	1	2	3	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	M_1	M_2	M_3	M_4	Bipartite graph
	1	2	3	1	1	1	1	
v^1	0	0	1	0	0	0	3	$H_{4/3}^1$
v^2	0	0	1	0	0	1	2	$H_{4/3}^1$
v^3	0	0	1	0	0	2	1	$H_{4/3}^1$
v^4	0	0	1	0	0	3	0	$H_{4/3}^1$
v^5	0	0	1	0	1	0	2	$H_{4/3}^1$
v^6	0	0	1	0	1	1	1	$H_{4/3}^1$
v^7	0	0	1	0	1	2	0	$H_{4/3}^1$
v^8	0	0	1	0	2	0	1	$H_{4/3}^1$
v^9	0	0	1	0	2	1	0	$H_{4/3}^1$
v^{10}	0	0	1	0	3	0	0	$H_{4/3}^1$
v^{11}	0	0	1	1	0	0	2	$H_{4/3}^1$
v^{12}	0	0	1	1	0	1	1	$H_{4/3}^1$
v^{13}	0	0	1	1	0	2	0	$H_{4/3}^1$
v^{14}	0	0	1	1	1	0	1	$H_{4/3}^1$
v^{15}	0	0	1	1	1	1	0	$H_{4/3}^1$
v^{16}	0	0	1	1	2	0	0	$H_{4/3}^1$
v^{17}	0	0	1	2	0	0	1	$H_{4/3}^1$

v^{18}	0	0	1	2	0	1	0	$H_{4/3}^1$
v^{19}	0	0	1	2	1	0	0	$H_{4/3}^1$
v^{20}	0	0	1	3	0	0	0	$H_{4/3}^1$
v^{21}	0	1	0	0	0	0	2	$H_{3/2}^1$
v^{22}	0	1	0	0	0	1	1	$H_{3/2}^1$
v^{23}	0	1	0	0	0	2	0	$H_{3/2}^1$
v^{24}	0	1	0	0	1	0	1	$H_{3/2}^1$
v^{25}	0	1	0	0	1	1	0	$H_{3/2}^1$
v^{26}	0	1	0	0	2	0	0	$H_{3/2}^1$
v^{27}	0	1	0	1	0	0	1	$H_{3/2}^1$
v^{28}	0	1	0	1	0	1	0	$H_{3/2}^1$
v^{29}	0	1	0	1	1	0	0	$H_{3/2}^1$
v^{30}	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{31}	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{32}	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{33}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{34}	1	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	R_1	R_2R_3	R_4
	dim	1	2	3	1	2	3
Λ_1	1	1	0	1	-1	0	-1
Λ_2	1	0	1	1	0	-1	-1
Λ_3	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	R_1	R_2R_3	R_4	
	1	2	3	1	2	3	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/3}^1$
v^2	0	0	1	1	1	0	$H_{3/3}^1$
v^3	0	1	0	0	1	0	$H_{2/2}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	1	$H_{3/3}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	R_1	R_2R_3	R_4
dim		1	1	1	1	1	2	3
S_1	1	1	1	1	1	-1	-2	-3

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	X_1	X_2	X_3	X_4	R_1	R_2R_3	R_4	Bipartite graph
	1	1	1	1	1	2	3	
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	0	2	0	1	0	$H_{3/2}^1$
v^3	0	0	0	3	0	0	1	$H_{4/3}^1$
v^4	0	0	1	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	1	0	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	1	$H_{4/3}^1$
v^7	0	0	2	0	0	1	0	$H_{3/2}^1$
v^8	0	0	2	1	0	0	1	$H_{4/3}^1$
v^9	0	0	3	0	0	0	1	$H_{4/3}^1$
v^{10}	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	1	0	1	0	1	0	$H_{3/2}^1$
v^{12}	0	1	0	2	0	0	1	$H_{4/3}^1$
v^{13}	0	1	1	0	0	1	0	$H_{3/2}^1$
v^{14}	0	1	1	1	0	0	1	$H_{4/3}^1$
v^{15}	0	1	2	0	0	0	1	$H_{4/3}^1$
v^{16}	0	2	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	2	0	1	0	0	1	$H_{4/3}^1$
v^{18}	0	2	1	0	0	0	1	$H_{4/3}^1$
v^{19}	0	3	0	0	0	0	1	$H_{4/3}^1$
v^{20}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{21}	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{22}	1	0	0	2	0	0	1	$H_{4/3}^1$
v^{23}	1	0	1	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	1	1	0	0	1	$H_{4/3}^1$
v^{25}	1	0	2	0	0	0	1	$H_{4/3}^1$
v^{26}	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{27}	1	1	0	1	0	0	1	$H_{4/3}^1$
v^{28}	1	1	1	0	0	0	1	$H_{4/3}^1$
v^{29}	1	2	0	0	0	0	1	$H_{4/3}^1$
v^{30}	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{31}	2	0	0	1	0	0	1	$H_{4/3}^1$
v^{32}	2	0	1	0	0	0	1	$H_{4/3}^1$
v^{33}	2	1	0	0	0	0	1	$H_{4/3}^1$
v^{34}	3	0	0	0	0	0	1	$H_{4/3}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	R_1	R_2R_3	R_4
	dim	1	1	1	1	1	2	3
T_1	1	1	0	1	0	-1	-2	-1
T_2	1	0	1	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	M_1	M_2	M_3	M_4	R_1	R_2R_3	R_4	Bipartite graph
	1	1	1	1	1	2	3	
v^1	0	0	1	0	1	0	0	$H_{2/1}^1$
v^2	0	0	1	2	0	0	1	$H_{4/3}^1$
v^3	0	0	2	0	0	1	0	$H_{3/2}^1$
v^4	0	1	1	1	0	0	1	$H_{4/3}^1$
v^5	0	2	1	0	0	0	1	$H_{4/3}^1$
v^6	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	2	0	0	1	$H_{4/3}^1$
v^8	1	0	1	0	0	1	0	$H_{3/2}^1$
v^9	1	1	0	1	0	0	1	$H_{4/3}^1$
v^{10}	1	2	0	0	0	0	1	$H_{4/3}^1$
v^{11}	2	0	0	0	0	1	0	$H_{3/2}^1$

SG 196

Space group $F23$, No. 196, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	X_1	X_2	X_3	X_4
	dim	1	2	3	1	1	1	1
Δ_1	1	1	2	1	-1	-1	0	0
Δ_2	1	0	0	2	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	X_1	X_2	X_3	X_4	
	1	2	3	1	1	1	1	Bipartite graph
v^1	0	0	1	0	1	0	2	$H_{4/3}^1$
v^2	0	0	1	0	1	1	1	$H_{4/3}^1$
v^3	0	0	1	0	1	2	0	$H_{4/3}^1$
v^4	0	0	1	1	0	0	2	$H_{4/3}^1$
v^5	0	0	1	1	0	1	1	$H_{4/3}^1$
v^6	0	0	1	1	0	2	0	$H_{4/3}^1$
v^7	0	1	0	0	2	0	0	$H_{3/2}^1$
v^8	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{10}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	L_1	L_2L_3
	dim	1	2	3	1	2
Λ_1	1	1	0	1	-1	0
Λ_2	1	0	1	1	0	-1
Λ_3	1	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	L_1	L_2L_3	
	1	2	3	1	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/3}^1$
v^2	0	1	0	0	1	$H_{2/2}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q : $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		L_1	L_2L_3	W_1	W_2
	dim	1	2	1	1
Q_1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1	L_2L_3	W_1	W_2	
	1	2	1	1	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	1	$H_{2/1}^1$
v^5	1	0	1	0	$H_{2/1}^1$

SG 197

Space group $I23$, No. 197, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	N_1	N_2
dim	1	2	3	1	1
Σ_1	1	2	3	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	N_1	N_2	
	1	2	3	1	1	Bipartite graph
v^1	0	0	1	0	3	$H_{4/3}^1$
v^2	0	0	1	1	2	$H_{4/3}^1$
v^3	0	0	1	2	1	$H_{4/3}^1$
v^4	0	0	1	3	0	$H_{4/3}^1$
v^5	0	1	0	0	2	$H_{3/2}^1$
v^6	0	1	0	1	1	$H_{3/2}^1$
v^7	0	1	0	2	0	$H_{3/2}^1$
v^8	1	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	H_1	H_2H_3	H_4
dim		1	2	3	1	2	3
Δ_1	1	1	2	1	-1	-2	-1
Δ_2	1	0	0	2	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	H_1	H_2H_3	H_4	
	1	2	3	1	2	3	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/3}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	0	1	0	2	0	0	$H_{3/2}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	2	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	P_1	P_2	P_3	P_4
dim		1	2	3	1	1	1	3
Λ_1	1	1	0	1	-1	0	0	-1
Λ_2	1	0	1	1	0	0	-1	-1
Λ_3	1	0	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	P_1	P_2	P_3	P_4	
	1	2	3	1	1	1	3	Bipartite graph
v^1	0	0	1	0	0	0	1	$H_{2/3}^1$
v^2	0	0	1	1	1	1	0	$H_{4/3}^1$
v^3	0	1	0	0	1	1	0	$H_{3/2}^1$
v^4	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	0	1	$H_{3/3}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix



corep		H_1	H_2H_3	H_4	N_1	N_2
	dim	1	2	3	1	1
G_1	1	1	2	3	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	H_1	H_2H_3	H_4	N_1	N_2	Bipartite graph
	1	2	3	1	1	
v^1	0	0	1	0	3	$H_{4/3}^1$
v^2	0	0	1	1	2	$H_{4/3}^1$
v^3	0	0	1	2	1	$H_{4/3}^1$
v^4	0	0	1	3	0	$H_{4/3}^1$
v^5	0	1	0	0	2	$H_{3/2}^1$
v^6	0	1	0	1	1	$H_{3/2}^1$
v^7	0	1	0	2	0	$H_{3/2}^1$
v^8	1	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1	H_2H_3	H_4	P_1	P_2	P_3	P_4
	dim	1	2	3	1	1	1	3
F_1	1	1	0	1	-1	0	0	-1
F_2	1	0	1	1	0	0	-1	-1
F_3	1	0	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	H_1	H_2H_3	H_4	P_1	P_2	P_3	P_4	Bipartite graph
	1	2	3	1	1	1	3	
v^1	0	0	1	0	0	0	1	$H_{2/3}^1$
v^2	0	0	1	1	1	1	0	$H_{4/3}^1$
v^3	0	1	0	0	1	1	0	$H_{3/2}^1$
v^4	1	0	0	1	0	0	0	$H_{2/1}^1$
v^5	1	1	0	0	0	0	1	$H_{3/3}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_1	P_2	P_3	P_4	N_1	N_2
	dim	1	1	1	3	1	1
D_1	1	1	1	1	1	-1	0
D_2	1	0	0	0	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	P_1	P_2	P_3	P_4	N_1	N_2	
	1	1	1	3	1	1	Bipartite graph
v^1	0	0	0	1	1	2	$H_{4/3}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$

SG 198

Space group $P2_13$, No. 198, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	X_1
	dim	1	2	3	2
Δ_1	1	1	2	1	-1
Δ_2	1	0	0	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	X_1	
	1	2	3	2	Bipartite graph
v^1	0	1	2	4	$H_{7/8}^{10}, H_{7/8}^3$
v^2	1	0	1	2	$H_{4/4}^4$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	M_1M_2	M_3M_4
dim	1	2	3	2	2
Σ_1	1	1	2	3	-2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	M_1M_2	M_3M_4	Bipartite graph
	1	2	3	2	2	
v^1	0	0	2	0	3	$H_{5/6}^2$
v^2	0	0	2	1	2	$H_{5/6}^2$
v^3	0	0	2	2	1	$H_{5/6}^2$
v^4	0	0	2	3	0	$H_{5/6}^2$
v^5	0	1	0	0	1	$H_{2/2}^1$
v^6	0	1	0	1	0	$H_{2/2}^1$
v^7	1	0	1	0	2	$H_{4/4}^4$
v^8	1	0	1	1	1	$H_{4/4}^4$
v^9	1	0	1	2	0	$H_{4/4}^4$
v^{10}	2	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	R_1R_1	R_2R_3
dim	1	2	3	4	4
Λ_1	1	1	0	1	0 -2
Λ_2	1	0	1	1	-2 -1
Λ_3	1	0	1	1	-2 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	R_1R_1	R_2R_3	Bipartite graph
	1	2	3	4	4	
v^1	0	0	4	1	2	$H_{7/12}^2, H_{7/12}^1$
v^2	0	1	2	1	1	$H_{5/8}^2, H_{5/8}^3$
v^3	0	2	0	1	0	$H_{3/4}^1$
v^4	1	0	1	0	1	$H_{3/4}^2$
v^5	2	1	0	0	1	$H_{4/4}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	R_1R_1	R_2R_3
dim	2	4	4
S_1S_1	2	1	-2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	R_1R_1	R_2R_3	
	2	4	4	Bipartite graph
v^1	2	0	1	$H_{3/2}^1$
v^2	2	1	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	M_1M_2	M_3M_4
dim	2	2	2
Z_1Z_2	2	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	M_1M_2	M_3M_4	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_1M_2	M_3M_4	R_1R_1	R_2R_3
dim	2	2	4	4
T_1T_1	2	1	0	-1 -1
T_2T_2	2	0	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_1M_2	M_3M_4	R_1R_1	R_2R_3	
	2	2	4	4	Bipartite graph
v^1	1	1	0	1	$H_{3/2}^1$
v^2	1	1	1	0	$H_{3/2}^1$

SG 199

Space group $I2_13$, No. 199, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	N_1	N_2
dim	1	2	3	1	1
Σ_1	1	1	2	3	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	N_1	N_2	
	1	2	3	1	1	Bipartite graph
v^1	0	0	1	0	3	$H_{4/3}^1$
v^2	0	0	1	1	2	$H_{4/3}^1$
v^3	0	0	1	2	1	$H_{4/3}^1$
v^4	0	0	1	3	0	$H_{4/3}^1$
v^5	0	1	0	0	2	$H_{3/2}^1$
v^6	0	1	0	1	1	$H_{3/2}^1$
v^7	0	1	0	2	0	$H_{3/2}^1$
v^8	1	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	H_1	H_2H_3	H_4
dim	1	2	3	1	2	3
Δ_1	1	1	2	1	0	0 -2
Δ_2	1	0	0	2	-1	-2 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	H_1	H_2H_3	H_4	
	1	2	3	1	2	3	Bipartite graph
v^1	0	0	2	1	1	1	$H_{5/6}^5, H_{5/6}^3$
v^2	0	0	2	3	0	1	$H_{6/6}^3$
v^3	0	0	4	0	3	2	$H_{9/12}^6, H_{9/12}^7, H_{9/12}^3, H_{9/12}^2, H_{9/12}^1$
v^4	0	1	2	0	1	2	$H_{6/8}^4, H_{6/8}^5, H_{6/8}^3$
v^5	0	1	2	2	0	2	$H_{7/8}^4, H_{7/8}^5, H_{7/8}^8, H_{7/8}^6$
v^6	0	2	2	1	0	3	$H_{8/10}^3, H_{8/10}^5, H_{8/10}^4, H_{8/10}^1, H_{8/10}^6$
v^7	0	3	2	0	0	4	$H_{9/12}^6, H_{9/12}^7, H_{9/12}^3, H_{9/12}^2, H_{9/12}^1$
v^8	1	0	1	1	0	1	$H_{4/4}^3$
v^9	1	0	3	0	2	2	$H_{8/10}^3, H_{8/10}^4, H_{8/10}^6, H_{8/10}^1, H_{8/10}^5$
v^{10}	1	1	1	0	0	2	$H_{5/6}^5, H_{5/6}^3$
v^{11}	2	0	2	0	1	2	$H_{7/8}^4, H_{7/8}^8, H_{7/8}^5, H_{7/8}^6$
v^{12}	3	0	1	0	0	2	$H_{6/6}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		Γ_1	$\Gamma_2\Gamma_3$	Γ_4	P_1	P_2	P_3
dim		1	2	3	2	2	2
Λ_1	1	1	0	1	0	-1	-1
Λ_2	1	0	1	1	-1	0	-1
Λ_3	1	0	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1	$\Gamma_2\Gamma_3$	Γ_4	P_1	P_2	P_3	
	1	2	3	2	2	2	Bipartite graph
v^1	0	0	2	1	1	1	$H_{5/6}^2$
v^2	0	1	0	1	0	0	$H_{2/2}^1$
v^3	1	0	1	0	1	1	$H_{4/4}^4$
v^4	2	1	0	0	1	1	$H_{5/4}^3$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep		H_1	H_2H_3	H_4	N_1	N_2
	dim	1	2	3	1	1
G_1	1	1	2	3	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	H_1	H_2H_3	H_4	N_1	N_2	
	1	2	3	1	1	Bipartite graph
v^1	0	0	1	0	3	$H_{4/3}^1$
v^2	0	0	1	1	2	$H_{4/3}^1$
v^3	0	0	1	2	1	$H_{4/3}^1$
v^4	0	0	1	3	0	$H_{4/3}^1$
v^5	0	1	0	0	2	$H_{3/2}^1$
v^6	0	1	0	1	1	$H_{3/2}^1$
v^7	0	1	0	2	0	$H_{3/2}^1$
v^8	1	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1	H_2H_3	H_4	P_1	P_2	P_3
	dim	1	2	3	2	2	2
F_1	1	1	0	1	0	-1	-1
F_2	1	0	1	1	-1	0	-1
F_3	1	0	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	H_1	H_2H_3	H_4	P_1	P_2	P_3	
	1	2	3	2	2	2	Bipartite graph
v^1	0	0	2	1	1	1	$H_{5/6}^2$
v^2	0	1	0	1	0	0	$H_{2/2}^1$
v^3	1	0	1	0	1	1	$H_{4/4}^4$
v^4	2	1	0	0	1	1	$H_{5/4}^3$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_1	P_2	P_3	N_1	N_2
dim		2	2	2	1	1
D_1	1	1	1	1	-1	0
D_2	1	1	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_1	P_2	P_3	N_1	N_2	
	2	2	2	1	1	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$

SG 200

Space group $Pm\bar{3}$, No. 200, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_3^+	Γ_2^-	Γ_3^-	Γ_4^+	Γ_4^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
dim		1	1	2	2	3	3	1	1	1	1	1	1	1	1	1	1
Δ_1	1	1	0	2	0	0	1	-1	0	0	-1	0	0	0	0	0	0
Δ_2	1	0	0	0	0	1	1	0	0	0	0	0	0	-1	-1	0	0
Δ_3	1	0	1	0	2	1	0	0	-1	-1	0	0	0	0	0	0	0
Δ_4	1	0	0	0	0	1	1	0	0	0	0	0	-1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 26.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_3^+	Γ_2^-	Γ_3^-	Γ_4^+	Γ_4^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	
	1	1	2	2	3	3	1	1	1	1	1	1	1	1	1	1	Bipartite graph
v^1	0	0	0	0	0	1	0	0	0	1	0	0	1	1			$H_{4/3}^1$
v^2	0	0	0	0	0	1	0	0	0	1	0	1	0	1			$H_{4/3}^1$
v^3	0	0	0	0	0	1	0	0	0	1	1	0	1	0			$H_{4/3}^1$
v^4	0	0	0	0	0	1	0	0	0	1	1	1	0	0			$H_{4/3}^1$
v^5	0	0	0	0	0	1	1	0	0	0	0	0	1	1			$H_{4/3}^1$
v^6	0	0	0	0	0	1	1	0	0	0	0	1	0	1			$H_{4/3}^1$
v^7	0	0	0	0	0	1	1	0	0	0	1	0	1	0			$H_{4/3}^1$
v^8	0	0	0	0	0	1	1	0	0	0	1	1	0	0			$H_{4/3}^1$
v^9	0	0	0	0	1	0	0	0	1	0	0	0	1	1			$H_{4/3}^1$
v^{10}	0	0	0	0	1	0	0	0	1	0	0	1	0	1			$H_{4/3}^1$

v^{11}	0	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{4/3}^1$
v^{12}	0	0	0	0	1	0	0	0	1	0	1	1	0	0	$H_{4/3}^1$
v^{13}	0	0	0	0	1	0	0	1	0	0	0	0	1	1	$H_{4/3}^1$
v^{14}	0	0	0	0	1	0	0	1	0	0	0	1	0	1	$H_{4/3}^1$
v^{15}	0	0	0	0	1	0	0	1	0	0	1	0	1	0	$H_{4/3}^1$
v^{16}	0	0	0	0	1	0	0	1	0	0	1	1	0	0	$H_{4/3}^1$
v^{17}	0	0	0	1	0	0	0	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	1	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{19}	0	0	0	1	0	0	0	2	0	0	0	0	0	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	0	0	0	0	2	0	0	0	0	$H_{3/2}^1$
v^{21}	0	0	1	0	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	2	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{24}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{26}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-
	dim	1	1	2	2	3	3	1	1	1	1	1	1	1	1
Σ_1	1	1	0	2	0	1	2	-1	0	0	-1	-1	0	0	-1
Σ_2	1	0	1	0	2	2	1	0	-1	-1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 108.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	Bipartite graph
	1	1	2	2	3	3	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	0	0	0	1	2	$H_{4/3}^1$
v^2	0	0	0	0	0	1	0	0	0	0	0	1	0	2	$H_{4/3}^1$
v^3	0	0	0	0	0	1	0	0	0	0	1	0	1	1	$H_{4/3}^1$
v^4	0	0	0	0	0	1	0	0	0	0	1	1	0	1	$H_{4/3}^1$
v^5	0	0	0	0	0	1	0	0	0	0	2	0	1	0	$H_{4/3}^1$
v^6	0	0	0	0	0	1	0	0	0	0	2	1	0	0	$H_{4/3}^1$
v^7	0	0	0	0	0	1	0	0	0	1	0	0	1	1	$H_{4/3}^1$
v^8	0	0	0	0	0	1	0	0	0	1	0	1	0	1	$H_{4/3}^1$
v^9	0	0	0	0	0	1	0	0	0	1	1	0	1	0	$H_{4/3}^1$
v^{10}	0	0	0	0	0	1	0	0	0	1	1	1	0	0	$H_{4/3}^1$
v^{11}	0	0	0	0	0	1	0	0	0	2	0	0	1	0	$H_{4/3}^1$
v^{12}	0	0	0	0	0	1	0	0	0	2	0	1	0	0	$H_{4/3}^1$

v^{13}	0	0	0	0	0	1	0	0	1	0	0	0	0	2	$H_{4/3}^1$
v^{14}	0	0	0	0	0	1	0	0	1	0	1	0	0	1	$H_{4/3}^1$
v^{15}	0	0	0	0	0	1	0	0	1	0	2	0	0	0	$H_{4/3}^1$
v^{16}	0	0	0	0	0	1	0	0	1	1	0	0	0	1	$H_{4/3}^1$
v^{17}	0	0	0	0	0	1	0	0	1	1	1	0	0	0	$H_{4/3}^1$
v^{18}	0	0	0	0	0	1	0	0	1	2	0	0	0	0	$H_{4/3}^1$
v^{19}	0	0	0	0	0	1	0	1	0	0	0	0	0	2	$H_{4/3}^1$
v^{20}	0	0	0	0	0	1	0	1	0	0	1	0	0	1	$H_{4/3}^1$
v^{21}	0	0	0	0	0	1	0	1	0	0	2	0	0	0	$H_{4/3}^1$
v^{22}	0	0	0	0	0	1	0	1	0	1	0	0	0	1	$H_{4/3}^1$
v^{23}	0	0	0	0	0	1	0	1	0	1	1	0	0	0	$H_{4/3}^1$
v^{24}	0	0	0	0	0	1	0	1	0	2	0	0	0	0	$H_{4/3}^1$
v^{25}	0	0	0	0	0	1	1	0	0	0	0	0	1	1	$H_{4/3}^1$
v^{26}	0	0	0	0	0	1	1	0	0	0	0	1	0	1	$H_{4/3}^1$
v^{27}	0	0	0	0	0	1	1	0	0	0	1	0	1	0	$H_{4/3}^1$
v^{28}	0	0	0	0	0	1	1	0	0	0	1	1	0	0	$H_{4/3}^1$
v^{29}	0	0	0	0	0	1	1	0	0	1	0	0	1	0	$H_{4/3}^1$
v^{30}	0	0	0	0	0	1	1	0	0	1	0	1	0	0	$H_{4/3}^1$
v^{31}	0	0	0	0	0	1	1	0	1	0	0	0	0	1	$H_{4/3}^1$
v^{32}	0	0	0	0	0	1	1	0	1	0	1	0	0	0	$H_{4/3}^1$
v^{33}	0	0	0	0	0	1	1	0	1	1	0	0	0	0	$H_{4/3}^1$
v^{34}	0	0	0	0	0	1	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{35}	0	0	0	0	0	1	1	1	0	0	1	0	0	0	$H_{4/3}^1$
v^{36}	0	0	0	0	0	1	1	1	0	1	0	0	0	0	$H_{4/3}^1$
v^{37}	0	0	0	0	0	1	2	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{38}	0	0	0	0	0	1	2	0	0	0	0	1	0	0	$H_{4/3}^1$
v^{39}	0	0	0	0	0	1	2	0	1	0	0	0	0	0	$H_{4/3}^1$
v^{40}	0	0	0	0	0	1	2	1	0	0	0	0	0	0	$H_{4/3}^1$
v^{41}	0	0	0	0	1	0	0	0	0	0	0	0	2	1	$H_{4/3}^1$
v^{42}	0	0	0	0	1	0	0	0	0	0	0	1	1	1	$H_{4/3}^1$
v^{43}	0	0	0	0	1	0	0	0	0	0	0	2	0	1	$H_{4/3}^1$
v^{44}	0	0	0	0	1	0	0	0	0	0	1	0	2	0	$H_{4/3}^1$
v^{45}	0	0	0	0	1	0	0	0	0	0	1	1	1	0	$H_{4/3}^1$
v^{46}	0	0	0	0	1	0	0	0	0	0	1	2	0	0	$H_{4/3}^1$
v^{47}	0	0	0	0	1	0	0	0	0	1	0	0	2	0	$H_{4/3}^1$
v^{48}	0	0	0	0	1	0	0	0	0	1	0	1	1	0	$H_{4/3}^1$
v^{49}	0	0	0	0	1	0	0	0	0	1	0	2	0	0	$H_{4/3}^1$
v^{50}	0	0	0	0	1	0	0	0	1	0	0	0	1	1	$H_{4/3}^1$
v^{51}	0	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{4/3}^1$
v^{52}	0	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{4/3}^1$
v^{53}	0	0	0	0	1	0	0	0	1	0	1	1	0	0	$H_{4/3}^1$
v^{54}	0	0	0	0	1	0	0	0	1	1	0	0	1	0	$H_{4/3}^1$

v^{55}	0	0	0	0	1	0	0	0	1	1	0	1	0	0	$H_{4/3}^1$
v^{56}	0	0	0	0	1	0	0	0	2	0	0	0	0	1	$H_{4/3}^1$
v^{57}	0	0	0	0	1	0	0	0	2	0	1	0	0	0	$H_{4/3}^1$
v^{58}	0	0	0	0	1	0	0	0	2	1	0	0	0	0	$H_{4/3}^1$
v^{59}	0	0	0	0	1	0	0	1	0	0	0	0	1	1	$H_{4/3}^1$
v^{60}	0	0	0	0	1	0	0	1	0	0	0	1	0	1	$H_{4/3}^1$
v^{61}	0	0	0	0	1	0	0	1	0	0	1	0	1	0	$H_{4/3}^1$
v^{62}	0	0	0	0	1	0	0	1	0	0	1	1	0	0	$H_{4/3}^1$
v^{63}	0	0	0	0	1	0	0	1	0	1	0	0	1	0	$H_{4/3}^1$
v^{64}	0	0	0	0	1	0	0	1	0	1	0	1	0	0	$H_{4/3}^1$
v^{65}	0	0	0	0	1	0	0	1	1	0	0	0	0	1	$H_{4/3}^1$
v^{66}	0	0	0	0	1	0	0	1	1	0	1	0	0	0	$H_{4/3}^1$
v^{67}	0	0	0	0	1	0	0	1	1	1	0	0	0	0	$H_{4/3}^1$
v^{68}	0	0	0	0	1	0	0	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{69}	0	0	0	0	1	0	0	2	0	0	1	0	0	0	$H_{4/3}^1$
v^{70}	0	0	0	0	1	0	0	2	0	1	0	0	0	0	$H_{4/3}^1$
v^{71}	0	0	0	0	1	0	1	0	0	0	0	0	2	0	$H_{4/3}^1$
v^{72}	0	0	0	0	1	0	1	0	0	0	0	1	1	0	$H_{4/3}^1$
v^{73}	0	0	0	0	1	0	1	0	0	0	0	2	0	0	$H_{4/3}^1$
v^{74}	0	0	0	0	1	0	1	0	1	0	0	0	1	0	$H_{4/3}^1$
v^{75}	0	0	0	0	1	0	1	0	1	0	0	1	0	0	$H_{4/3}^1$
v^{76}	0	0	0	0	1	0	1	0	2	0	0	0	0	0	$H_{4/3}^1$
v^{77}	0	0	0	0	1	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{78}	0	0	0	0	1	0	1	1	0	0	0	1	0	0	$H_{4/3}^1$
v^{79}	0	0	0	0	1	0	1	1	1	0	0	0	0	0	$H_{4/3}^1$
v^{80}	0	0	0	0	1	0	1	2	0	0	0	0	0	0	$H_{4/3}^1$
v^{81}	0	0	0	1	0	0	0	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{82}	0	0	0	1	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{83}	0	0	0	1	0	0	0	0	0	0	0	2	0	0	$H_{3/2}^1$
v^{84}	0	0	0	1	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{85}	0	0	0	1	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{86}	0	0	0	1	0	0	0	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{87}	0	0	0	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{88}	0	0	0	1	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{89}	0	0	0	1	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{90}	0	0	0	1	0	0	0	2	0	0	0	0	0	0	$H_{3/2}^1$
v^{91}	0	0	1	0	0	0	0	0	0	0	0	0	0	2	$H_{3/2}^1$
v^{92}	0	0	1	0	0	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{93}	0	0	1	0	0	0	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{94}	0	0	1	0	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{95}	0	0	1	0	0	0	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{96}	0	0	1	0	0	0	0	0	0	2	0	0	0	0	$H_{3/2}^1$

v^{97}	0	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{98}	0	0	1	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{99}	0	0	1	0	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{100}	0	0	1	0	0	0	2	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{101}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{102}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{103}	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{104}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{105}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{106}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{107}	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{108}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	R_1^+	R_1^-	$R_2^+R_3^+$	$R_2^-R_3^-$	R_4^+	R_4^-
dim		1	1	2	2	3	3	1	1	2	2	3	3
Λ_1	1	1	1	0	0	1	1	-1	-1	0	0	-1	-1
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	1	0	0	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 28.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	R_1^+	R_1^-	$R_2^+R_3^+$	$R_2^-R_3^-$	R_4^+	R_4^-	Bipartite graph
	1	1	2	2	3	3	1	1	2	2	3	3	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^9	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

v^{17}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{20}	0	1	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{21}	0	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{22}	0	1	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{23}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{27}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{28}	1	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-
	dim	1	1	1	1	1	1	1	1	1	1	2	2	3	3
S_1	1	1	0	1	0	0	1	0	1	-1	0	-2	0	-1	-2
S_2	1	0	1	0	1	1	0	1	0	0	-1	0	-2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 108.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	2	2	3	3	
v^1	0	0	0	0	0	0	0	1	1	0	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	2	0	0	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	1	2	0	0	0	0	0	1	$H_{4/3}^1$
v^5	0	0	0	0	0	0	2	0	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	2	1	0	0	0	0	1	0	$H_{4/3}^1$
v^7	0	0	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^8	0	0	0	0	0	1	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	1	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{10}	0	0	0	0	0	1	2	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{11}	0	0	0	0	0	2	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	0	2	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{13}	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	1	0	0	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{15}	0	0	0	0	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	0	0	0	1	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$

v^{17}	0	0	0	0	1	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{18}	0	0	0	0	1	1	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{19}	0	0	0	0	1	2	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{20}	0	0	0	0	2	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{21}	0	0	0	0	2	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{22}	0	0	0	0	2	1	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{23}	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	0	0	0	1	0	0	0	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{25}	0	0	0	1	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{26}	0	0	0	1	0	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{27}	0	0	0	1	0	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{28}	0	0	0	1	0	1	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{29}	0	0	0	1	0	2	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{30}	0	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{31}	0	0	0	1	1	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{32}	0	0	0	1	1	1	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{33}	0	0	0	2	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{34}	0	0	0	2	0	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{35}	0	0	0	2	0	1	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{36}	0	0	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{37}	0	0	1	0	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{38}	0	0	1	0	0	0	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{39}	0	0	1	0	0	0	2	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{40}	0	0	1	0	0	1	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{41}	0	0	1	0	0	1	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{42}	0	0	1	0	1	0	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{43}	0	0	1	0	1	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{44}	0	0	1	0	1	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{45}	0	0	1	0	2	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{46}	0	0	1	1	0	0	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{47}	0	0	1	1	0	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{48}	0	0	1	1	0	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{49}	0	0	1	1	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{50}	0	0	1	2	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{51}	0	0	2	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{52}	0	0	2	0	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{53}	0	0	2	0	1	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{54}	0	0	2	1	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{55}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{56}	0	1	0	0	0	0	0	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{57}	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{58}	0	1	0	0	0	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$

v^{59}	0	1	0	0	0	1	0	1	0	0	0	0	1	$H_{4/3}^1$
v^{60}	0	1	0	0	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{61}	0	1	0	0	0	2	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{62}	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{63}	0	1	0	0	1	0	0	1	0	0	0	1	0	$H_{4/3}^1$
v^{64}	0	1	0	0	1	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{65}	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{66}	0	1	0	1	0	0	0	1	0	0	0	1	0	$H_{4/3}^1$
v^{67}	0	1	0	1	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{68}	0	1	1	0	0	0	0	1	0	0	0	0	1	$H_{4/3}^1$
v^{69}	0	1	1	0	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{70}	0	1	1	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{71}	0	1	1	0	1	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{72}	0	1	1	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{73}	0	1	2	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{74}	0	2	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{75}	0	2	0	0	0	0	0	1	0	0	0	1	0	$H_{4/3}^1$
v^{76}	0	2	0	0	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{77}	0	2	1	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{78}	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{79}	1	0	0	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{80}	1	0	0	0	0	0	1	1	0	0	0	0	1	$H_{4/3}^1$
v^{81}	1	0	0	0	0	0	2	0	0	0	0	1	0	$H_{4/3}^1$
v^{82}	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{83}	1	0	0	0	0	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{84}	1	0	0	0	1	0	0	1	0	0	0	0	1	$H_{4/3}^1$
v^{85}	1	0	0	0	1	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{86}	1	0	0	0	1	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{87}	1	0	0	0	2	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{88}	1	0	0	1	0	0	0	1	0	0	0	0	1	$H_{4/3}^1$
v^{89}	1	0	0	1	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{90}	1	0	0	1	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{91}	1	0	0	1	1	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{92}	1	0	0	2	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{93}	1	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{94}	1	0	1	0	0	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{95}	1	0	1	0	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{96}	1	0	1	1	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{97}	1	1	0	0	0	0	0	1	0	0	0	0	1	$H_{4/3}^1$
v^{98}	1	1	0	0	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{99}	1	1	0	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{100}	1	1	0	0	1	0	0	0	0	0	0	1	0	$H_{4/3}^1$

v^{101}	1	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{102}	1	1	1	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{103}	1	2	0	0	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{104}	2	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{105}	2	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{106}	2	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{107}	2	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{108}	2	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-
	dim	1	1	1	1	1	1	1	1	1	1	2	2	3	3
T_1	1	1	0	0	0	0	1	0	0	-1	0	-2	0	0	-1
T_2	1	0	0	1	0	0	0	0	1	0	0	0	0	-1	-1
T_3	1	0	1	0	0	1	0	0	0	0	-1	0	-2	-1	0
T_4	1	0	0	0	1	0	0	1	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 26.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-	Bipartite graph
	1	1	1	1	1	1	1	1	1	1	2	2	3	3	
v^1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^3	0	0	0	0	0	2	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	0	1	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^6	0	0	0	0	2	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	1	0	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^8	0	0	0	1	1	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^9	0	0	1	0	0	1	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{10}	0	0	1	0	1	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{11}	0	0	1	1	0	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{12}	0	0	1	1	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{15}	0	1	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	1	0	1	0	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{17}	0	1	1	0	0	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{18}	0	1	1	1	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$

v^{19}	0	2	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{20}	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{21}	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{22}	1	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{23}	1	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{24}	1	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{25}	1	0	1	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{26}	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$

SG 201

Space group $Pn\bar{3}$, No. 201, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	X_1	X_2
dim		1	1	2	2	3	3	2	2
Δ_1	1	1	0	2	0	0	1	-1	0
Δ_2	1	0	0	0	0	1	1	0	-1
Δ_3	1	0	1	0	2	1	0	-1	0
Δ_4	1	0	0	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	X_1	X_2	
	1	1	2	2	3	3	2	2	Bipartite graph
v^1	0	0	0	0	1	1	1	2	$H_{5/6}^2$
v^2	0	0	0	1	0	2	2	2	$H_{7/8}^3$
v^3	0	0	1	0	2	0	2	2	$H_{7/8}^3$
v^4	0	0	1	1	0	0	2	0	$H_{4/4}^2$
v^5	0	1	0	0	0	1	1	1	$H_{4/4}^4$
v^6	0	1	1	0	1	0	2	1	$H_{6/6}^8$
v^7	0	2	1	0	0	0	2	0	$H_{5/4}^3$
v^8	1	0	0	0	1	0	1	1	$H_{4/4}^4$
v^9	1	0	0	1	0	1	2	1	$H_{6/6}^8$
v^{10}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	1	0	0	2	0	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	M_1	M_2
	dim	1	1	2	2	3	3	2	2
Σ_1	1	1	0	2	0	1	2	-1	-1
Σ_2	1	0	1	0	2	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 39.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	M_1	M_2	Bipartite graph
	1	1	2	2	3	3	2	2	
v^1	0	0	0	0	1	1	0	3	$H_{5/6}^2$
v^2	0	0	0	0	1	1	1	2	$H_{5/6}^2$
v^3	0	0	0	0	1	1	2	1	$H_{5/6}^2$
v^4	0	0	0	0	1	1	3	0	$H_{5/6}^2$
v^5	0	0	0	1	0	2	0	4	$H_{7/8}^{10}, H_{7/8}^3$
v^6	0	0	0	1	0	2	1	3	$H_{7/8}^{10}, H_{7/8}^3$
v^7	0	0	0	1	0	2	2	2	$H_{7/8}^{10}, H_{7/8}^3$
v^8	0	0	0	1	0	2	3	1	$H_{7/8}^{10}, H_{7/8}^3$
v^9	0	0	0	1	0	2	4	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{10}	0	0	1	0	2	0	0	4	$H_{7/8}^{10}, H_{7/8}^3$
v^{11}	0	0	1	0	2	0	1	3	$H_{7/8}^{10}, H_{7/8}^3$
v^{12}	0	0	1	0	2	0	2	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{13}	0	0	1	0	2	0	3	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{14}	0	0	1	0	2	0	4	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{15}	0	0	1	1	0	0	0	2	$H_{4/4}^2$
v^{16}	0	0	1	1	0	0	1	1	$H_{4/4}^2$
v^{17}	0	0	1	1	0	0	2	0	$H_{4/4}^2$
v^{18}	0	1	0	0	0	1	0	2	$H_{4/4}^4$
v^{19}	0	1	0	0	0	1	1	1	$H_{4/4}^4$
v^{20}	0	1	0	0	0	1	2	0	$H_{4/4}^4$
v^{21}	0	1	1	0	1	0	0	3	$H_{6/6}^8, H_{6/6}^7$
v^{22}	0	1	1	0	1	0	1	2	$H_{6/6}^8, H_{6/6}^7$
v^{23}	0	1	1	0	1	0	2	1	$H_{6/6}^8, H_{6/6}^7$
v^{24}	0	1	1	0	1	0	3	0	$H_{6/6}^8, H_{6/6}^7$
v^{25}	0	2	1	0	0	0	0	2	$H_{5/4}^3$
v^{26}	0	2	1	0	0	0	1	1	$H_{5/4}^3$
v^{27}	0	2	1	0	0	0	2	0	$H_{5/4}^3$
v^{28}	1	0	0	0	1	0	0	2	$H_{4/4}^4$
v^{29}	1	0	0	0	1	0	1	1	$H_{4/4}^4$
v^{30}	1	0	0	0	1	0	2	0	$H_{4/4}^4$

v^{31}	1	0	0	1	0	1	0	3	$H_{6/6}^8, H_{6/6}^7$
v^{32}	1	0	0	1	0	1	1	2	$H_{6/6}^8, H_{6/6}^7$
v^{33}	1	0	0	1	0	1	2	1	$H_{6/6}^8, H_{6/6}^7$
v^{34}	1	0	0	1	0	1	3	0	$H_{6/6}^8, H_{6/6}^7$
v^{35}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{36}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{37}	2	0	0	1	0	0	0	2	$H_{5/4}^3$
v^{38}	2	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{39}	2	0	0	1	0	0	2	0	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	R_1^+	R_1^-	$R_2^+R_3^+$	$R_2^-R_3^-$	R_4^+	R_4^-
dim		1	1	2	2	3	3	1	1	2	2	3	3
Λ_1	1	1	1	0	0	1	1	-1	-1	0	0	-1	-1
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	1	0	0	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 28.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	R_1^+	R_1^-	$R_2^+R_3^+$	$R_2^-R_3^-$	R_4^+	R_4^-	Bipartite graph
	1	1	2	2	3	3	1	1	2	2	3	3	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^9	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

v^{20}	0	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	0	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{23}	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{27}	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{28}	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-		
dim	2	2	1	1	2	2	3	3		
S_1	1	1	1	1	0	-1	0	-2	-2	-1
S_2	1	1	1	1	-1	0	-2	0	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 39.

	X_1	X_2	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-	Bipartite graph
	2	2	1	1	2	2	3	3	
v^1	0	1	1	1	0	0	0	0	$H_{3/2}^1$
v^2	0	2	0	0	1	1	0	0	$H_{4/4}^2$
v^3	0	2	0	1	0	0	0	1	$H_{4/4}^4$
v^4	0	2	0	2	1	0	0	0	$H_{5/4}^3$
v^5	0	2	1	0	0	0	1	0	$H_{4/4}^4$
v^6	0	2	2	0	0	1	0	0	$H_{5/4}^3$
v^7	0	3	0	0	0	0	1	1	$H_{5/6}^2$
v^8	0	3	0	1	1	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^9	0	3	1	0	0	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{10}	0	4	0	0	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{11}	0	4	0	0	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{12}	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{13}	1	1	0	0	1	1	0	0	$H_{4/4}^2$
v^{14}	1	1	0	1	0	0	0	1	$H_{4/4}^4$
v^{15}	1	1	0	2	1	0	0	0	$H_{5/4}^3$
v^{16}	1	1	1	0	0	0	1	0	$H_{4/4}^4$
v^{17}	1	1	2	0	0	1	0	0	$H_{5/4}^3$
v^{18}	1	2	0	0	0	0	1	1	$H_{5/6}^2$
v^{19}	1	2	0	1	1	0	1	0	$H_{6/6}^8, H_{6/6}^7$

v^{20}	1	2	1	0	0	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{21}	1	3	0	0	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{22}	1	3	0	0	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{23}	2	0	0	0	1	1	0	0	$H_{4/4}^2$
v^{24}	2	0	0	1	0	0	0	1	$H_{4/4}^4$
v^{25}	2	0	0	2	1	0	0	0	$H_{5/4}^3$
v^{26}	2	0	1	0	0	0	1	0	$H_{4/4}^4$
v^{27}	2	0	2	0	0	1	0	0	$H_{5/4}^3$
v^{28}	2	1	0	0	0	0	1	1	$H_{5/6}^2$
v^{29}	2	1	0	1	1	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{30}	2	1	1	0	0	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{31}	2	2	0	0	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{32}	2	2	0	0	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{33}	3	0	0	0	0	0	1	1	$H_{5/6}^2$
v^{34}	3	0	0	1	1	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{35}	3	0	1	0	0	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{36}	3	1	0	0	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{37}	3	1	0	0	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{38}	4	0	0	0	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{39}	4	0	0	0	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	M_1	M_2
dim	2	2	2	2
Z_1	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_1	X_2	M_1	M_2	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_1	M_2	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-
dim	2	2	1	1	2	2	3	3
T_1	1	1	0	-1	0	-2	-1	0
T_2	1	1	0	0	-2	0	0	-1
T_3	1	0	1	0	0	0	-1	-1
T_4	1	0	1	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	M_1	M_2	R_1^+	R_1^-	$R_2^+ R_3^+$	$R_2^- R_3^-$	R_4^+	R_4^-	Bipartite graph
	2	2	1	1	2	2	3	3	
v^1	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^2	1	1	0	1	0	0	0	1	$H_{4/4}^4$
v^3	1	1	1	0	0	0	1	0	$H_{4/4}^4$
v^4	1	2	0	0	0	0	1	1	$H_{5/6}^2$
v^5	2	0	0	0	1	1	0	0	$H_{4/4}^2$
v^6	2	0	0	2	1	0	0	0	$H_{5/4}^3$
v^7	2	0	2	0	0	1	0	0	$H_{5/4}^3$
v^8	2	1	0	1	1	0	1	0	$H_{6/6}^8$
v^9	2	1	1	0	0	1	0	1	$H_{6/6}^8$
v^{10}	2	2	0	0	0	1	0	2	$H_{7/8}^3$
v^{11}	2	2	0	0	1	0	2	0	$H_{7/8}^3$

SG 202

Space group $Fm\bar{3}$, No. 202, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+ \Gamma_3^+$	$\Gamma_2^- \Gamma_3^-$	Γ_4^+	Γ_4^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-
dim	1	1	2	2	3	3	1	1	1	1	1	1	1	1
Δ_1	1	1	0	2	0	0	1	-1	0	0	-1	0	0	0
Δ_2	1	0	0	0	0	1	1	0	0	0	0	0	-1	-1
Δ_3	1	0	1	0	2	1	0	0	-1	-1	0	0	0	0
Δ_4	1	0	0	0	0	1	1	0	0	0	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 26.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	Bipartite graph
	1	1	2	2	3	3	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	0	1	0	0	1	1	$H_{4/3}^1$
v^2	0	0	0	0	0	1	0	0	0	1	0	1	0	1	$H_{4/3}^1$
v^3	0	0	0	0	0	1	0	0	0	1	1	0	1	0	$H_{4/3}^1$
v^4	0	0	0	0	0	1	0	0	0	1	1	1	0	0	$H_{4/3}^1$
v^5	0	0	0	0	0	1	1	0	0	0	0	0	1	1	$H_{4/3}^1$
v^6	0	0	0	0	0	1	1	0	0	0	0	1	0	1	$H_{4/3}^1$
v^7	0	0	0	0	0	1	1	0	0	0	1	0	1	0	$H_{4/3}^1$
v^8	0	0	0	0	0	1	1	0	0	0	1	1	0	0	$H_{4/3}^1$
v^9	0	0	0	0	1	0	0	0	1	0	0	0	1	1	$H_{4/3}^1$
v^{10}	0	0	0	0	1	0	0	0	1	0	0	1	0	1	$H_{4/3}^1$
v^{11}	0	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{4/3}^1$
v^{12}	0	0	0	0	1	0	0	0	1	0	1	1	0	0	$H_{4/3}^1$
v^{13}	0	0	0	0	1	0	0	1	0	0	0	0	1	1	$H_{4/3}^1$
v^{14}	0	0	0	0	1	0	0	1	0	0	0	1	0	1	$H_{4/3}^1$
v^{15}	0	0	0	0	1	0	0	1	0	0	1	0	1	0	$H_{4/3}^1$
v^{16}	0	0	0	0	1	0	0	1	0	0	1	1	0	0	$H_{4/3}^1$
v^{17}	0	0	0	1	0	0	0	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	1	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{19}	0	0	0	1	0	0	0	2	0	0	0	0	0	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	0	0	0	0	2	0	0	0	0	$H_{3/2}^1$
v^{21}	0	0	1	0	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	0	2	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{24}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{25}	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{26}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	L_1^+	L_1^-	$L_2^+L_3^+$	$L_2^-L_3^-$
	dim	1	1	2	2	3	3	1	1	2	2
Λ_1	1	1	1	0	0	1	1	-1	-1	0	0
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	L_1^+	L_1^-	$L_2^+L_3^+$	$L_2^-L_3^-$	
	1	1	2	2	3	3	1	1	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep	L_1^+	L_1^-	$L_2^+L_3^+$	$L_2^-L_3^-$	W_1	W_2	W_3	W_4
dim	1	1	2	2	1	1	1	1
Q_1	1	1	2	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 28.

	L_1^+	L_1^-	$L_2^+L_3^+$	$L_2^-L_3^-$	W_1	W_2	W_3	W_4	
	1	1	2	2	1	1	1	1	Bipartite graph
v^1	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	1	2	0	0	0	$H_{3/2}^1$

v^{11}	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^{12}	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^{13}	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^{14}	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^{15}	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^{16}	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^{17}	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^{19}	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^{20}	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{23}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{24}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{26}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{28}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 203

Space group $Fd\bar{3}$, No. 203, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	X_1	X_2
dim	1	1	2	2	3	3	2	2
Δ_1	1	0	2	0	0	1	-1	0
Δ_2	1	0	0	0	1	1	0	-1
Δ_3	1	0	1	0	2	1	-1	0
Δ_4	1	0	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	X_1	X_2	
	1	1	2	2	3	3	2	2	Bipartite graph
v^1	0	0	0	0	1	1	1	2	$H_{5/6}^2$
v^2	0	0	0	1	0	2	2	2	$H_{7/8}^3$
v^3	0	0	1	0	2	0	2	2	$H_{7/8}^3$
v^4	0	0	1	1	0	0	2	0	$H_{4/4}^2$

v^5	0	1	0	0	0	1	1	1	$H_{4/4}^4$
v^6	0	1	1	0	1	0	2	1	$H_{6/6}^8$
v^7	0	2	1	0	0	0	2	0	$H_{5/4}^3$
v^8	1	0	0	0	1	0	1	1	$H_{4/4}^4$
v^9	1	0	0	1	0	1	2	1	$H_{6/6}^8$
v^{10}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	1	0	0	2	0	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	L_1^+	L_1^-	$L_2^+L_3^+$	$L_2^-L_3^-$
	dim	1	1	2	2	3	3	1	1	2	2
Λ_1	1	1	1	0	0	1	1	-1	-1	0	0
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	L_1^+	L_1^-	$L_2^+L_3^+$	$L_2^-L_3^-$	Bipartite graph
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_1	X_2	W_1
dim	2	2	2
Z_1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_1	X_2	W_1	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep	L_1^+	L_1^-	$L_2^+ L_3^+$	$L_2^- L_3^-$	W_1
dim	1	1	2	2	2
Q_1	1	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1^+	L_1^-	$L_2^+ L_3^+$	$L_2^- L_3^-$	W_1	
	1	1	2	2	2	Bipartite graph
v^1	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	0	1	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$
v^5	2	0	0	0	1	$H_{3/2}^1$

SG 204

Space group $Im\bar{3}$, No. 204, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+ \Gamma_3^+$	$\Gamma_2^- \Gamma_3^-$	Γ_4^+	Γ_4^-	N_1^+	N_1^-	N_2^+	N_2^-
dim	1	1	2	2	3	3	1	1	1	1
Σ_1	1	1	0	2	0	1	2	-1	0	-1
Σ_2	1	0	1	0	2	2	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 22.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	N_1^+	N_1^-	N_2^+	N_2^-	Bipartite graph
	1	1	2	2	3	3	1	1	1	1	
v^1	0	0	0	0	0	1	0	0	1	2	$H_{4/3}^1$
v^2	0	0	0	0	0	1	0	1	0	2	$H_{4/3}^1$
v^3	0	0	0	0	0	1	1	0	1	1	$H_{4/3}^1$
v^4	0	0	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v^5	0	0	0	0	0	1	2	0	1	0	$H_{4/3}^1$
v^6	0	0	0	0	0	1	2	1	0	0	$H_{4/3}^1$
v^7	0	0	0	0	1	0	0	0	2	1	$H_{4/3}^1$
v^8	0	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v^9	0	0	0	0	1	0	0	2	0	1	$H_{4/3}^1$
v^{10}	0	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$
v^{11}	0	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^{12}	0	0	0	0	1	0	1	2	0	0	$H_{4/3}^1$
v^{13}	0	0	0	1	0	0	0	0	2	0	$H_{3/2}^1$
v^{14}	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^{15}	0	0	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^{16}	0	0	1	0	0	0	0	0	0	2	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{19}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{20}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{21}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-	
dim	1	1	2	2	3	3	1	1	2	2	3	3	
Δ_1	1	1	0	2	0	0	1	-1	0	-2	0	0	-1
Δ_2	1	0	0	0	0	1	1	0	0	0	0	-1	-1
Δ_3	1	0	1	0	2	1	0	0	-1	0	-2	-1	0
Δ_4	1	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-	Bipartite graph
	1	1	2	2	3	3	1	1	2	2	3	3	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/3}^1$
v^2	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/3}^1$
v^3	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^4	0	0	0	1	0	0	0	2	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	1	1	1	0	0	1	0	$H_{5/5}^1$
v^6	0	0	0	1	0	2	0	0	1	0	2	0	$H_{6/8}^3$
v^7	0	0	0	1	0	2	2	0	0	0	2	0	$H_{7/8}^6$
v^8	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/2}^1$
v^9	0	0	1	0	0	0	2	0	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	1	0	1	0	1	1	0	0	0	1	$H_{5/5}^1$
v^{11}	0	0	1	0	2	0	0	0	0	1	0	2	$H_{6/8}^3$
v^{12}	0	0	1	0	2	0	0	2	0	0	0	2	$H_{7/8}^6$
v^{13}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	1	1	0	0	0	1	0	$H_{4/4}^3$
v^{15}	0	1	1	0	1	0	1	0	0	1	0	1	$H_{6/6}^6$
v^{16}	0	2	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{17}	0	2	0	0	0	2	0	0	1	0	2	0	$H_{7/8}^6$
v^{18}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	1	0	0	1	0	0	0	1	$H_{4/4}^3$
v^{20}	1	0	0	1	0	1	0	1	1	0	1	0	$H_{6/6}^6$
v^{21}	1	1	0	0	0	1	0	0	1	0	1	0	$H_{5/5}^1$
v^{22}	1	1	0	0	1	0	0	0	0	1	0	1	$H_{5/5}^1$
v^{23}	2	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{24}	2	0	0	0	2	0	0	0	0	1	0	2	$H_{7/8}^6$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	P_1	P_2P_3	P_4
dim	1	1	2	2	3	3	1	2	3
Λ_1	1	1	0	0	1	1	-1	0	-1
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	P_1	P_2P_3	P_4	Bipartite graph
1	1	2	2	3	3	1	2	3	

v ¹	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v ²	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v ³	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v ⁴	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v ⁵	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v ⁶	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v ⁷	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v ⁸	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v ⁹	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v ¹⁰	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v ¹¹	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v ¹²	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep		H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-	N_1^+	N_1^-	N_2^+	N_2^-
	dim	1	1	2	2	3	3	1	1	1	1
G_1	1	1	0	2	0	1	2	-1	0	0	-1
G_2	1	0	1	0	2	2	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 22.

	H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-	N_1^+	N_1^-	N_2^+	N_2^-	Bipartite graph
	1	1	2	2	3	3	1	1	1	1	
v ¹	0	0	0	0	0	1	0	0	1	2	$H_{4/3}^1$
v ²	0	0	0	0	0	1	0	1	0	2	$H_{4/3}^1$
v ³	0	0	0	0	0	1	1	0	1	1	$H_{4/3}^1$
v ⁴	0	0	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v ⁵	0	0	0	0	0	1	2	0	1	0	$H_{4/3}^1$
v ⁶	0	0	0	0	0	1	2	1	0	0	$H_{4/3}^1$
v ⁷	0	0	0	0	1	0	0	0	2	1	$H_{4/3}^1$
v ⁸	0	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v ⁹	0	0	0	0	1	0	0	2	0	1	$H_{4/3}^1$
v ¹⁰	0	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$
v ¹¹	0	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v ¹²	0	0	0	0	1	0	1	2	0	0	$H_{4/3}^1$
v ¹³	0	0	0	1	0	0	0	0	2	0	$H_{3/2}^1$
v ¹⁴	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v ¹⁵	0	0	0	1	0	0	0	2	0	0	$H_{3/2}^1$

v^{16}	0	0	1	0	0	0	0	0	0	0	2	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{19}	0	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{20}	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{21}	1	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-	P_1	P_2P_3	P_4
	dim	1	1	2	2	3	3	1	2	3
F_1	1	1	1	0	0	1	1	-1	0	-1
F_2F_3	2	0	0	1	1	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-	P_1	P_2P_3	P_4	
	1	1	2	2	3	3	1	2	3	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep	P_1	P_2P_3	P_4	N_1^+	N_1^-	N_2^+	N_2^-
dim	1	2	3	1	1	1	1

D_1	1	1	2	1	-1	-1	0	0
D_2	1	0	0	2	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	P_1	$P_2 P_3$	P_4	N_1^+	N_1^-	N_2^+	N_2^-	Bipartite graph
	1	2	3	1	1	1	1	
v^1	0	0	1	0	1	0	2	$H_{4/3}^1$
v^2	0	0	1	0	1	1	1	$H_{4/3}^1$
v^3	0	0	1	0	1	2	0	$H_{4/3}^1$
v^4	0	0	1	1	0	0	2	$H_{4/3}^1$
v^5	0	0	1	1	0	1	1	$H_{4/3}^1$
v^6	0	0	1	1	0	2	0	$H_{4/3}^1$
v^7	0	1	0	0	2	0	0	$H_{3/2}^1$
v^8	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{10}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	1	0	0	0	$H_{2/1}^1$

SG 205

Space group $Pa\bar{3}$, No. 205, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+ \Gamma_3^+$	$\Gamma_2^- \Gamma_3^-$	Γ_4^+	Γ_4^-	X_1	X_2
dim		1	1	2	2	3	3	2	2
Δ_1	1	1	0	2	0	0	1	-1	0
Δ_2	1	0	0	0	0	1	1	0	-1
Δ_3	1	0	1	0	2	1	0	0	-1
Δ_4	1	0	0	0	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_1^+	Γ_1^-	$\Gamma_2^+ \Gamma_3^+$	$\Gamma_2^- \Gamma_3^-$	Γ_4^+	Γ_4^-	X_1	X_2	Bipartite graph
	1	1	2	2	3	3	2	2	
v^1	0	0	0	1	0	2	2	2	$H_{7/8}^3$
v^2	0	0	1	0	2	0	2	2	$H_{7/8}^3$
v^3	0	1	0	0	0	1	1	1	$H_{4/4}^4$
v^4	1	0	0	0	1	0	1	1	$H_{4/4}^4$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	M_1M_2	
dim	1	1	2	2	3	3	4	
Σ_1	1	1	0	2	0	1	2	-2
Σ_2	1	0	1	0	2	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	M_1M_2	Bipartite graph
	1	1	2	2	3	3	4	
v^1	0	0	0	0	2	2	3	$H_{7/12}^1$
v^2	0	0	0	1	0	2	2	$H_{5/8}^2, H_{5/8}^1$
v^3	0	0	1	0	2	0	2	$H_{5/8}^2, H_{5/8}^1$
v^4	0	0	1	1	0	0	1	$H_{3/4}^1$
v^5	0	1	0	0	0	1	1	$H_{3/4}^2$
v^6	0	2	1	0	0	0	1	$H_{4/4}^1$
v^7	1	0	0	0	1	0	1	$H_{3/4}^2$
v^8	2	0	0	1	0	0	1	$H_{4/4}^1$
v^9	2	2	0	0	0	0	1	$H_{5/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$R_1^+R_1^+$	$R_1^-R_1^-$	$R_2^+R_3^+$	$R_2^-R_3^-$
dim	1	1	2	2	3	3	4	4	4	4
Λ_1	1	1	0	0	1	1	0	0	-2	-2
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	-2	-2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 80.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	$R_1^+R_1^+$	$R_1^-R_1^-$	$R_2^+R_3^+$	$R_2^-R_3^-$	Bipartite graph
	1	1	2	2	3	3	4	4	4	4	
v^1	0	0	0	0	0	4	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^2	0	0	0	0	0	4	0	1	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^3	0	0	0	0	0	4	0	1	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^4	0	0	0	0	0	4	1	0	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^5	0	0	0	0	0	4	1	0	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^6	0	0	0	0	0	4	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$

v^7	0	0	0	0	1	3	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^8	0	0	0	0	1	3	0	1	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^9	0	0	0	0	1	3	0	1	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{10}	0	0	0	0	1	3	1	0	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{11}	0	0	0	0	1	3	1	0	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{12}	0	0	0	0	1	3	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{13}	0	0	0	0	2	2	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{14}	0	0	0	0	2	2	0	1	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{15}	0	0	0	0	2	2	0	1	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{16}	0	0	0	0	2	2	1	0	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{17}	0	0	0	0	2	2	1	0	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{18}	0	0	0	0	2	2	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{19}	0	0	0	0	3	1	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{20}	0	0	0	0	3	1	0	1	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{21}	0	0	0	0	3	1	0	1	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{22}	0	0	0	0	3	1	1	0	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{23}	0	0	0	0	3	1	1	0	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{24}	0	0	0	0	3	1	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{25}	0	0	0	0	4	0	0	1	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{26}	0	0	0	0	4	0	0	1	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{27}	0	0	0	0	4	0	0	1	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{28}	0	0	0	0	4	0	1	0	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{29}	0	0	0	0	4	0	1	0	1	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{30}	0	0	0	0	4	0	1	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{31}	0	0	0	1	0	2	0	1	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{32}	0	0	0	1	0	2	0	1	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{33}	0	0	0	1	0	2	1	0	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{34}	0	0	0	1	0	2	1	0	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{35}	0	0	0	1	1	1	0	1	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{36}	0	0	0	1	1	1	0	1	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{37}	0	0	0	1	1	1	1	0	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{38}	0	0	0	1	1	1	1	0	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{39}	0	0	0	1	2	0	0	1	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{40}	0	0	0	1	2	0	0	1	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{41}	0	0	0	1	2	0	1	0	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{42}	0	0	0	1	2	0	1	0	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{43}	0	0	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{44}	0	0	0	2	0	0	1	0	0	0	$H_{3/2}^1$
v^{45}	0	0	1	0	0	2	0	1	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{46}	0	0	1	0	0	2	0	1	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{47}	0	0	1	0	0	2	1	0	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{48}	0	0	1	0	0	2	1	0	1	0	$H_{5/5}^3, H_{5/5}^2$

v^{49}	0	0	1	0	1	1	0	1	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{50}	0	0	1	0	1	1	0	1	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{51}	0	0	1	0	1	1	1	0	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{52}	0	0	1	0	1	1	1	0	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{53}	0	0	1	0	2	0	0	1	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{54}	0	0	1	0	2	0	0	1	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{55}	0	0	1	0	2	0	1	0	0	1	$H_{5/5}^3, H_{5/5}^2$
v^{56}	0	0	1	0	2	0	1	0	1	0	$H_{5/5}^3, H_{5/5}^2$
v^{57}	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{58}	0	0	1	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{59}	0	0	2	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{60}	0	0	2	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{61}	0	1	0	0	0	1	0	0	0	1	$H_{3/3}^1$
v^{62}	0	1	0	0	0	1	0	0	1	0	$H_{3/3}^1$
v^{63}	0	1	0	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{64}	0	1	0	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{65}	0	2	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{66}	0	2	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{67}	0	2	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{68}	0	2	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{69}	1	0	0	0	0	1	0	0	0	1	$H_{3/3}^1$
v^{70}	1	0	0	0	0	1	0	0	1	0	$H_{3/3}^1$
v^{71}	1	0	0	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{72}	1	0	0	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{73}	1	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{74}	1	1	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{75}	1	1	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{76}	1	1	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{77}	2	0	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{78}	2	0	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{79}	2	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{80}	2	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1	X_2	$R_1^+ R_1^+$	$R_1^- R_1^-$	$R_2^+ R_3^+$	$R_2^- R_3^-$
dim	2	2	4	4	4	4
$S_1 S_2$	2	1	1	-2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	$R_1^+ R_1^+$	$R_1^- R_1^-$	$R_2^+ R_3^+$	$R_2^- R_3^-$	
	2	2	4	4	4	4	Bipartite graph
v^1	0	2	0	0	0	1	$H_{3/2}^1$
v^2	0	2	0	0	1	0	$H_{3/2}^1$
v^3	0	2	0	1	0	0	$H_{3/2}^1$
v^4	0	2	1	0	0	0	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$
v^6	1	1	0	0	1	0	$H_{3/2}^1$
v^7	1	1	0	1	0	0	$H_{3/2}^1$
v^8	1	1	1	0	0	0	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	0	$H_{3/2}^1$
v^{11}	2	0	0	1	0	0	$H_{3/2}^1$
v^{12}	2	0	1	0	0	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	$M_1 M_2$
dim	2	2	4
$Z_1 Z_2$	2	1	0
$Z_3 Z_4$	2	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_1	X_2	$M_1 M_2$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$M_1 M_2$	$R_1^+ R_1^+$	$R_1^- R_1^-$	$R_2^+ R_3^+$	$R_2^- R_3^-$
dim	4	4	4	4	4
$T_1 T_1$	4	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_1 M_2$	$R_1^+ R_1^+$	$R_1^- R_1^-$	$R_2^+ R_3^+$	$R_2^- R_3^-$	
	4	4	4	4	4	Bipartite graph
v^1	1	0	0	0	1	$H_{2/1}^1$
v^2	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	1	0	0	$H_{2/1}^1$
v^4	1	1	0	0	0	$H_{2/1}^1$

SG 206

Space group $Ia\bar{3}$, No. 206, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+ \Gamma_3^+$	$\Gamma_2^- \Gamma_3^-$	Γ_4^+	Γ_4^-	N_1
	dim	1	1	2	2	3	3	2
Σ_1	1	1	0	2	0	1	2	-1
Σ_2	1	0	1	0	2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1^+	Γ_1^-	$\Gamma_2^+ \Gamma_3^+$	$\Gamma_2^- \Gamma_3^-$	Γ_4^+	Γ_4^-	N_1	
	1	1	2	2	3	3	2	Bipartite graph
v^1	0	0	0	0	1	1	3	$H_{5/6}^2$
v^2	0	0	0	1	0	2	4	$H_{7/8}^{10}, H_{7/8}^3$
v^3	0	0	1	0	2	0	4	$H_{7/8}^{10}, H_{7/8}^3$
v^4	0	0	1	1	0	0	2	$H_{4/4}^2$
v^5	0	1	0	0	0	1	2	$H_{4/4}^4$
v^6	0	1	1	0	1	0	3	$H_{6/6}^8, H_{6/6}^7$
v^7	0	2	1	0	0	0	2	$H_{5/4}^3$
v^8	1	0	0	0	1	0	2	$H_{4/4}^4$
v^9	1	0	0	1	0	1	3	$H_{6/6}^8, H_{6/6}^7$
v^{10}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	1	0	0	2	$H_{5/4}^3$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-
dim		1	1	2	2	3	3	1	1	2	2	3	3
Δ_1	1	1	0	2	0	0	1	0	0	0	0	-1	-1
Δ_2	1	0	0	0	0	1	1	0	-1	0	-2	-1	0
Δ_3	1	0	1	0	2	1	0	0	0	0	0	-1	-1
Δ_4	1	0	0	0	0	1	1	-1	0	-2	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 44.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	H_1^+	H_1^-	$H_2^+H_3^+$	$H_2^-H_3^-$	H_4^+	H_4^-	Bipartite graph
	1	1	2	2	3	3	1	1	2	2	3	3	
v^1	0	0	0	0	1	1	0	1	1	0	1	0	$H_{5/6}^5$
v^2	0	0	0	0	1	1	1	0	0	1	0	1	$H_{5/6}^5$
v^3	0	0	0	0	1	1	1	2	0	0	0	1	$H_{6/6}^3$
v^4	0	0	0	0	1	1	2	1	0	0	1	0	$H_{6/6}^3$
v^5	0	0	0	0	2	2	0	0	1	2	0	2	$H_{9/12}^{10}, H_{9/12}^8, H_{9/12}^9, H_{9/12}^1$
v^6	0	0	0	0	2	2	0	0	2	1	2	0	$H_{9/12}^{10}, H_{9/12}^8, H_{9/12}^9, H_{9/12}^1$
v^7	0	0	0	0	2	2	0	2	1	1	0	2	$H_{10/12}^7, H_{10/12}^6, H_{10/12}^1, H_{10/12}^5, H_{10/12}^3, H_{10/12}^2, H_{10/12}^4$
v^8	0	0	0	0	2	2	0	4	1	0	0	2	$H_{11/12}^1, H_{11/12}^2$
v^9	0	0	0	0	2	2	2	0	1	1	2	0	$H_{10/12}^7, H_{10/12}^6, H_{10/12}^1, H_{10/12}^5, H_{10/12}^3, H_{10/12}^2, H_{10/12}^4$
v^{10}	0	0	0	0	2	2	4	0	0	1	2	0	$H_{11/12}^1, H_{11/12}^2$
v^{11}	0	0	0	1	0	2	0	0	0	1	0	2	$H_{6/8}^3$
v^{12}	0	0	0	1	0	2	0	0	1	0	2	0	$H_{6/8}^3$
v^{13}	0	0	0	1	0	2	0	2	0	0	0	2	$H_{7/8}^6$
v^{14}	0	0	0	1	0	2	1	1	0	0	1	1	$H_{7/8}^8, H_{7/8}^6, H_{7/8}^7$
v^{15}	0	0	0	1	0	2	2	0	0	0	2	0	$H_{7/8}^6$
v^{16}	0	0	1	0	2	0	0	0	0	1	0	2	$H_{6/8}^3$
v^{17}	0	0	1	0	2	0	0	0	1	0	2	0	$H_{6/8}^3$
v^{18}	0	0	1	0	2	0	0	2	0	0	0	2	$H_{7/8}^6$
v^{19}	0	0	1	0	2	0	1	1	0	0	1	1	$H_{7/8}^8, H_{7/8}^6, H_{7/8}^7$
v^{20}	0	0	1	0	2	0	2	0	0	0	2	0	$H_{7/8}^6$
v^{21}	0	0	1	1	1	1	0	1	0	0	1	2	$H_{8/10}^9, H_{8/10}^8, H_{8/10}^7, H_{8/10}^6, H_{8/10}^{10}$
v^{22}	0	0	1	1	1	1	1	0	0	0	2	1	$H_{8/10}^9, H_{8/10}^8, H_{8/10}^6, H_{8/10}^7, H_{8/10}^{10}$
v^{23}	0	0	1	2	0	2	0	0	0	0	2	2	$H_{9/12}^{10}, H_{9/12}^9, H_{9/12}^8, H_{9/12}^1$
v^{24}	0	0	2	1	2	0	0	0	0	0	2	2	$H_{9/12}^{10}, H_{9/12}^9, H_{9/12}^8, H_{9/12}^1$
v^{25}	0	1	0	0	0	1	0	1	0	0	0	1	$H_{4/4}^3$
v^{26}	0	1	0	0	0	1	1	0	0	0	1	0	$H_{4/4}^3$
v^{27}	0	1	0	0	1	2	0	0	1	1	1	1	$H_{8/10}^9, H_{8/10}^7, H_{8/10}^6, H_{8/10}^{10}, H_{8/10}^8$
v^{28}	0	1	1	0	1	0	0	0	0	0	1	1	$H_{5/6}^5$
v^{29}	0	2	0	0	0	2	0	0	0	1	0	2	$H_{7/8}^6$
v^{30}	0	2	0	0	0	2	0	0	1	0	2	0	$H_{7/8}^6$
v^{31}	0	2	1	1	0	2	0	0	0	0	2	2	$H_{10/12}^7, H_{10/12}^5, H_{10/12}^6, H_{10/12}^3, H_{10/12}^1, H_{10/12}^2, H_{10/12}^4$

v^{32}	0	4	1	0	0	2	0	0	0	0	2	2	$H_{11/12}^1, H_{11/12}^2$
v^{33}	1	0	0	0	1	0	0	1	0	0	0	1	$H_{4/4}^3$
v^{34}	1	0	0	0	1	0	1	0	0	0	1	0	$H_{4/4}^3$
v^{35}	1	0	0	0	2	1	0	0	1	1	1	1	$H_{8/10}^9, H_{8/10}^6, H_{8/10}^{10}, H_{8/10}^8, H_{8/10}^7$
v^{36}	1	0	0	1	0	1	0	0	0	0	1	1	$H_{5/6}^5$
v^{37}	1	1	0	0	1	1	0	0	0	1	0	2	$H_{7/8}^8, H_{7/8}^6, H_{7/8}^7$
v^{38}	1	1	0	0	1	1	0	0	1	0	2	0	$H_{7/8}^8, H_{7/8}^6, H_{7/8}^7$
v^{39}	1	2	0	0	0	1	0	0	0	0	1	1	$H_{6/6}^3$
v^{40}	2	0	0	0	2	0	0	0	0	1	0	2	$H_{7/8}^6$
v^{41}	2	0	0	0	2	0	0	0	1	0	2	0	$H_{7/8}^6$
v^{42}	2	0	1	1	2	0	0	0	0	0	2	2	$H_{10/12}^7, H_{10/12}^5, H_{10/12}^6, H_{10/12}^3, H_{10/12}^1, H_{10/12}^2, H_{10/12}^4$
v^{43}	2	1	0	0	1	0	0	0	0	0	1	1	$H_{6/6}^3$
v^{44}	4	0	0	1	2	0	0	0	0	0	2	2	$H_{11/12}^1, H_{11/12}^2$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	P_1P_1	P_2P_3
	dim	1	1	2	2	3	3	4	4
Λ_1	1	1	1	0	0	1	1	0	-2
$\Lambda_2\Lambda_3$	2	0	0	1	1	1	1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	Γ_1^+	Γ_1^-	$\Gamma_2^+\Gamma_3^+$	$\Gamma_2^-\Gamma_3^-$	Γ_4^+	Γ_4^-	P_1P_1	P_2P_3	
	1	1	2	2	3	3	4	4	Bipartite graph
v^1	0	0	0	0	0	4	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^2	0	0	0	0	1	3	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^3	0	0	0	0	2	2	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^4	0	0	0	0	3	1	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^5	0	0	0	0	4	0	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^6	0	0	0	1	0	2	1	1	$H_{5/5}^3, H_{5/5}^2$
v^7	0	0	0	1	1	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^8	0	0	0	1	2	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^9	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^{10}	0	0	1	0	0	2	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{11}	0	0	1	0	1	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{12}	0	0	1	0	2	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{13}	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	0	1	0	1	$H_{3/3}^1$

v^{16}	0	1	0	0	1	0	0	1	$H_{3/3}^1$
v^{17}	0	2	0	1	0	0	0	1	$H_{4/3}^1$
v^{18}	0	2	1	0	0	0	0	1	$H_{4/3}^1$
v^{19}	1	0	0	0	0	1	0	1	$H_{3/3}^1$
v^{20}	1	0	0	0	1	0	0	1	$H_{3/3}^1$
v^{21}	1	1	0	1	0	0	0	1	$H_{4/3}^1$
v^{22}	1	1	1	0	0	0	0	1	$H_{4/3}^1$
v^{23}	2	0	0	1	0	0	0	1	$H_{4/3}^1$
v^{24}	2	0	1	0	0	0	0	1	$H_{4/3}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep		H_1^+	H_1^-	$H_2^+ H_3^+$	$H_2^- H_3^-$	H_4^+	H_4^-	N_1
	dim	1	1	2	2	3	3	2
G_1	1	1	0	2	0	1	2	-1
G_2	1	0	1	0	2	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	H_1^+	H_1^-	$H_2^+ H_3^+$	$H_2^- H_3^-$	H_4^+	H_4^-	N_1	
	1	1	2	2	3	3	2	Bipartite graph
v^1	0	0	0	0	1	1	3	$H_{5/6}^2$
v^2	0	0	0	1	0	2	4	$H_{7/8}^{10}, H_{7/8}^3$
v^3	0	0	1	0	2	0	4	$H_{7/8}^{10}, H_{7/8}^3$
v^4	0	0	1	1	0	0	2	$H_{4/4}^2$
v^5	0	1	0	0	0	1	2	$H_{4/4}^4$
v^6	0	1	1	0	1	0	3	$H_{6/6}^8, H_{6/6}^7$
v^7	0	2	1	0	0	0	2	$H_{5/4}^3$
v^8	1	0	0	0	1	0	2	$H_{4/4}^4$
v^9	1	0	0	1	0	1	3	$H_{6/6}^8, H_{6/6}^7$
v^{10}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	1	0	0	2	$H_{5/4}^3$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		H_1^+	H_1^-	$H_2^+ H_3^+$	$H_2^- H_3^-$	H_4^+	H_4^-	$P_1 P_1$	$P_2 P_3$
	dim	1	1	2	2	3	3	4	4
F_1	1	1	1	0	0	1	1	0	-2
$F_2 F_3$	2	0	0	1	1	1	1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	H_1^+	H_1^-	$H_2^+ H_3^+$	$H_2^- H_3^-$	H_4^+	H_4^-	$P_1 P_1$	$P_2 P_3$	
	1	1	2	2	3	3	4	4	Bipartite graph
v^1	0	0	0	0	0	4	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^2	0	0	0	0	1	3	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^3	0	0	0	0	2	2	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^4	0	0	0	0	3	1	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^5	0	0	0	0	4	0	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^6	0	0	0	1	0	2	1	1	$H_{5/5}^3, H_{5/5}^2$
v^7	0	0	0	1	1	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^8	0	0	0	1	2	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^9	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^{10}	0	0	1	0	0	2	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{11}	0	0	1	0	1	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{12}	0	0	1	0	2	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{13}	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^{14}	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	0	1	0	1	$H_{3/3}^1$
v^{16}	0	1	0	0	1	0	0	1	$H_{3/3}^1$
v^{17}	0	2	0	1	0	0	0	1	$H_{4/3}^1$
v^{18}	0	2	1	0	0	0	0	1	$H_{4/3}^1$
v^{19}	1	0	0	0	0	1	0	1	$H_{3/3}^1$
v^{20}	1	0	0	0	1	0	0	1	$H_{3/3}^1$
v^{21}	1	1	0	1	0	0	0	1	$H_{4/3}^1$
v^{22}	1	1	1	0	0	0	0	1	$H_{4/3}^1$
v^{23}	2	0	0	1	0	0	0	1	$H_{4/3}^1$
v^{24}	2	0	1	0	0	0	0	1	$H_{4/3}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep	$P_1 P_1$	$P_2 P_3$	N_1
dim	4	4	2
$D_1 D_2$	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$P_1 P_1$	$P_2 P_3$	N_1	
	4	4	2	Bipartite graph
v^1	0	1	2	$H_{3/2}^1$
v^2	1	0	2	$H_{3/2}^1$

SG 207

Space group $P432$, No. 207, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5
	dim	1	1	2	3	3	1	1	1	1	2
Δ_1	1	1	0	1	1	0	-1	-1	0	0	0
Δ_2	1	0	0	0	1	1	0	0	0	0	-1
Δ_3	1	0	1	1	0	1	0	0	-1	-1	0
Δ_4	1	0	0	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5	
	1	1	2	3	3	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	1	1	$H_{3/3}^1$
v^2	0	0	0	0	1	0	0	1	0	1	$H_{3/3}^1$
v^3	0	0	0	1	0	0	1	0	0	1	$H_{3/3}^1$
v^4	0	0	0	1	0	1	0	0	0	1	$H_{3/3}^1$
v^5	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^6	0	0	1	0	0	0	1	1	0	0	$H_{3/2}^1$
v^7	0	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
dim		1	1	2	3	3	1	1	1	1	2
Σ_1	1	1	0	1	1	2	-1	0	0	-1	-1
Σ_2	1	0	1	1	2	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	1	1	$H_{3/3}^1$
v^2	0	0	0	0	1	0	0	1	2	0	$H_{4/3}^1$
v^3	0	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/3}^1$
v^5	0	0	0	0	1	1	0	1	1	0	$H_{4/3}^1$
v^6	0	0	0	0	1	1	1	0	1	0	$H_{4/3}^1$
v^7	0	0	0	0	1	2	0	1	0	0	$H_{4/3}^1$
v^8	0	0	0	0	1	2	1	0	0	0	$H_{4/3}^1$
v^9	0	0	0	1	0	0	0	1	0	1	$H_{3/3}^1$
v^{10}	0	0	0	1	0	0	0	2	1	0	$H_{4/3}^1$
v^{11}	0	0	0	1	0	0	1	0	0	1	$H_{3/3}^1$
v^{12}	0	0	0	1	0	0	1	1	1	0	$H_{4/3}^1$
v^{13}	0	0	0	1	0	0	2	0	1	0	$H_{4/3}^1$
v^{14}	0	0	0	1	0	1	0	2	0	0	$H_{4/3}^1$
v^{15}	0	0	0	1	0	1	1	1	0	0	$H_{4/3}^1$
v^{16}	0	0	0	1	0	1	2	0	0	0	$H_{4/3}^1$
v^{17}	0	0	0	1	1	0	0	0	0	3	$H_{5/6}^2$
v^{18}	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{19}	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{21}	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{24}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	0	1	0	0	1	0	0	0	0	2	$H_{4/4}^4$
v^{26}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{28}	1	0	0	1	0	0	0	0	0	2	$H_{4/4}^4$
v^{29}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1	R_2	R_3	R_4	R_5
dim	1	1	2	3	3	1	1	2	3	3
Λ_1	1	1	0	1	1	-1	-1	0	-1	-1
Λ_2	1	0	0	1	1	0	0	-1	-1	-1
Λ_3	1	0	0	1	1	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1	R_2	R_3	R_4	R_5	Bipartite graph
	1	1	2	3	3	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/3}^1$
v^2	0	0	0	0	1	0	0	0	1	0	$H_{2/3}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/3}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/3}^1$
v^5	0	0	0	1	0	0	0	0	0	1	$H_{2/3}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/3}^1$
v^7	0	0	0	1	0	0	1	1	0	0	$H_{3/3}^1$
v^8	0	0	0	1	0	1	0	1	0	0	$H_{3/3}^1$
v^9	0	0	1	0	0	0	0	1	0	0	$H_{2/2}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	1	1	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{13}	0	1	1	0	0	0	0	0	1	0	$H_{3/3}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{17}	1	0	1	0	0	0	0	0	1	0	$H_{3/3}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	X_3	X_4	X_5	R_1	R_2	R_3	R_4	R_5
dim	1	1	1	1	2	1	1	2	3	3
S_1	1	1	0	0	1	1	-1	0	-1	-2
S_2	1	0	1	1	0	1	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	X_1	X_2	X_3	X_4	X_5	R_1	R_2	R_3	R_4	R_5	
	1	1	1	1	2	1	1	2	3	3	Bipartite graph
v^1	0	0	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^2	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	0	1	$H_{4/4}^4$
v^4	0	0	0	0	2	1	0	0	1	0	$H_{4/4}^4$
v^5	0	0	0	0	3	0	0	0	1	1	$H_{5/6}^2$
v^6	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	1	1	0	0	0	0	1	$H_{3/3}^1$
v^8	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{10}	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	1	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{12}	0	0	2	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{13}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{15}	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	1	0	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{17}	0	1	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{18}	0	2	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{19}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{21}	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	1	0	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{23}	1	0	2	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{24}	1	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{25}	1	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{26}	1	1	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{27}	1	2	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{28}	2	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{29}	2	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1	X_2	X_3	X_4	X_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Z_1	1	1	0	1	0	1	-1	0	-1	0	-1
Z_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	X_1	X_2	X_3	X_4	X_5	M_1	M_2	M_3	M_4	M_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	R_1	R_2	R_3	R_4	R_5
	dim	1	1	1	1	2	1	1	2	3	3
T_1	1	1	1	0	0	0	-1	0	-1	-1	0
T_2	1	0	0	0	0	1	0	0	0	-1	-1
T_3	1	0	0	1	1	0	0	-1	-1	0	-1
T_4	1	0	0	0	0	1	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	R_1	R_2	R_3	R_4	R_5	
	1	1	1	1	2	1	1	2	3	3	Bipartite graph
v^1	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	0	1	$H_{3/3}^1$
v^3	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	1	0	0	0	0	1	$H_{3/3}^1$
v^5	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$

v^6	0	1	0	0	1	0	0	0	1	0	$H_{3/3}^1$
v^7	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{11}	1	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$

SG 208

Space group $P4_232$, No. 208, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5
	dim	1	1	2	3	3	1	1	1	1	2
Δ_1	1	1	0	1	1	0	0	0	0	0	-1
Δ_2	1	0	0	0	1	1	-1	-1	0	0	0
Δ_3	1	0	1	1	0	1	0	0	0	0	-1
Δ_4	1	0	0	0	1	1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 19.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	1	1	0	2	0	2	1	$H_{7/6}^2$
v^2	0	0	0	1	1	0	2	1	1	1	$H_{7/6}^2$
v^3	0	0	0	1	1	0	2	2	0	1	$H_{7/6}^2$
v^4	0	0	0	1	1	1	1	0	2	1	$H_{7/6}^2$
v^5	0	0	0	1	1	1	1	1	1	1	$H_{7/6}^2$
v^6	0	0	0	1	1	1	1	2	0	1	$H_{7/6}^2$
v^7	0	0	0	1	1	2	0	0	2	1	$H_{7/6}^2$
v^8	0	0	0	1	1	2	0	1	1	1	$H_{7/6}^2$
v^9	0	0	0	1	1	2	0	2	0	1	$H_{7/6}^2$
v^{10}	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{11}	0	1	0	1	0	0	1	0	1	1	$H_{5/4}^2$
v^{12}	0	1	0	1	0	0	1	1	0	1	$H_{5/4}^2$
v^{13}	0	1	0	1	0	1	0	0	1	1	$H_{5/4}^2$
v^{14}	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^2$
v^{15}	1	0	0	0	1	0	1	0	1	1	$H_{5/4}^2$

v^{16}	1	0	0	0	1	0	1	1	0	1	$H_{5/4}^2$
v^{17}	1	0	0	0	1	1	0	0	1	1	$H_{5/4}^2$
v^{18}	1	0	0	0	1	1	0	1	0	1	$H_{5/4}^2$
v^{19}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	2	3	3	1	1	1	1	2
Σ_1	1	1	0	1	1	2	0	-1	-1	0	-1
Σ_2	1	0	1	1	2	1	-1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	1	0	1	$H_{3/3}^1$
v^2	0	0	0	0	1	0	0	2	1	0	$H_{4/3}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/3}^1$
v^4	0	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^5	0	0	0	0	1	0	2	0	1	0	$H_{4/3}^1$
v^6	0	0	0	0	1	1	0	2	0	0	$H_{4/3}^1$
v^7	0	0	0	0	1	1	1	1	0	0	$H_{4/3}^1$
v^8	0	0	0	0	1	1	2	0	0	0	$H_{4/3}^1$
v^9	0	0	0	1	0	0	0	0	1	1	$H_{3/3}^1$
v^{10}	0	0	0	1	0	0	0	1	2	0	$H_{4/3}^1$
v^{11}	0	0	0	1	0	0	1	0	2	0	$H_{4/3}^1$
v^{12}	0	0	0	1	0	1	0	0	0	1	$H_{3/3}^1$
v^{13}	0	0	0	1	0	1	0	1	1	0	$H_{4/3}^1$
v^{14}	0	0	0	1	0	1	1	0	1	0	$H_{4/3}^1$
v^{15}	0	0	0	1	0	2	0	1	0	0	$H_{4/3}^1$
v^{16}	0	0	0	1	0	2	1	0	0	0	$H_{4/3}^1$
v^{17}	0	0	0	1	1	0	0	0	0	3	$H_{5/6}^2$
v^{18}	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{19}	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{21}	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{24}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$

v^{25}	0	1	0	0	1	0	0	0	0	2	$H_{4/4}^4$
v^{26}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	1	0	0	1	0	0	0	0	0	2	$H_{4/4}^4$
v^{29}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1	R_2	R_3	R_4	R_5
	dim	1	1	2	3	3	1	1	2	3	3
Λ_1	1	1	1	0	1	1	-1	-1	0	-1	-1
Λ_2	1	0	0	1	1	1	0	0	-1	-1	-1
Λ_3	1	0	0	1	1	1	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1	R_2	R_3	R_4	R_5	Bipartite graph
	1	1	2	3	3	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/3}^1$
v^2	0	0	0	0	1	0	0	0	1	0	$H_{2/3}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/3}^1$
v^4	0	0	0	0	1	1	0	1	0	0	$H_{3/3}^1$
v^5	0	0	0	1	0	0	0	0	0	1	$H_{2/3}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/3}^1$
v^7	0	0	0	1	0	0	1	1	0	0	$H_{3/3}^1$
v^8	0	0	0	1	0	1	0	1	0	0	$H_{3/3}^1$
v^9	0	0	1	0	0	0	0	1	0	0	$H_{2/2}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	1	1	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{13}	0	1	1	0	0	0	0	0	1	0	$H_{3/3}^1$
v^{14}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	1	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{17}	1	0	1	0	0	0	0	0	1	0	$H_{3/3}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	X_5	R_1	R_2	R_3	R_4	R_5
	dim	1	1	1	1	2	1	1	2	3	3
S_1	1	1	0	0	1	1	0	-1	-1	-2	-1
S_2	1	0	1	1	0	1	-1	0	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	X_1	X_2	X_3	X_4	X_5	R_1	R_2	R_3	R_4	R_5	Bipartite graph
	1	1	1	1	2	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^2	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	0	1	$H_{4/4}^4$
v^4	0	0	0	0	2	1	0	0	1	0	$H_{4/4}^4$
v^5	0	0	0	0	3	0	0	0	1	1	$H_{5/6}^2$
v^6	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	0	1	1	0	0	0	1	0	$H_{3/3}^1$
v^8	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{10}	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	1	2	0	0	0	0	1	0	$H_{4/3}^1$
v^{12}	0	0	2	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{13}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{15}	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	1	0	2	0	0	0	0	1	0	$H_{4/3}^1$
v^{17}	0	1	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{18}	0	2	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{19}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{21}	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	1	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{23}	1	0	2	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{24}	1	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{25}	1	1	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{26}	1	1	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{27}	1	2	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{28}	2	0	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{29}	2	1	0	0	0	0	0	0	1	0	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_1	X_2	X_3	X_4	X_5	M_1	M_2	M_3	M_4	M_5	
dim	1	1	1	1	2	1	1	1	1	2	
Z_1	1	0	1	0	1	1	-1	0	-1	0	-1
Z_2	1	1	0	1	0	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	X_1	X_2	X_3	X_4	X_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	1	1	2	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_1	M_2	M_3	M_4	M_5	R_1	R_2	R_3	R_4	R_5
dim	1	1	1	1	2	1	1	2	3	3
T_1	1	0	0	1	1	0	0	0	-1	-1
T_2	1	1	1	0	0	0	0	0	-1	-1
T_3	1	0	0	0	0	1	-1	0	-1	0
T_4	1	0	0	0	0	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 19.

	M_1	M_2	M_3	M_4	M_5	R_1	R_2	R_3	R_4	R_5	Bipartite graph
	1	1	1	1	2	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^2	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^3	0	1	0	1	1	0	1	0	1	0	$H_{5/4}^2$
v^4	0	1	0	1	1	1	0	0	0	1	$H_{5/4}^2$
v^5	0	1	1	0	1	0	1	0	1	0	$H_{5/4}^2$
v^6	0	1	1	0	1	1	0	0	0	1	$H_{5/4}^2$
v^7	0	2	0	2	1	0	0	0	1	1	$H_{7/6}^2$
v^8	0	2	1	1	1	0	0	0	1	1	$H_{7/6}^2$
v^9	0	2	2	0	1	0	0	0	1	1	$H_{7/6}^2$
v^{10}	1	0	0	1	1	0	1	0	1	0	$H_{5/4}^2$
v^{11}	1	0	0	1	1	1	0	0	0	1	$H_{5/4}^2$
v^{12}	1	0	1	0	1	0	1	0	1	0	$H_{5/4}^2$
v^{13}	1	0	1	0	1	1	0	0	0	1	$H_{5/4}^2$
v^{14}	1	1	0	2	1	0	0	0	1	1	$H_{7/6}^2$
v^{15}	1	1	1	1	1	0	0	0	1	1	$H_{7/6}^2$
v^{16}	1	1	2	0	1	0	0	0	1	1	$H_{7/6}^2$
v^{17}	2	0	0	2	1	0	0	0	1	1	$H_{7/6}^2$
v^{18}	2	0	1	1	1	0	0	0	1	1	$H_{7/6}^2$
v^{19}	2	0	2	0	1	0	0	0	1	1	$H_{7/6}^2$

SG 209

Space group $F432$, No. 209, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5
	dim	1	1	2	3	3	1	1	1	1	2
Δ_1	1	1	0	1	1	0	-1	-1	0	0	0
Δ_2	1	0	0	0	1	1	0	0	0	0	-1
Δ_3	1	0	1	1	0	1	0	0	-1	-1	0
Δ_4	1	0	0	0	1	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5	Bipartite graph
1	1	2	3	3	1	1	1	1	2	

v^1	0 0 0 0 1	0 0 0 1 1	$H_{3/3}^1$
v^2	0 0 0 0 1	0 0 1 0 1	$H_{3/3}^1$
v^3	0 0 0 1 0	0 1 0 0 1	$H_{3/3}^1$
v^4	0 0 0 1 0	1 0 0 0 1	$H_{3/3}^1$
v^5	0 0 1 0 0	0 1 0 1 0	$H_{3/2}^1$
v^6	0 0 1 0 0	0 1 1 0 0	$H_{3/2}^1$
v^7	0 0 1 0 0	1 0 0 1 0	$H_{3/2}^1$
v^8	0 0 1 0 0	1 0 1 0 0	$H_{3/2}^1$
v^9	0 1 0 0 0	0 0 0 1 0	$H_{2/1}^1$
v^{10}	0 1 0 0 0	0 0 1 0 0	$H_{2/1}^1$
v^{11}	1 0 0 0 0	0 1 0 0 0	$H_{2/1}^1$
v^{12}	1 0 0 0 0	1 0 0 0 0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1	L_2	L_3
	dim	1	1	2	3	3	1	1	2
Λ_1	1	1	1	0	1	1	-1	-1	0
Λ_2	1	0	0	1	1	1	0	0	-1
Λ_3	1	0	0	1	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1	L_2	L_3	Bipartite graph
	1	1	2	3	3	1	1	2	
v^1	0	0	0	0	1	0	1	1	$H_{3/3}^1$
v^2	0	0	0	0	1	1	0	1	$H_{3/3}^1$
v^3	0	0	0	1	0	0	1	1	$H_{3/3}^1$
v^4	0	0	0	1	0	1	0	1	$H_{3/3}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		X_1	X_2	X_3	X_4	X_5	W_1	W_2W_4	W_3
	dim	1	1	1	1	2	1	2	1
Z_1	1	1	0	1	0	1	-1	0	-1
Z_2	1	0	1	0	1	1	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	X_1	X_2	X_3	X_4	X_5	W_1	W_2W_4	W_3	Bipartite graph
	1	1	1	1	2	1	2	1	
v^1	0	0	0	0	2	0	1	2	$H_{5/4}^3$
v^2	0	0	0	0	2	1	1	1	$H_{5/4}^3$
v^3	0	0	0	0	2	2	1	0	$H_{5/4}^3$
v^4	0	0	0	1	1	0	1	1	$H_{4/3}^2$
v^5	0	0	0	1	1	1	1	0	$H_{4/3}^2$
v^6	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^9	0	1	0	0	1	0	1	1	$H_{4/3}^2$
v^{10}	0	1	0	0	1	1	1	0	$H_{4/3}^2$
v^{11}	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^{12}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	1	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q : $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		L_1	L_2	L_3	W_1	W_2W_4	W_3
	dim	1	1	2	1	2	1
Q_1	1	1	0	1	-1	-1	0
Q_2	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1	L_2	L_3	W_1	W_2W_4	W_3	Bipartite graph
	1	1	2	1	2	1	
v^1	0	0	1	0	1	0	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	1	0	0	1	0	0	$H_{2/1}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$

SG 210

Space group $F4_132$, No. 210, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5
dim		1	1	2	3	3	1	1	1	1	2
Δ_1	1	1	0	1	1	0	0	0	0	0	-1
Δ_2	1	0	0	0	1	1	0	0	-1	-1	0
Δ_3	1	0	1	1	0	1	0	0	0	0	-1
Δ_4	1	0	0	0	1	1	-1	-1	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 19.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	1	1	0	2	0	2	1	$H_{7/6}^2$
v^2	0	0	0	1	1	0	2	1	1	1	$H_{7/6}^2$
v^3	0	0	0	1	1	0	2	2	0	1	$H_{7/6}^2$
v^4	0	0	0	1	1	1	1	0	2	1	$H_{7/6}^2$
v^5	0	0	0	1	1	1	1	1	1	1	$H_{7/6}^2$
v^6	0	0	0	1	1	1	1	2	0	1	$H_{7/6}^2$
v^7	0	0	0	1	1	2	0	0	2	1	$H_{7/6}^2$
v^8	0	0	0	1	1	2	0	1	1	1	$H_{7/6}^2$
v^9	0	0	0	1	1	2	0	2	0	1	$H_{7/6}^2$
v^{10}	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{11}	0	1	0	1	0	0	1	0	1	1	$H_{5/4}^2$
v^{12}	0	1	0	1	0	0	1	1	0	1	$H_{5/4}^2$
v^{13}	0	1	0	1	0	1	0	0	1	1	$H_{5/4}^2$
v^{14}	0	1	0	1	0	1	0	1	0	1	$H_{5/4}^2$
v^{15}	1	0	0	0	1	0	1	0	1	1	$H_{5/4}^2$
v^{16}	1	0	0	0	1	0	1	1	0	1	$H_{5/4}^2$
v^{17}	1	0	0	0	1	1	0	0	1	1	$H_{5/4}^2$
v^{18}	1	0	0	0	1	1	0	1	0	1	$H_{5/4}^2$
v^{19}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1	L_2	L_3
	dim	1	1	2	3	3	1	1	2
Λ_1	1	1	1	0	1	1	-1	-1	0
Λ_2	1	0	0	1	1	1	0	0	-1
Λ_3	1	0	0	1	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1	L_2	L_3	
	1	1	2	3	3	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	1	1	$H_{3/3}^1$
v^2	0	0	0	0	1	1	0	1	$H_{3/3}^1$
v^3	0	0	0	1	0	0	1	1	$H_{3/3}^1$
v^4	0	0	0	1	0	1	0	1	$H_{3/3}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^6	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	0	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z: $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		X_1	X_2	X_3	X_4	X_5	W_1
	dim	1	1	1	1	2	2
Z_1	1	0	1	0	1	1	-1
Z_2	1	1	0	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	X_1	X_2	X_3	X_4	X_5	W_1	
	1	1	1	1	2	2	Bipartite graph
v^1	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		L_1	L_2	L_3	W_1
	dim	1	1	2	2
Q_1	1	1	0	1	-1
Q_2	1	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1	L_2	L_3	W_1	
	1	1	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	1	1	0	1	$H_{3/2}^1$

SG 211

Space group $I432$, No. 211, without SOC

 Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1	N_2	N_3	N_4
	dim	1	1	2	3	3	1	1	1	1
Σ_1	1	1	0	1	1	2	-1	0	0	-1
Σ_2	1	0	1	1	2	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1	N_2	N_3	N_4	
	1	1	2	3	3	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	0	1	2	$H_{4/3}^1$
v^2	0	0	0	0	1	0	1	0	2	$H_{4/3}^1$
v^3	0	0	0	0	1	1	0	1	1	$H_{4/3}^1$
v^4	0	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v^5	0	0	0	0	1	2	0	1	0	$H_{4/3}^1$
v^6	0	0	0	0	1	2	1	0	0	$H_{4/3}^1$
v^7	0	0	0	1	0	0	0	2	1	$H_{4/3}^1$
v^8	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v^9	0	0	0	1	0	0	2	0	1	$H_{4/3}^1$
v^{10}	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$

v^{11}	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^{12}	0	0	0	1	0	1	2	0	0	$H_{4/3}^1$
v^{13}	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^{15}	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{16}	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0)$, $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1	H_2	H_3	H_4	H_5
	dim	1	1	2	3	3	1	1	2	3	3
Δ_1	1	1	0	1	1	0	-1	0	-1	-1	0
Δ_2	1	0	0	0	1	1	0	0	0	-1	-1
Δ_3	1	0	1	1	0	1	0	-1	-1	0	-1
Δ_4	1	0	0	0	1	1	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1	H_2	H_3	H_4	H_5	Bipartite graph
	1	1	2	3	3	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/3}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/3}^1$
v^3	0	0	1	0	0	0	0	1	0	0	$H_{2/2}^1$
v^4	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^5	0	0	1	0	1	0	2	0	1	0	$H_{5/5}^1$
v^6	0	0	1	1	0	2	0	0	0	1	$H_{5/5}^1$
v^7	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	1	0	1	0	1	0	0	0	1	$H_{4/4}^3$
v^9	0	2	0	1	0	0	0	1	0	1	$H_{5/5}^1$
v^{10}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	1	0	1	0	$H_{4/4}^3$
v^{12}	1	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{13}	2	0	0	0	1	0	0	1	1	0	$H_{5/5}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3	P_4
	dim	1	1	2	3	3	1	1	1	3
Λ_1	1	1	1	0	1	1	-1	0	0	-1
Λ_2	1	0	0	1	1	1	0	0	-1	-1
Λ_3	1	0	0	1	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3	P_4	
	1	1	2	3	3	1	1	1	3	Bipartite graph
v^1	0	0	0	0	1	0	0	0	1	$H_{2/3}^1$
v^2	0	0	0	0	1	1	1	1	0	$H_{4/3}^1$
v^3	0	0	0	1	0	0	0	0	1	$H_{2/3}^1$
v^4	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^5	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	1	0	0	0	0	0	1	$H_{3/3}^1$
v^8	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	1	0	0	0	0	0	1	$H_{3/3}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$
A matrix

corep		H_1	H_2	H_3	H_4	H_5	N_1	N_2	N_3	N_4
	dim	1	1	2	3	3	1	1	1	1
G_1	1	1	0	1	1	2	-1	-1	0	0
G_2	1	0	1	1	2	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	H_1	H_2	H_3	H_4	H_5	N_1	N_2	N_3	N_4	
	1	1	2	3	3	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	2	0	1	$H_{4/3}^1$
v^2	0	0	0	0	1	0	2	1	0	$H_{4/3}^1$
v^3	0	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v^4	0	0	0	0	1	1	1	1	0	$H_{4/3}^1$
v^5	0	0	0	0	1	2	0	0	1	$H_{4/3}^1$

v^6	0	0	0	0	1	2	0	1	0	$H_{4/3}^1$
v^7	0	0	0	1	0	0	1	0	2	$H_{4/3}^1$
v^8	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v^9	0	0	0	1	0	0	1	2	0	$H_{4/3}^1$
v^{10}	0	0	0	1	0	1	0	0	2	$H_{4/3}^1$
v^{11}	0	0	0	1	0	1	0	1	1	$H_{4/3}^1$
v^{12}	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$
v^{13}	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^{15}	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{16}	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1	H_2	H_3	H_4	H_5	P_1	P_2	P_3	P_4
	dim	1	1	2	3	3	1	1	1	3
F_1	1	1	1	0	1	1	-1	0	0	-1
F_2	1	0	0	1	1	1	0	0	-1	-1
F_3	1	0	0	1	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	H_1	H_2	H_3	H_4	H_5	P_1	P_2	P_3	P_4	Bipartite graph
	1	1	2	3	3	1	1	1	3	
v^1	0	0	0	0	1	0	0	0	1	$H_{2/3}^1$
v^2	0	0	0	0	1	1	1	1	0	$H_{4/3}^1$
v^3	0	0	0	1	0	0	0	0	1	$H_{2/3}^1$
v^4	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^5	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	1	0	0	0	0	0	1	$H_{3/3}^1$
v^8	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	1	0	1	0	0	0	0	0	1	$H_{3/3}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_1	P_2	P_3	P_4	N_1	N_2	N_3	N_4
	dim	1	1	1	3	1	1	1	1
D_1	1	1	1	1	1	-1	0	-1	0
D_2	1	0	0	0	2	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	P_1	P_2	P_3	P_4	N_1	N_2	N_3	N_4	
	1	1	1	3	1	1	1	1	Bipartite graph
v^1	0	0	0	1	0	0	1	2	$H_{4/3}^1$
v^2	0	0	0	1	0	1	1	1	$H_{4/3}^1$
v^3	0	0	0	1	0	2	1	0	$H_{4/3}^1$
v^4	0	0	0	1	1	0	0	2	$H_{4/3}^1$
v^5	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v^6	0	0	0	1	1	2	0	0	$H_{4/3}^1$
v^7	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 212

Space group $P4_332$, No. 212, without SOC

 Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2
	dim	1	1	2	3	3	2	2
Δ_1	1	1	0	1	1	0	-1	0
Δ_2	1	0	0	0	1	1	0	-1
Δ_3	1	0	1	1	0	1	0	-1
Δ_4	1	0	0	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	
	1	1	2	3	3	2	2	Bipartite graph
v^1	0	0	1	1	1	2	2	$H_{7/8}^{10}, H_{7/8}^3$
v^2	0	1	0	1	0	1	1	$H_{4/4}^4$
v^3	1	0	0	0	1	1	1	$H_{4/4}^4$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
dim	1	1	2	3	3	2	2	2	
Σ_1	1	1	0	1	1	2	0	-2	-1
Σ_2	1	0	1	1	2	1	-2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 21.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
	1	1	2	3	3	2	2	2	Bipartite graph
v^1	0	0	0	0	2	0	1	2	$H_{5/6}^2$
v^2	0	0	0	0	2	1	2	0	$H_{5/6}^2$
v^3	0	0	0	1	1	0	0	3	$H_{5/6}^2$
v^4	0	0	0	1	1	1	1	1	$H_{5/6}^2, H_{5/6}^4$
v^5	0	0	0	2	0	1	0	2	$H_{5/6}^2$
v^6	0	0	0	2	0	2	1	0	$H_{5/6}^2$
v^7	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	1	1	1	2	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^9	0	0	2	0	0	1	1	0	$H_{4/4}^2$
v^{10}	0	1	0	0	1	0	0	2	$H_{4/4}^4$
v^{11}	0	1	0	0	1	1	1	0	$H_{4/4}^4$
v^{12}	0	1	0	1	0	1	0	1	$H_{4/4}^4$
v^{13}	0	1	1	1	0	2	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{14}	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^{15}	1	0	0	0	1	0	1	1	$H_{4/4}^4$
v^{16}	1	0	0	1	0	0	0	2	$H_{4/4}^4$
v^{17}	1	0	0	1	0	1	1	0	$H_{4/4}^4$
v^{18}	1	0	1	0	1	1	2	0	$H_{6/6}^8, H_{6/6}^7$
v^{19}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	1	1	1	0	0	1	1	0	$H_{5/4}^3$
v^{21}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	$R_1 R_2$	R_3
dim		1	1	2	3	3	4	4
Λ_1	1	1	1	0	1	1	0	-2
Λ_2	1	0	0	1	1	1	-2	-1
Λ_3	1	0	0	1	1	1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	$R_1 R_2$	R_3	Bipartite graph
	1	1	2	3	3	4	4	
v^1	0	0	0	0	4	1	2	$H_{7/12}^2, H_{7/12}^1$
v^2	0	0	0	1	3	1	2	$H_{7/12}^2, H_{7/12}^1$
v^3	0	0	0	2	2	1	2	$H_{7/12}^2, H_{7/12}^1$
v^4	0	0	0	3	1	1	2	$H_{7/12}^2, H_{7/12}^1$
v^5	0	0	0	4	0	1	2	$H_{7/12}^2, H_{7/12}^1$
v^6	0	0	1	0	2	1	1	$H_{5/8}^2, H_{5/8}^3$
v^7	0	0	1	1	1	1	1	$H_{5/8}^2, H_{5/8}^3$
v^8	0	0	1	2	0	1	1	$H_{5/8}^2, H_{5/8}^3$
v^9	0	0	2	0	0	1	0	$H_{3/4}^1$
v^{10}	0	1	0	0	1	0	1	$H_{3/4}^2$
v^{11}	0	1	0	1	0	0	1	$H_{3/4}^2$
v^{12}	0	2	1	0	0	0	1	$H_{4/4}^1$
v^{13}	1	0	0	0	1	0	1	$H_{3/4}^2$
v^{14}	1	0	0	1	0	0	1	$H_{3/4}^2$
v^{15}	1	1	1	0	0	0	1	$H_{4/4}^1$
v^{16}	2	0	1	0	0	0	1	$H_{4/4}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	$R_1 R_2$	R_3
dim		2	2	4	4
$S_1 S_2$	2	1	1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	X_1	X_2	$R_1 R_2$	R_3	
	2	2	4	4	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	0	2	1	0	$H_{3/2}^1$
v^3	1	1	0	1	$H_{3/2}^1$
v^4	1	1	1	0	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$
v^6	2	0	1	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	$M_1 M_2$	$M_3 M_4$	M_5
dim	2	2	2	2	2
$Z_1 Z_2$	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	X_1	X_2	$M_1 M_2$	$M_3 M_4$	M_5	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0)$, $R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$M_1 M_2$	$M_3 M_4$	M_5	$R_1 R_2$	R_3
dim	2	2	2	4	4
$T_1 T_3$	2	0	0	1	-1
$T_2 T_4$	2	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1M_2	M_3M_4	M_5	R_1R_2	R_3	
	2	2	2	4	4	Bipartite graph
v^1	0	1	1	0	1	$H_{3/2}^1$
v^2	0	1	1	1	0	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	1	0	$H_{3/2}^1$

SG 213

Space group $P4_132$, No. 213, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2
	dim	1	1	2	3	3	2	2
Δ_1	1	1	0	1	1	0	0	-1
Δ_2	1	0	0	0	1	1	0	-1
Δ_3	1	0	1	1	0	1	-1	0
Δ_4	1	0	0	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	
	1	1	2	3	3	2	2	Bipartite graph
v^1	0	0	1	1	1	2	2	$H_{7/8}^{10}, H_{7/8}^3$
v^2	0	1	0	1	0	1	1	$H_{4/4}^4$
v^3	1	0	0	0	1	1	1	$H_{4/4}^4$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5
	dim	1	1	2	3	3	2	2	2
Σ_1	1	1	0	1	1	2	-2	0	-1
Σ_2	1	0	1	1	2	1	0	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 21.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1M_2	M_3M_4	M_5	
	1	1	2	3	3	2	2	2	Bipartite graph
v^1	0	0	0	0	2	1	0	2	$H_{5/6}^2$
v^2	0	0	0	0	2	2	1	0	$H_{5/6}^2$
v^3	0	0	0	1	1	0	0	3	$H_{5/6}^2$
v^4	0	0	0	1	1	1	1	1	$H_{5/6}^2, H_{5/6}^4$
v^5	0	0	0	2	0	0	1	2	$H_{5/6}^2$
v^6	0	0	0	2	0	1	2	0	$H_{5/6}^2$
v^7	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	1	1	1	2	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^9	0	0	2	0	0	1	1	0	$H_{4/4}^2$
v^{10}	0	1	0	0	1	0	0	2	$H_{4/4}^4$
v^{11}	0	1	0	0	1	1	1	0	$H_{4/4}^4$
v^{12}	0	1	0	1	0	0	1	1	$H_{4/4}^4$
v^{13}	0	1	1	1	0	1	2	0	$H_{6/6}^8, H_{6/6}^7$
v^{14}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	1	1	0	1	$H_{4/4}^4$
v^{16}	1	0	0	1	0	0	0	2	$H_{4/4}^4$
v^{17}	1	0	0	1	0	1	1	0	$H_{4/4}^4$
v^{18}	1	0	1	0	1	2	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{19}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	1	1	1	0	0	1	1	0	$H_{5/4}^3$
v^{21}	2	0	0	0	0	1	0	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1R_2	R_3
	dim	1	1	2	3	3	4	4
Λ_1	1	1	1	0	1	1	0	-2
Λ_2	1	0	0	1	1	1	-2	-1
Λ_3	1	0	0	1	1	1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1R_2	R_3	
	1	1	2	3	3	4	4	Bipartite graph
v^1	0	0	0	0	4	1	2	$H_{7/12}^2, H_{7/12}^1$
v^2	0	0	0	1	3	1	2	$H_{7/12}^2, H_{7/12}^1$
v^3	0	0	0	2	2	1	2	$H_{7/12}^2, H_{7/12}^1$

v^4	0	0	0	3	1	1	2	$H_{7/12}^2, H_{7/12}^1$
v^5	0	0	0	4	0	1	2	$H_{7/12}^2, H_{7/12}^1$
v^6	0	0	1	0	2	1	1	$H_{5/8}^2, H_{5/8}^3$
v^7	0	0	1	1	1	1	1	$H_{5/8}^2, H_{5/8}^3$
v^8	0	0	1	2	0	1	1	$H_{5/8}^2, H_{5/8}^3$
v^9	0	0	2	0	0	1	0	$H_{3/4}^1$
v^{10}	0	1	0	0	1	0	1	$H_{3/4}^2$
v^{11}	0	1	0	1	0	0	1	$H_{3/4}^2$
v^{12}	0	2	1	0	0	0	1	$H_{4/4}^1$
v^{13}	1	0	0	0	1	0	1	$H_{3/4}^2$
v^{14}	1	0	0	1	0	0	1	$H_{3/4}^2$
v^{15}	1	1	1	0	0	0	1	$H_{4/4}^1$
v^{16}	2	0	1	0	0	0	1	$H_{4/4}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	R_1R_2	R_3
dim	2	2	4	4
S_1S_2	2	1	1	-2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	X_1	X_2	R_1R_2	R_3	
	2	2	4	4	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	0	2	1	0	$H_{3/2}^1$
v^3	1	1	0	1	$H_{3/2}^1$
v^4	1	1	1	0	$H_{3/2}^1$
v^5	2	0	0	1	$H_{3/2}^1$
v^6	2	0	1	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	M_1M_2	M_3M_4	M_5
dim	2	2	2	2	2
Z_1Z_2	2	1	1	-1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	X_1	X_2	M_1M_2	M_3M_4	M_5	Bipartite graph
	2	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1M_2	M_3M_4	M_5	R_1R_2	R_3
	dim	2	2	2	4	4
T_1T_3	2	0	0	1	-1	-1
T_2T_4	2	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_1M_2	M_3M_4	M_5	R_1R_2	R_3	Bipartite graph
	2	2	2	4	4	
v^1	0	1	1	0	1	$H_{3/2}^1$
v^2	0	1	1	1	0	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	0	1	1	0	$H_{3/2}^1$

SG 214

Space group $I4_132$, No. 214, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1	N_2	N_3	N_4
	dim	1	1	2	3	3	1	1	1	1
Σ_1	1	1	0	1	1	2	-1	0	-1	0
Σ_2	1	0	1	1	2	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1	N_2	N_3	N_4	Bipartite graph
	1	1	2	3	3	1	1	1	1	
v^1	0	0	0	0	1	0	0	2	1	$H_{4/3}^1$
v^2	0	0	0	0	1	0	1	2	0	$H_{4/3}^1$
v^3	0	0	0	0	1	1	0	1	1	$H_{4/3}^1$
v^4	0	0	0	0	1	1	1	1	0	$H_{4/3}^1$
v^5	0	0	0	0	1	2	0	0	1	$H_{4/3}^1$
v^6	0	0	0	0	1	2	1	0	0	$H_{4/3}^1$
v^7	0	0	0	1	0	0	0	1	2	$H_{4/3}^1$
v^8	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v^9	0	0	0	1	0	0	2	1	0	$H_{4/3}^1$
v^{10}	0	0	0	1	0	1	0	0	2	$H_{4/3}^1$
v^{11}	0	0	0	1	0	1	1	0	1	$H_{4/3}^1$
v^{12}	0	0	0	1	0	1	2	0	0	$H_{4/3}^1$
v^{13}	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^{15}	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{16}	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1	H_2	H_3	H_4	H_5
	dim	1	1	2	3	3	1	1	2	3	3
Δ_1	1	1	0	1	1	0	0	0	0	-1	-1
Δ_2	1	0	0	0	1	1	0	-1	-1	-1	0
Δ_3	1	0	1	1	0	1	0	0	0	-1	-1
Δ_4	1	0	0	0	1	1	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 25.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1	H_2	H_3	H_4	H_5	Bipartite graph
	1	1	2	3	3	1	1	2	3	3	
v^1	0	0	0	1	1	0	1	1	0	1	$H_{5/6}^5, H_{5/6}^3$
v^2	0	0	0	1	1	1	0	1	1	0	$H_{5/6}^5, H_{5/6}^3$
v^3	0	0	0	1	1	1	2	0	0	1	$H_{6/6}^3$

v^4	0 0 0 1 1	2 1 0 1 0	$H_{6/6}^3$
v^5	0 0 0 2 2	0 0 3 1 1	$H_{9/12}^6, H_{9/12}^2, H_{9/12}^5, H_{9/12}^4, H_{9/12}^8, H_{9/12}^7, H_{9/12}^3, H_{9/12}^1$
v^6	0 0 1 1 1	0 0 1 1 1	$H_{6/8}^4, H_{6/8}^5, H_{6/8}^3$
v^7	0 0 1 1 1	0 2 0 0 2	$H_{7/8}^5, H_{7/8}^6$
v^8	0 0 1 1 1	1 1 0 1 1	$H_{7/8}^4, H_{7/8}^5, H_{7/8}^8, H_{7/8}^6, H_{7/8}^7$
v^9	0 0 1 1 1	2 0 0 2 0	$H_{7/8}^5, H_{7/8}^6$
v^{10}	0 0 2 1 1	0 1 0 1 2	$H_{8/10}^3, H_{8/10}^5, H_{8/10}^1, H_{8/10}^4, H_{8/10}^2, H_{8/10}^7, H_{8/10}^6$
v^{11}	0 0 2 1 1	1 0 0 2 1	$H_{8/10}^3, H_{8/10}^5, H_{8/10}^4, H_{8/10}^1, H_{8/10}^2, H_{8/10}^6, H_{8/10}^7$
v^{12}	0 0 3 1 1	0 0 0 2 2	$H_{9/12}^6, H_{9/12}^2, H_{9/12}^7, H_{9/12}^5, H_{9/12}^3, H_{9/12}^4, H_{9/12}^8, H_{9/12}^1$
v^{13}	0 1 0 1 0	0 1 0 0 1	$H_{4/4}^3$
v^{14}	0 1 0 1 0	1 0 0 1 0	$H_{4/4}^3$
v^{15}	0 1 0 2 1	0 0 2 1 1	$H_{8/10}^3, H_{8/10}^4, H_{8/10}^6, H_{8/10}^1, H_{8/10}^5, H_{8/10}^7, H_{8/10}^2$
v^{16}	0 1 1 1 0	0 0 0 1 1	$H_{5/6}^5, H_{5/6}^3$
v^{17}	0 2 0 2 0	0 0 1 1 1	$H_{7/8}^5, H_{7/8}^6$
v^{18}	1 0 0 0 1	0 1 0 0 1	$H_{4/4}^3$
v^{19}	1 0 0 0 1	1 0 0 1 0	$H_{4/4}^3$
v^{20}	1 0 0 1 2	0 0 2 1 1	$H_{8/10}^3, H_{8/10}^1, H_{8/10}^5, H_{8/10}^7, H_{8/10}^2, H_{8/10}^4, H_{8/10}^6$
v^{21}	1 0 1 0 1	0 0 0 1 1	$H_{5/6}^5, H_{5/6}^3$
v^{22}	1 1 0 1 1	0 0 1 1 1	$H_{7/8}^4, H_{7/8}^8, H_{7/8}^5, H_{7/8}^7, H_{7/8}^6$
v^{23}	1 2 0 1 0	0 0 0 1 1	$H_{6/6}^3$
v^{24}	2 0 0 0 2	0 0 1 1 1	$H_{7/8}^5, H_{7/8}^6$
v^{25}	2 1 0 0 1	0 0 0 1 1	$H_{6/6}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3
dim	1	1	2	3	3	2	2	2
Λ_1	1	1	0	1	1	0	-1	-1
Λ_2	1	0	0	1	1	-1	0	-1
Λ_3	1	0	0	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3	Bipartite graph
	1	1	2	3	3	2	2	2	
v^1	0	0	0	0	2	1	1	1	$H_{5/6}^2$
v^2	0	0	0	1	1	1	1	1	$H_{5/6}^2$
v^3	0	0	0	2	0	1	1	1	$H_{5/6}^2$
v^4	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^5	0	1	0	0	1	0	1	1	$H_{4/4}^4$

v^6	0	1	0	1	0	0	1	1	$H_{4/4}^4$
v^7	0	2	1	0	0	0	1	1	$H_{5/4}^3$
v^8	1	0	0	0	1	0	1	1	$H_{4/4}^4$
v^9	1	0	0	1	0	0	1	1	$H_{4/4}^4$
v^{10}	1	1	1	0	0	0	1	1	$H_{5/4}^3$
v^{11}	2	0	1	0	0	0	1	1	$H_{5/4}^3$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep		H_1	H_2	H_3	H_4	H_5	N_1	N_2	N_3	N_4
	dim	1	1	2	3	3	1	1	1	1
G_1	1	1	0	1	2	1	-1	-1	0	0
G_2	1	0	1	1	1	2	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	H_1	H_2	H_3	H_4	H_5	N_1	N_2	N_3	N_4	
	1	1	2	3	3	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	1	0	2	$H_{4/3}^1$
v^2	0	0	0	0	1	0	1	1	1	$H_{4/3}^1$
v^3	0	0	0	0	1	0	1	2	0	$H_{4/3}^1$
v^4	0	0	0	0	1	1	0	0	2	$H_{4/3}^1$
v^5	0	0	0	0	1	1	0	1	1	$H_{4/3}^1$
v^6	0	0	0	0	1	1	0	2	0	$H_{4/3}^1$
v^7	0	0	0	1	0	0	2	0	1	$H_{4/3}^1$
v^8	0	0	0	1	0	0	2	1	0	$H_{4/3}^1$
v^9	0	0	0	1	0	1	1	0	1	$H_{4/3}^1$
v^{10}	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^{11}	0	0	0	1	0	2	0	0	1	$H_{4/3}^1$
v^{12}	0	0	0	1	0	2	0	1	0	$H_{4/3}^1$
v^{13}	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^{15}	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{16}	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1	H_2	H_3	H_4	H_5	P_1	P_2	P_3
	dim	1	1	2	3	3	2	2	2
F_1	1	1	1	0	1	1	0	-1	-1
F_2	1	0	0	1	1	1	-1	0	-1
F_3	1	0	0	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	H_1	H_2	H_3	H_4	H_5	P_1	P_2	P_3	Bipartite graph
	1	1	2	3	3	2	2	2	
v^1	0	0	0	0	2	1	1	1	$H_{5/6}^2$
v^2	0	0	0	1	1	1	1	1	$H_{5/6}^2$
v^3	0	0	0	2	0	1	1	1	$H_{5/6}^2$
v^4	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^5	0	1	0	0	1	0	1	1	$H_{4/4}^4$
v^6	0	1	0	1	0	0	1	1	$H_{4/4}^4$
v^7	0	2	1	0	0	0	1	1	$H_{5/4}^3$
v^8	1	0	0	0	1	0	1	1	$H_{4/4}^4$
v^9	1	0	0	1	0	0	1	1	$H_{4/4}^4$
v^{10}	1	1	1	0	0	0	1	1	$H_{5/4}^3$
v^{11}	2	0	1	0	0	0	1	1	$H_{5/4}^3$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_1	P_2	P_3	N_1	N_2	N_3	N_4
	dim	2	2	2	1	1	1	1
D_1	1	1	1	1	-1	0	0	-1
D_2	1	1	1	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	P_1	P_2	P_3	N_1	N_2	N_3	N_4	Bipartite graph
	2	2	2	1	1	1	1	
v^1	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	0	1	1	0	1	0	$H_{3/2}^1$

v^4	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	1	0	0	0	1	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	1	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	$H_{3/2}^1$
v^8	0	1	0	1	1	0	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	1	$H_{3/2}^1$
v^{10}	1	0	0	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{12}	1	0	0	1	1	0	0	$H_{3/2}^1$

SG 215

Space group $P\bar{4}3m$, No. 215, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5
	dim	1	1	2	3	3	1	1	1	1	2
Δ_1	1	1	0	1	0	1	-1	0	0	-1	0
$\Delta_2\Delta_4$	2	0	0	0	1	1	0	0	0	0	-1
Δ_3	1	0	1	1	1	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^7	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
dim		1	1	2	3	3	1	1	1	1	2
Σ_1	1	1	0	1	1	2	-1	0	0	-1	-1
Σ_2	1	0	1	1	2	1	0	-1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	1	1	$H_{3/3}^1$
v^2	0	0	0	0	1	0	0	1	2	0	$H_{4/3}^1$
v^3	0	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$
v^4	0	0	0	0	1	1	0	0	0	1	$H_{3/3}^1$
v^5	0	0	0	0	1	1	0	1	1	0	$H_{4/3}^1$
v^6	0	0	0	0	1	1	1	0	1	0	$H_{4/3}^1$
v^7	0	0	0	0	1	2	0	1	0	0	$H_{4/3}^1$
v^8	0	0	0	0	1	2	1	0	0	0	$H_{4/3}^1$
v^9	0	0	0	1	0	0	0	1	0	1	$H_{3/3}^1$
v^{10}	0	0	0	1	0	0	0	2	1	0	$H_{4/3}^1$
v^{11}	0	0	0	1	0	0	1	0	0	1	$H_{3/3}^1$
v^{12}	0	0	0	1	0	0	1	1	1	0	$H_{4/3}^1$
v^{13}	0	0	0	1	0	0	2	0	1	0	$H_{4/3}^1$
v^{14}	0	0	0	1	0	1	0	2	0	0	$H_{4/3}^1$
v^{15}	0	0	0	1	0	1	1	1	0	0	$H_{4/3}^1$
v^{16}	0	0	0	1	0	1	2	0	0	0	$H_{4/3}^1$
v^{17}	0	0	0	1	1	0	0	0	0	3	$H_{5/6}^2$
v^{18}	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{19}	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{21}	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{24}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	0	1	0	0	1	0	0	0	0	2	$H_{4/4}^4$
v^{26}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{28}	1	0	0	1	0	0	0	0	0	2	$H_{4/4}^4$
v^{29}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1	R_2	R_3	R_4	R_5
	dim	1	1	2	3	3	1	1	2	3	3
Λ_1	1	1	0	0	0	1	-1	0	0	0	-1
Λ_2	1	0	1	0	1	0	0	-1	0	-1	0
Λ_3	2	0	0	1	1	1	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1	R_2	R_3	R_4	R_5	Bipartite graph
	1	1	2	3	3	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	1	1	0	0	1	0	$H_{4/3}^2$
v^8	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	1	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^{11}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	X_5	R_1	R_2	R_3	R_4	R_5
	dim	1	1	1	1	2	1	1	2	3	3
S_1	1	1	0	0	1	1	-1	0	-1	-1	-2
S_2	1	0	1	1	0	1	0	-1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	X_1	X_2	X_3	X_4	X_5	R_1	R_2	R_3	R_4	R_5	Bipartite graph
	1	1	1	1	2	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	1	0	0	$H_{2/2}^1$
v^2	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	2	0	1	0	0	1	$H_{4/4}^4$

v^4	0	0	0	0	2	1	0	0	1	0	$H_{4/4}^4$
v^5	0	0	0	0	3	0	0	0	1	1	$H_{5/6}^2$
v^6	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^7	0	0	0	1	1	0	0	0	0	1	$H_{3/3}^1$
v^8	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{10}	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	0	0	1	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{12}	0	0	2	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{13}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	1	0	0	0	1	0	$H_{3/3}^1$
v^{15}	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{16}	0	1	0	2	0	0	0	0	0	1	$H_{4/3}^1$
v^{17}	0	1	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{18}	0	2	0	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{19}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{21}	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{22}	1	0	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{23}	1	0	2	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{24}	1	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{25}	1	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{26}	1	1	1	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{27}	1	2	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{28}	2	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{29}	2	1	0	0	0	0	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1	X_2	X_3	X_4	X_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	1	1	2	1	1	1	1	2
Z_1	1	1	0	1	0	1	-1	0	-1	0	-1
Z_2	1	0	1	0	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	X_1	X_2	X_3	X_4	X_5	M_1	M_2	M_3	M_4	M_5	
	1	1	1	1	2	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$

v^3	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	M_5	R_1	R_2	R_3	R_4	R_5
	dim	1	1	1	1	2	1	1	2	3	3
T_1	1	1	0	0	1	0	-1	0	-1	0	-1
T_2T_4	2	0	0	0	0	1	0	0	0	-1	-1
T_3	1	0	1	1	0	0	0	-1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	M_1	M_2	M_3	M_4	M_5	R_1	R_2	R_3	R_4	R_5	Bipartite graph
	1	1	1	1	2	1	1	2	3	3	
v^1	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^3	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^4	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^6	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{11}	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	1	0	0	$H_{3/2}^1$

SG 216

Space group $F\bar{4}3m$, No. 216, without SOC Δ High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$ **A** matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5
	dim	1	1	2	3	3	1	1	1	1	2
Δ_1	1	1	0	1	0	1	-1	0	0	-1	0
$\Delta_2\Delta_4$	2	0	0	0	1	1	0	0	0	0	-1
Δ_3	1	0	1	1	1	0	0	-1	-1	0	0

 $\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^7	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

 Λ High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ **A** matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1	L_2	L_3
	dim	1	1	2	3	3	1	1	2
Λ_1	1	1	0	0	0	1	-1	0	0
Λ_2	1	0	1	0	1	0	0	-1	0
Λ_3	2	0	0	1	1	1	0	0	-1

 $\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1	L_2	L_3	
	1	1	2	3	3	1	1	2	Bipartite graph
v^1	0	0	0	0	1	1	0	1	$H_{3/2}^1$
v^2	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	0	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z: $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		X_1	X_2	X_3	X_4	X_5	W_1	W_2	W_3	W_4
	dim	1	1	1	1	2	1	1	1	1
Z_1	1	1	0	1	0	1	-1	0	-1	0
Z_2	1	0	1	0	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_1	X_2	X_3	X_4	X_5	W_1	W_2	W_3	W_4	
	1	1	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix



corep		L_1	L_2	L_3	W_1	W_2	W_3	W_4
	dim	1	1	2	1	1	1	1
Q_1	1	1	1	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	L_1	L_2	L_3	W_1	W_2	W_3	W_4	
	1	1	2	1	1	1	1	Bipartite graph
v^1	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	1	$H_{3/2}^1$
v^5	0	0	1	0	1	1	0	$H_{3/2}^1$
v^6	0	0	1	0	2	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	1	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	1	2	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{17}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	1	0	0	1	0	0	0	$H_{2/1}^1$

SG 217

Space group $I\bar{4}3m$, No. 217, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1	N_2	N_3	N_4
	dim	1	1	2	3	3	1	1	1	1
Σ_1	1	1	0	1	1	2	-1	-1	0	0
Σ_2	1	0	1	1	2	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1	N_2	N_3	N_4	Bipartite graph
	1	1	2	3	3	1	1	1	1	
v^1	0	0	0	0	1	0	2	0	1	$H_{4/3}^1$
v^2	0	0	0	0	1	0	2	1	0	$H_{4/3}^1$
v^3	0	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v^4	0	0	0	0	1	1	1	1	0	$H_{4/3}^1$
v^5	0	0	0	0	1	2	0	0	1	$H_{4/3}^1$
v^6	0	0	0	0	1	2	0	1	0	$H_{4/3}^1$
v^7	0	0	0	1	0	0	1	0	2	$H_{4/3}^1$
v^8	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v^9	0	0	0	1	0	0	1	2	0	$H_{4/3}^1$
v^{10}	0	0	0	1	0	1	0	0	2	$H_{4/3}^1$
v^{11}	0	0	0	1	0	1	0	1	1	$H_{4/3}^1$
v^{12}	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$
v^{13}	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^{15}	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{16}	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1	H_2	H_3	H_4	H_5
	dim	1	1	2	3	3	1	1	2	3	3
Δ_1	1	1	0	1	0	1	-1	0	-1	0	-1
$\Delta_2\Delta_4$	2	0	0	0	1	1	0	0	0	-1	-1
Δ_3	1	0	1	1	1	0	0	-1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1	H_2	H_3	H_4	H_5	Bipartite graph
	1	1	2	3	3	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	1	0	0	0	0	1	0	0	$H_{2/2}^1$
v^4	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$

v^5	0 0 1 0 1	2 0 0 1 0	$H_{5/4}^3$
v^6	0 0 1 1 0	0 2 0 0 1	$H_{5/4}^3$
v^7	0 1 0 0 0	0 1 0 0 0	$H_{2/1}^1$
v^8	0 1 0 0 1	1 0 0 1 0	$H_{4/3}^2$
v^9	0 2 0 0 1	0 0 1 1 0	$H_{5/4}^3$
v^{10}	1 0 0 0 0	1 0 0 0 0	$H_{2/1}^1$
v^{11}	1 0 0 1 0	0 1 0 0 1	$H_{4/3}^2$
v^{12}	1 1 0 0 0	0 0 1 0 0	$H_{3/2}^1$
v^{13}	2 0 0 1 0	0 0 1 0 1	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3	P_4	P_5
	dim	1	1	2	3	3	1	1	2	3	3
Λ_1	1	1	0	0	0	1	-1	0	0	0	-1
Λ_2	1	0	1	0	1	0	0	-1	0	-1	0
Λ_3	2	0	0	1	1	1	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3	P_4	P_5	Bipartite graph
	1	1	2	3	3	1	1	2	3	3	
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	1	1	0	0	1	0	$H_{4/3}^2$
v^8	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	1	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^{11}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep		H_1	H_2	H_3	H_4	H_5	N_1	N_2	N_3	N_4
dim		1	1	2	3	3	1	1	1	1
G_1	1	1	0	1	1	2	-1	0	0	-1
G_2	1	0	1	1	2	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	H_1	H_2	H_3	H_4	H_5	N_1	N_2	N_3	N_4	Bipartite graph
	1	1	2	3	3	1	1	1	1	
v^1	0	0	0	0	1	0	0	1	2	$H_{4/3}^1$
v^2	0	0	0	0	1	0	1	0	2	$H_{4/3}^1$
v^3	0	0	0	0	1	1	0	1	1	$H_{4/3}^1$
v^4	0	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v^5	0	0	0	0	1	2	0	1	0	$H_{4/3}^1$
v^6	0	0	0	0	1	2	1	0	0	$H_{4/3}^1$
v^7	0	0	0	1	0	0	0	2	1	$H_{4/3}^1$
v^8	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v^9	0	0	0	1	0	0	2	0	1	$H_{4/3}^1$
v^{10}	0	0	0	1	0	1	0	2	0	$H_{4/3}^1$
v^{11}	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^{12}	0	0	0	1	0	1	2	0	0	$H_{4/3}^1$
v^{13}	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^{14}	0	0	1	0	0	0	1	0	1	$H_{3/2}^1$
v^{15}	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{16}	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^{17}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1	H_2	H_3	H_4	H_5	P_1	P_2	P_3	P_4	P_5
dim		1	1	2	3	3	1	1	2	3	3
F_1	1	1	0	0	0	1	-1	0	0	0	-1
F_2	1	0	1	0	1	0	0	-1	0	-1	0
F_3	2	0	0	1	1	1	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	H_1	H_2	H_3	H_4	H_5	P_1	P_2	P_3	P_4	P_5	
	1	1	2	3	3	1	1	2	3	3	Bipartite graph
v^1	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	1	0	1	0	0	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^4	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	1	0	0	1	1	0	0	1	0	$H_{4/3}^2$
v^8	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^9	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	1	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^{11}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$

A matrix

corep		P_1	P_2	P_3	P_4	P_5	N_1	N_2	N_3	N_4
	dim	1	1	2	3	3	1	1	1	1
D_1	1	1	0	1	0	1	-1	0	0	0
D_2	1	0	0	0	1	1	0	-1	0	0
D_3	1	0	1	1	1	0	0	0	-1	0
D_4	1	0	0	0	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	P_1	P_2	P_3	P_4	P_5	N_1	N_2	N_3	N_4	
	1	1	2	3	3	1	1	1	1	Bipartite graph
v^1	0	0	0	0	1	1	1	0	1	$H_{4/3}^1$
v^2	0	0	0	1	0	0	1	1	1	$H_{4/3}^1$
v^3	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^4	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$

SG 218

Space group $P\bar{4}3n$, No. 218, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1X_2	X_3X_4	X_5
	dim	1	1	2	3	3	2	2	2
Δ_1	1	1	0	1	0	1	0	0	-1
$\Delta_2\Delta_4$	2	0	0	0	1	1	-1	-1	0
Δ_3	1	0	1	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1X_2	X_3X_4	X_5	Bipartite graph
	1	1	2	3	3	2	2	2	
v^1	0	0	0	1	1	0	2	1	$H_{5/4}^3$
v^2	0	0	0	1	1	1	1	1	$H_{5/4}^3$
v^3	0	0	0	1	1	2	0	1	$H_{5/4}^3$
v^4	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^5	0	1	0	0	1	0	1	1	$H_{4/3}^2$
v^6	0	1	0	0	1	1	0	1	$H_{4/3}^2$
v^7	1	0	0	1	0	0	1	1	$H_{4/3}^2$
v^8	1	0	0	1	0	1	0	1	$H_{4/3}^2$
v^9	1	1	0	0	0	0	0	1	$H_{3/2}^1$

 Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5
	dim	1	1	2	3	3	1	1	1	1	2
Σ_1	1	1	0	1	1	2	0	-1	-1	0	-1
Σ_2	1	0	1	1	2	1	-1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	1	1	2	3	3	1	1	1	1	2	
v^1	0	0	0	0	1	0	0	1	0	1	$H_{3/3}^1$
v^2	0	0	0	0	1	0	0	2	1	0	$H_{4/3}^1$
v^3	0	0	0	0	1	0	1	0	0	1	$H_{3/3}^1$
v^4	0	0	0	0	1	0	1	1	1	0	$H_{4/3}^1$
v^5	0	0	0	0	1	0	2	0	1	0	$H_{4/3}^1$

v^6	0	0	0	0	1	1	0	2	0	0	$H_{4/3}^1$
v^7	0	0	0	0	1	1	1	1	0	0	$H_{4/3}^1$
v^8	0	0	0	0	1	1	2	0	0	0	$H_{4/3}^1$
v^9	0	0	0	1	0	0	0	0	1	1	$H_{3/3}^1$
v^{10}	0	0	0	1	0	0	0	1	2	0	$H_{4/3}^1$
v^{11}	0	0	0	1	0	0	1	0	2	0	$H_{4/3}^1$
v^{12}	0	0	0	1	0	1	0	0	0	1	$H_{3/3}^1$
v^{13}	0	0	0	1	0	1	0	1	1	0	$H_{4/3}^1$
v^{14}	0	0	0	1	0	1	1	0	1	0	$H_{4/3}^1$
v^{15}	0	0	0	1	0	2	0	1	0	0	$H_{4/3}^1$
v^{16}	0	0	0	1	0	2	1	0	0	0	$H_{4/3}^1$
v^{17}	0	0	0	1	1	0	0	0	0	3	$H_{5/6}^2$
v^{18}	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{19}	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{21}	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{24}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{25}	0	1	0	0	1	0	0	0	0	2	$H_{4/4}^4$
v^{26}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	1	0	0	1	0	0	0	0	0	2	$H_{4/4}^4$
v^{29}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1R_2	R_3R_3	R_4R_5
	dim	1	1	2	3	3	2	4	6
Λ_1	1	1	0	0	0	1	-1	0	-1
Λ_2	1	0	1	0	1	0	-1	0	-1
Λ_3	2	0	0	1	1	1	0	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	R_1R_2	R_3R_3	R_4R_5	
	1	1	2	3	3	2	4	6	Bipartite graph
v^1	0	0	0	1	1	0	0	1	$H_{3/4}^1$
v^2	0	0	0	1	1	1	1	0	$H_{4/4}^2$
v^3	0	0	2	0	0	0	1	0	$H_{3/2}^1$

v^4	0	1	1	0	1	0	0	1	$H_{4/4}^1$
v^5	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^6	0	2	0	0	2	1	0	1	$H_{6/6}^9$
v^7	0	2	0	0	2	2	1	0	$H_{7/6}^3$
v^8	1	0	1	1	0	0	0	1	$H_{4/4}^1$
v^9	1	0	1	1	0	1	1	0	$H_{5/4}^3$
v^{10}	1	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{11}	1	1	2	0	0	0	0	1	$H_{5/4}^1$
v^{12}	2	0	0	2	0	1	0	1	$H_{6/6}^9$
v^{13}	2	0	0	2	0	2	1	0	$H_{7/6}^3$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_1X_2	X_3X_4	X_5	R_1R_2	R_3R_3	R_4R_5	
dim	2	2	2	2	4	6	
S_1S_2	2	1	1	1	-1	-2	-3

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 19.

	X_1X_2	X_3X_4	X_5	R_1R_2	R_3R_3	R_4R_5	Bipartite graph
	2	2	2	2	4	6	
v^1	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	2	0	1	0	$H_{3/2}^1$
v^3	0	0	3	0	0	1	$H_{4/3}^1$
v^4	0	1	0	1	0	0	$H_{2/1}^1$
v^5	0	1	1	0	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	1	$H_{4/3}^1$
v^7	0	2	0	0	1	0	$H_{3/2}^1$
v^8	0	2	1	0	0	1	$H_{4/3}^1$
v^9	0	3	0	0	0	1	$H_{4/3}^1$
v^{10}	1	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	1	0	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	1	$H_{4/3}^1$
v^{13}	1	1	0	0	1	0	$H_{3/2}^1$
v^{14}	1	1	1	0	0	1	$H_{4/3}^1$
v^{15}	1	2	0	0	0	1	$H_{4/3}^1$
v^{16}	2	0	0	0	1	0	$H_{3/2}^1$
v^{17}	2	0	1	0	0	1	$H_{4/3}^1$
v^{18}	2	1	0	0	0	1	$H_{4/3}^1$
v^{19}	3	0	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_1X_2	X_3X_4	X_5	M_1	M_2	M_3	M_4	M_5
	dim	2	2	2	1	1	1	1	2
Z_1	1	0	2	1	-1	0	-1	0	-1
Z_2	1	2	0	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	X_1X_2	X_3X_4	X_5	M_1	M_2	M_3	M_4	M_5	Bipartite graph
	2	2	2	1	1	1	1	2	
v^1	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	1	0	0	0	2	0	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	1	0	2	0	0	0	0	$H_{3/2}^1$
v^9	1	0	0	0	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	0	2	0	0	0	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	2	$H_{4/4}^2$
v^{13}	1	1	0	0	0	1	1	1	$H_{5/4}^3$
v^{14}	1	1	0	0	1	1	0	1	$H_{5/4}^3$
v^{15}	1	1	0	1	0	0	1	1	$H_{5/4}^3$
v^{16}	1	1	0	1	1	0	0	1	$H_{5/4}^3$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	M_5	R_1R_2	R_3R_3	R_4R_5
	dim	1	1	1	1	2	2	4	6
T_1T_3	2	0	0	0	0	1	-1	-2	-1
T_2	1	1	0	0	1	0	0	0	-2
T_4	1	0	1	1	0	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	M_1	M_2	M_3	M_4	M_5	R_1R_2	R_3R_3	R_4R_5	
	1	1	1	1	2	2	4	6	Bipartite graph
v^1	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	0	0	2	0	1	0	$H_{3/2}^1$
v^3	0	0	2	2	1	0	0	1	$H_{6/5}^2$
v^4	0	1	1	2	1	0	0	1	$H_{6/5}^2$
v^5	0	2	0	2	1	0	0	1	$H_{6/5}^2$
v^6	1	0	2	1	1	0	0	1	$H_{6/5}^2$
v^7	1	1	1	1	1	0	0	1	$H_{6/5}^2$
v^8	1	2	0	1	1	0	0	1	$H_{6/5}^2$
v^9	2	0	2	0	1	0	0	1	$H_{6/5}^2$
v^{10}	2	1	1	0	1	0	0	1	$H_{6/5}^2$
v^{11}	2	2	0	0	1	0	0	1	$H_{6/5}^2$

SG 219

Space group $F\bar{4}3c$, No. 219, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5
	dim	1	1	2	3	3	1	1	1	1	2
Δ_1	1	1	0	1	0	1	0	-1	-1	0	0
$\Delta_2\Delta_4$	2	0	0	0	1	1	0	0	0	0	-1
Δ_3	1	0	1	1	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	X_1	X_2	X_3	X_4	X_5	
	1	1	2	3	3	1	1	1	1	2	Bipartite graph
v^1	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^4	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	1	0	0	0	1	0	1	0	$H_{3/2}^1$
v^7	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^8	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$

v^{11}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1L_2	L_3L_3
	dim	1	1	2	3	3	2	4
Λ_1	1	1	0	0	0	1	-1	0
Λ_2	1	0	1	0	1	0	-1	0
Λ_3	2	0	0	1	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	L_1L_2	L_3L_3	
	1	1	2	3	3	2	4	Bipartite graph
v^1	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	2	0	0	0	1	$H_{3/2}^1$
v^3	0	1	1	0	1	1	1	$H_{5/4}^3$
v^4	0	2	0	0	2	2	1	$H_{7/6}^3$
v^5	1	0	1	1	0	1	1	$H_{5/4}^3$
v^6	1	1	0	0	0	1	0	$H_{3/2}^1$
v^7	2	0	0	2	0	2	1	$H_{7/6}^3$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		X_1	X_2	X_3	X_4	X_5	W_1W_3	W_2W_4
	dim	1	1	1	1	2	2	2
Z_1	1	1	0	1	0	1	-2	0
Z_2	1	0	1	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	X_1	X_2	X_3	X_4	X_5	W_1W_3	W_2W_4	
	1	1	1	1	2	2	2	Bipartite graph

v^1	0	0	0	0	2	1	1	$H_{4/4}^2$
v^2	0	0	0	2	0	0	1	$H_{3/2}^1$
v^3	0	0	1	1	1	1	1	$H_{5/4}^3$
v^4	0	0	2	0	0	1	0	$H_{3/2}^1$
v^5	0	1	0	1	0	0	1	$H_{3/2}^1$
v^6	0	1	1	0	1	1	1	$H_{5/4}^3$
v^7	0	2	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	1	1	1	1	$H_{5/4}^3$
v^9	1	0	1	0	0	1	0	$H_{3/2}^1$
v^{10}	1	1	0	0	1	1	1	$H_{5/4}^3$
v^{11}	2	0	0	0	0	1	0	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	L_1L_2	L_3L_3	W_1W_3	W_2W_4
dim	2	4	2	2
Q_1Q_1	2	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_1L_2	L_3L_3	W_1W_3	W_2W_4	
	2	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	1	$H_{2/1}^1$
v^5	1	0	1	0	$H_{2/1}^1$

SG 220

Space group $I\bar{4}3d$, No. 220, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1
	dim	1	1	2	3	3	2
Σ_1	1	1	0	1	1	2	-1
Σ_2	1	0	1	1	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	N_1	
	1	1	2	3	3	2	Bipartite graph
v^1	0	0	0	1	1	3	$H_{5/6}^2$
v^2	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	1	0	0	1	2	$H_{4/4}^4$
v^4	1	0	0	1	0	2	$H_{4/4}^4$
v^5	1	1	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1H_2	H_3H_3	H_4H_5
	dim	1	1	2	3	3	2	4	6
Δ_1	1	1	0	1	0	1	0	0	-2
$\Delta_2\Delta_4$	2	0	0	0	1	1	-1	-2	-1
Δ_3	1	0	1	1	1	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	H_1H_2	H_3H_3	H_4H_5	
	1	1	2	3	3	2	4	6	Bipartite graph
v^1	0	0	0	2	2	1	1	1	$H_{7/8}^{13}$
v^2	0	0	0	2	2	3	0	1	$H_{8/8}^2$
v^3	0	0	0	4	4	0	3	2	$H_{13/16}^3, H_{13/16}^2, H_{13/16}^5, H_{13/16}^4, H_{13/16}^1$
v^4	0	0	1	1	1	1	0	1	$H_{5/6}^6$
v^5	0	0	1	3	3	0	2	2	$H_{11/14}^6, H_{11/14}^5, H_{11/14}^1, H_{11/14}^2, H_{11/14}^3, H_{11/14}^4$
v^6	0	0	2	2	2	0	1	2	$H_{9/12}^{11}, H_{9/12}^{13}, H_{9/12}^{12}, H_{9/12}^{14}$
v^7	0	0	3	1	1	0	0	2	$H_{7/10}^1$
v^8	0	1	0	1	2	0	1	1	$H_{6/7}^1$
v^9	0	1	0	1	2	2	0	1	$H_{7/7}^1$
v^{10}	0	1	1	0	1	0	0	1	$H_{4/5}^1$
v^{11}	0	2	0	0	2	1	0	1	$H_{6/6}^{10}$
v^{12}	0	4	0	0	4	0	1	2	$H_{11/12}^4, H_{11/12}^3$
v^{13}	1	0	0	2	1	0	1	1	$H_{6/7}^1$

v^{14}	1	0	0	2	1	2	0	1	$H_{7/7}^1$
v^{15}	1	0	1	1	0	0	0	1	$H_{4/5}^1$
v^{16}	1	1	0	1	1	1	0	1	$H_{6/6}^{10}$
v^{17}	1	2	0	0	1	0	0	1	$H_{5/5}^4$
v^{18}	2	0	0	2	0	1	0	1	$H_{6/6}^{10}$
v^{19}	2	1	0	1	0	0	0	1	$H_{5/5}^4$
v^{20}	4	0	0	4	0	0	1	2	$H_{11/12}^4, H_{11/12}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3
	dim	1	1	2	3	3	2	2	4
Λ_1	1	1	0	0	0	1	0	0	-1
Λ_2	1	0	1	0	1	0	0	0	-1
Λ_3	2	0	0	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_1	Γ_2	Γ_3	Γ_4	Γ_5	P_1	P_2	P_3	Bipartite graph
	1	1	2	3	3	2	2	4	
v^1	0	0	0	1	1	0	1	1	$H_{4/4}^4$
v^2	0	0	0	1	1	1	0	1	$H_{4/4}^4$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^5	0	1	0	0	1	0	0	1	$H_{3/3}^1$
v^6	1	0	0	1	0	0	0	1	$H_{3/3}^1$
v^7	1	1	1	0	0	0	0	1	$H_{4/3}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$
A matrix

corep		H_1H_2	H_3H_3	H_4H_5	N_1
	dim	2	4	6	2
G_1G_2	2	1	2	3	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	H_1H_2	H_3H_3	H_4H_5	N_1	
	2	4	6	2	Bipartite graph
v^1	0	0	1	3	$H_{4/3}^1$
v^2	0	1	0	2	$H_{3/2}^1$
v^3	1	0	0	1	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1H_2	H_3H_3	H_4H_5	P_1	P_2	P_3
	dim	2	4	6	2	2	4
F_1	1	1	0	1	0	0	-1
F_2	1	1	0	1	0	0	-1
F_3	2	0	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	H_1H_2	H_3H_3	H_4H_5	P_1	P_2	P_3	
	2	4	6	2	2	4	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/4}^2$
v^2	0	0	1	1	0	1	$H_{3/4}^2$
v^3	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	1	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	2	0	0	$H_{3/2}^1$
v^6	1	0	1	0	0	2	$H_{4/6}^1$
v^7	1	1	0	0	1	1	$H_{4/4}^4$
v^8	1	1	0	1	0	1	$H_{4/4}^4$
v^9	2	1	0	0	0	2	$H_{5/6}^2$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_1	P_2	P_3	N_1
	dim	2	2	4	2
D_1	2	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_1	P_2	P_3	N_1	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/2}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

SG 221

Space group $Pm\bar{3}m$, No. 221, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	
dim		1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
Δ_1	1	1	0	0	0	1	0	0	1	0	0	-1	0	0	-1	0	0	0	0	0	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	-1	-1	0	0	0	0	0	0	0	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	0	0	0	-1	0	0	-1	0	0	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	0	0	-1	-1	0	0	0	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	0	0	0	0	0	0	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$

v^{17}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{19}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{20}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{21}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{22}	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{23}	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{24}	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{25}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{26}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-
	dim	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2
Σ_1	1	1	0	0	0	1	0	0	1	1	1	-1	0	0	0	0	0	-1	0	0	-1
Σ_2	1	0	0	1	0	1	0	1	1	0	1	0	0	-1	0	-1	0	0	0	0	-1
Σ_3	1	0	1	0	0	0	1	1	0	1	1	0	-1	0	0	0	0	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	1	1	1	0	0	0	0	-1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 108.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	1	$H_{3/3}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	1	0	0	0	$H_{4/3}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	$H_{4/3}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/3}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	1	0	0	0	$H_{4/3}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	1	0	0	0	$H_{4/3}^1$
v^7	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	1	0	0	$H_{4/3}^1$
v^8	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	1	0	0	$H_{4/3}^1$
v^9	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	0	0	0	$H_{4/3}^1$
v^{10}	0	0	0	0	0	0	0	0	0	1	1	1	1	0	0	0	0	0	0	0	$H_{4/3}^1$

v^{11}	0 0 0 0 0 0 0 0 1 0	0 0 0 0 0 0 1 0 1 0	$H_{3/3}^1$
v^{12}	0 0 0 0 0 0 0 0 1 0	0 0 0 0 0 1 1 1 0 0	$H_{4/3}^1$
v^{13}	0 0 0 0 0 0 0 0 1 0	0 0 0 1 0 0 1 1 0 0	$H_{4/3}^1$
v^{14}	0 0 0 0 0 0 0 0 1 0	0 1 0 0 0 1 1 0 0 0	$H_{4/3}^1$
v^{15}	0 0 0 0 0 0 0 0 1 0	0 1 0 1 0 0 1 0 0 0	$H_{4/3}^1$
v^{16}	0 0 0 0 0 0 0 0 1 0	1 0 0 0 0 0 0 0 1 0	$H_{3/3}^1$
v^{17}	0 0 0 0 0 0 0 0 1 0	1 0 0 0 0 1 0 1 0 0	$H_{4/3}^1$
v^{18}	0 0 0 0 0 0 0 0 1 0	1 0 0 1 0 0 0 1 0 0	$H_{4/3}^1$
v^{19}	0 0 0 0 0 0 0 0 1 0	1 1 0 0 0 1 0 0 0 0	$H_{4/3}^1$
v^{20}	0 0 0 0 0 0 0 0 1 0	1 1 0 1 0 0 0 0 0 0	$H_{4/3}^1$
v^{21}	0 0 0 0 0 0 0 1 0 0	0 0 0 0 0 1 0 0 0 1	$H_{3/3}^1$
v^{22}	0 0 0 0 0 0 0 1 0 0	0 0 0 0 1 1 1 0 0 0	$H_{4/3}^1$
v^{23}	0 0 0 0 0 0 0 1 0 0	0 0 0 1 0 0 0 0 0 1	$H_{3/3}^1$
v^{24}	0 0 0 0 0 0 0 1 0 0	0 0 0 1 1 0 1 0 0 0	$H_{4/3}^1$
v^{25}	0 0 0 0 0 0 0 1 0 0	0 0 1 0 0 1 1 0 0 0	$H_{4/3}^1$
v^{26}	0 0 0 0 0 0 0 1 0 0	0 0 1 1 0 0 1 0 0 0	$H_{4/3}^1$
v^{27}	0 0 0 0 0 0 0 1 0 0	1 0 0 0 1 1 0 0 0 0	$H_{4/3}^1$
v^{28}	0 0 0 0 0 0 0 1 0 0	1 0 0 1 1 0 0 0 0 0	$H_{4/3}^1$
v^{29}	0 0 0 0 0 0 0 1 0 0	1 0 1 0 0 1 0 0 0 0	$H_{4/3}^1$
v^{30}	0 0 0 0 0 0 0 1 0 0	1 0 1 1 0 0 0 0 0 0	$H_{4/3}^1$
v^{31}	0 0 0 0 0 0 0 1 0 1	0 0 0 0 0 0 0 0 1 2	$H_{5/6}^2$
v^{32}	0 0 0 0 0 0 0 1 0 1	0 0 0 0 1 0 1 0 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{33}	0 0 0 0 0 0 0 1 0 1	0 0 0 0 2 0 2 0 1 0	$H_{7/6}^2$
v^{34}	0 0 0 0 0 0 0 1 0 1	0 0 1 0 0 0 1 0 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{35}	0 0 0 0 0 0 0 1 0 1	0 0 1 0 1 0 2 0 1 0	$H_{7/6}^2$
v^{36}	0 0 0 0 0 0 0 1 0 1	0 0 2 0 0 0 2 0 1 0	$H_{7/6}^2$
v^{37}	0 0 0 0 0 0 0 1 0 1	1 0 0 0 1 0 0 0 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{38}	0 0 0 0 0 0 0 1 0 1	1 0 0 0 2 0 1 0 1 0	$H_{7/6}^2$
v^{39}	0 0 0 0 0 0 0 1 0 1	1 0 1 0 0 0 0 0 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{40}	0 0 0 0 0 0 0 1 0 1	1 0 1 0 1 0 1 0 1 0	$H_{7/6}^2$
v^{41}	0 0 0 0 0 0 0 1 0 1	1 0 2 0 0 0 1 0 1 0	$H_{7/6}^2$
v^{42}	0 0 0 0 0 0 0 1 0 1	2 0 0 0 2 0 0 0 1 0	$H_{7/6}^2$
v^{43}	0 0 0 0 0 0 0 1 0 1	2 0 1 0 1 0 0 0 1 0	$H_{7/6}^2$
v^{44}	0 0 0 0 0 0 0 1 0 1	2 0 2 0 0 0 0 0 1 0	$H_{7/6}^2$
v^{45}	0 0 0 0 0 0 1 0 0 0	0 0 0 0 1 0 0 0 1 0	$H_{3/3}^1$
v^{46}	0 0 0 0 0 0 1 0 0 0	0 0 0 0 1 1 0 1 0 0	$H_{4/3}^1$
v^{47}	0 0 0 0 0 0 1 0 0 0	0 0 0 1 1 0 0 1 0 0	$H_{4/3}^1$
v^{48}	0 0 0 0 0 0 1 0 0 0	0 0 1 0 0 0 0 0 1 0	$H_{3/3}^1$
v^{49}	0 0 0 0 0 0 1 0 0 0	0 0 1 0 0 1 0 1 0 0	$H_{4/3}^1$
v^{50}	0 0 0 0 0 0 1 0 0 0	0 0 1 1 0 0 0 1 0 0	$H_{4/3}^1$
v^{51}	0 0 0 0 0 0 1 0 0 0	0 1 0 0 1 1 0 0 0 0	$H_{4/3}^1$
v^{52}	0 0 0 0 0 0 1 0 0 0	0 1 0 1 1 0 0 0 0 0	$H_{4/3}^1$

v ⁵³	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	1	0	0	0	0	0	$H_{4/3}^1$
v ⁵⁴	0	0	0	0	0	0	1	0	0	0	0	0	1	1	1	0	0	0	0	0	0	$H_{4/3}^1$
v ⁵⁵	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	2	1	$H_{5/6}^2$
v ⁵⁶	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	1	1	1	1	$H_{6/6}^A, H_{6/6}^2$
v ⁵⁷	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	2	0	2	0	1	1	$H_{7/6}^2$
v ⁵⁸	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0	0	1	1	1	1	$H_{6/6}^A, H_{6/6}^2$
v ⁵⁹	0	0	0	0	0	0	1	0	1	0	0	0	1	0	1	0	2	0	1	1	1	$H_{7/6}^2$
v ⁶⁰	0	0	0	0	0	0	1	0	1	0	0	0	0	2	0	0	0	2	0	1	1	$H_{7/6}^2$
v ⁶¹	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	1	0	0	1	1	1	$H_{6/6}^A, H_{6/6}^2$
v ⁶²	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	2	0	1	0	1	1	$H_{7/6}^2$
v ⁶³	0	0	0	0	0	0	1	0	1	0	0	1	0	1	0	0	0	0	1	1	1	$H_{6/6}^A, H_{6/6}^2$
v ⁶⁴	0	0	0	0	0	0	1	0	1	0	0	1	0	1	0	1	0	1	0	1	1	$H_{7/6}^2$
v ⁶⁵	0	0	0	0	0	0	1	0	1	0	0	1	0	2	0	0	0	1	0	1	1	$H_{7/6}^2$
v ⁶⁶	0	0	0	0	0	0	1	0	1	0	0	2	0	0	0	2	0	0	0	1	1	$H_{7/6}^2$
v ⁶⁷	0	0	0	0	0	0	1	0	1	0	0	2	0	1	0	1	0	0	0	1	1	$H_{7/6}^2$
v ⁶⁸	0	0	0	0	0	0	1	0	1	0	0	2	0	2	0	0	0	0	0	1	1	$H_{7/6}^2$
v ⁶⁹	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v ⁷⁰	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v ⁷¹	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v ⁷²	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v ⁷³	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v ⁷⁴	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v ⁷⁵	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v ⁷⁶	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v ⁷⁷	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v ⁷⁸	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v ⁷⁹	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v ⁸⁰	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v ⁸¹	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	1	1	$H_{4/4}^A$
v ⁸²	0	0	0	1	0	0	0	0	0	1	0	0	0	0	1	0	1	0	1	0	0	$H_{5/4}^2$
v ⁸³	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	1	0	1	0	0	$H_{5/4}^2$
v ⁸⁴	0	0	0	1	0	0	0	0	0	1	1	0	0	0	1	0	0	0	1	0	0	$H_{5/4}^2$
v ⁸⁵	0	0	0	1	0	0	0	0	0	1	1	0	1	0	0	0	0	0	1	0	0	$H_{5/4}^2$
v ⁸⁶	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v ⁸⁷	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v ⁸⁸	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	1	1	$H_{4/4}^A$
v ⁸⁹	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	1	1	$H_{5/4}^2$
v ⁹⁰	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	1	0	1	$H_{5/4}^2$
v ⁹¹	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	1	1	$H_{5/4}^2$
v ⁹²	0	0	1	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	1	1	$H_{5/4}^2$
v ⁹³	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v ⁹⁴	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

v^{95}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	1	$H_{4/4}^4$
v^{96}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^2$
v^{97}	0	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	1	0	$H_{5/4}^2$
v^{98}	0	1	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	1	0	$H_{5/4}^2$
v^{99}	0	1	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	1	0	$H_{5/4}^2$
v^{100}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{101}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{102}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{103}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	1	$H_{4/4}^4$
v^{104}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^2$
v^{105}	1	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	1	0	1	$H_{5/4}^2$
v^{106}	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	0	1	$H_{5/4}^2$
v^{107}	1	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	0	0	1	$H_{5/4}^2$
v^{108}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-
	dim	1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	3	3	3	3
Λ_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	0	-1	0	0	0	-1	-1	0
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1	-1	0	0	0	-1	0	0	-1
Λ_3	2	0	0	0	0	1	1	1	1	1	1	0	0	0	0	-1	-1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 84.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	3	3	3	3	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$

v^{14}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^{15}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{19}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{20}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/2}^1$
v^{21}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{22}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{23}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{24}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{25}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{26}	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{28}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{29}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{30}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	1	$H_{4/3}^2$
v^{32}	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	1	$H_{4/3}^2$
v^{33}	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	1	$H_{4/3}^2$
v^{34}	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	1	$H_{4/3}^2$
v^{35}	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	0	0	1	$H_{4/3}^2$
v^{36}	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	0	1	0	$H_{4/3}^2$
v^{37}	0	0	0	1	0	0	1	0	0	0	0	1	0	0	0	0	0	1	$H_{4/3}^2$
v^{38}	0	0	0	1	0	0	1	0	0	0	0	1	0	0	0	0	1	0	$H_{4/3}^2$
v^{39}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{40}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{41}	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{42}	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{43}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{44}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{45}	0	0	1	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{46}	0	0	1	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	$H_{4/3}^2$
v^{47}	0	0	1	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{48}	0	0	1	0	0	0	0	1	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{49}	0	0	1	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{50}	0	0	1	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	$H_{4/3}^2$
v^{51}	0	0	1	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{52}	0	0	1	0	0	0	1	0	0	1	0	0	0	0	1	0	0	0	$H_{4/3}^2$
v^{53}	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{54}	0	0	1	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{55}	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

v^{56}	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{57}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{58}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{59}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{60}	0	1	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{61}	0	1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{62}	0	1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{63}	0	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{4/3}^2$
v^{64}	0	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{4/3}^2$
v^{65}	0	1	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{66}	0	1	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{67}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{68}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{69}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{70}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{71}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{72}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{73}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{74}	1	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{75}	1	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{76}	1	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{77}	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{4/3}^2$
v^{78}	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	1	0	$H_{4/3}^2$
v^{79}	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{80}	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{81}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{82}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{83}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{84}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	2	2	3	3	3	3
S_1	1	1	0	0	0	0	0	1	0	0	1	-1	0	0	0	-1	0	0	-1	-1	-1
S_2	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0	-1	0	-1	-1	0	-1
S_3	1	0	1	0	0	0	0	0	1	1	0	0	-1	0	0	0	-1	-1	0	-1	-1
S_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1	0	-1	-1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 108.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	2	2	3	3	3	3	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0	$H_{4/4}^4$
v^6	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	0	0	1	$H_{4/4}^4$
v^7	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	1	0	0	$H_{4/4}^4$
v^8	0	0	0	0	0	0	0	0	0	1	1	1	0	0	0	0	1	0	0	0	$H_{4/4}^4$
v^9	0	0	0	0	0	0	0	0	0	1	2	0	0	0	0	0	0	1	0	1	$H_{5/6}^2$
v^{10}	0	0	0	0	0	0	0	0	0	2	1	0	0	0	0	0	1	0	1	0	$H_{5/6}^2$
v^{11}	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{13}	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/3}^1$
v^{15}	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{16}	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/3}^1$
v^{17}	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	0	0	1	0	1	0	1	0	0	0	1	0	0	0	0	1	0	$H_{5/4}^2$
v^{19}	0	0	0	0	0	1	0	1	0	1	1	0	0	0	0	0	1	0	0	0	$H_{5/4}^2$
v^{20}	0	0	0	0	0	1	0	1	1	1	1	0	0	0	0	0	1	0	1	0	$H_{6/6}^A, H_{6/6}^2$
v^{21}	0	0	0	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{22}	0	0	0	0	0	2	0	2	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{23}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{24}	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/3}^1$
v^{25}	0	0	0	0	1	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{26}	0	0	0	0	1	0	1	0	1	0	0	0	0	1	0	0	0	0	0	1	$H_{5/4}^2$
v^{27}	0	0	0	0	1	0	1	0	1	0	0	0	1	0	0	0	0	1	0	0	$H_{5/4}^2$
v^{28}	0	0	0	0	1	0	1	0	1	1	0	0	0	0	0	0	0	1	0	1	$H_{6/6}^A, H_{6/6}^2$
v^{29}	0	0	0	0	1	0	1	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{30}	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{31}	0	0	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{4/3}^1$
v^{32}	0	0	0	0	2	0	2	0	1	0	0	0	0	0	0	0	0	1	0	1	$H_{7/6}^2$
v^{33}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{34}	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/3}^1$
v^{35}	0	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{36}	0	0	0	1	0	0	0	1	0	1	0	0	1	0	0	0	0	0	1	0	$H_{5/4}^2$
v^{37}	0	0	0	1	0	0	0	1	0	1	1	0	0	0	0	0	1	0	0	0	$H_{5/4}^2$
v^{38}	0	0	0	1	0	0	0	1	1	1	0	0	0	0	0	0	1	0	1	0	$H_{6/6}^A, H_{6/6}^2$
v^{39}	0	0	0	1	0	0	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{40}	0	0	0	1	0	1	0	2	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$

v^{41}	0	0	0	1	1	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{42}	0	0	0	1	1	0	1	0	0	0	0	0	0	0	0	1	0	0	$H_{4/3}^1$
v^{43}	0	0	0	2	0	0	0	2	0	1	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{44}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{45}	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	$H_{3/3}^1$
v^{46}	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{47}	0	0	1	0	0	0	1	0	1	0	0	0	1	0	0	0	0	1	$H_{5/4}^2$
v^{48}	0	0	1	0	0	0	1	0	1	0	0	0	0	0	1	0	0	0	$H_{5/4}^2$
v^{49}	0	0	1	0	0	0	1	0	1	1	0	0	0	0	0	1	0	1	$H_{6/6}^A, H_{6/6}^2$
v^{50}	0	0	1	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{51}	0	0	1	0	0	1	0	1	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{52}	0	0	1	0	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{4/3}^1$
v^{53}	0	0	1	0	1	0	2	0	1	0	0	0	0	0	0	1	0	1	$H_{7/6}^2$
v^{54}	0	0	1	1	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{55}	0	0	1	1	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{56}	0	0	2	0	0	0	2	0	1	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{57}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{58}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{59}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{60}	0	1	0	0	0	1	0	0	0	1	0	0	1	0	0	0	1	0	$H_{5/4}^2$
v^{61}	0	1	0	0	0	1	0	0	0	1	1	0	0	0	0	1	0	0	$H_{5/4}^2$
v^{62}	0	1	0	0	0	1	0	0	1	1	0	0	0	0	1	0	1	0	$H_{6/6}^A, H_{6/6}^2$
v^{63}	0	1	0	0	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{4/3}^1$
v^{64}	0	1	0	0	0	2	0	1	0	1	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{65}	0	1	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{66}	0	1	0	0	1	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{4/3}^1$
v^{67}	0	1	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{68}	0	1	0	1	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{5/4}^2$
v^{69}	0	1	0	1	0	0	0	0	0	1	1	0	0	0	1	0	0	0	$H_{5/4}^2$
v^{70}	0	1	0	1	0	0	0	0	1	1	0	0	0	0	1	0	1	0	$H_{6/6}^A, H_{6/6}^2$
v^{71}	0	1	0	1	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{72}	0	1	0	1	0	1	0	1	0	1	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{73}	0	1	0	1	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{4/3}^1$
v^{74}	0	1	0	2	0	0	0	1	0	1	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{75}	0	1	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{76}	0	1	1	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{4/3}^1$
v^{77}	0	1	1	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{4/3}^1$
v^{78}	0	2	0	0	0	2	0	0	0	1	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{79}	0	2	0	1	0	1	0	0	0	1	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{80}	0	2	0	2	0	0	0	0	0	1	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{81}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{82}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/3}^1$

v^{83}	1	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{4/3}^1$
v^{84}	1	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{85}	1	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{5/4}^2$
v^{86}	1	0	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{5/4}^2$
v^{87}	1	0	0	0	1	0	0	0	1	1	0	0	0	0	0	0	1	0	1	$H_{6/6}^A, H_{6/6}^2$
v^{88}	1	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{89}	1	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{90}	1	0	0	0	2	0	1	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{91}	1	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	$H_{4/3}^1$
v^{92}	1	0	0	1	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{93}	1	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{94}	1	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{5/4}^2$
v^{95}	1	0	1	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{5/4}^2$
v^{96}	1	0	1	0	0	0	0	0	1	1	0	0	0	0	0	1	0	1	0	$H_{6/6}^A, H_{6/6}^2$
v^{97}	1	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{98}	1	0	1	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{99}	1	0	1	0	1	0	1	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{100}	1	0	1	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{4/3}^1$
v^{101}	1	0	2	0	0	0	1	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{102}	1	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{4/3}^1$
v^{103}	1	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{104}	1	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{4/3}^1$
v^{105}	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{106}	2	0	0	0	2	0	0	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{107}	2	0	1	0	1	0	0	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$
v^{108}	2	0	2	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	$H_{7/6}^2$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-
dim		1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2
Z_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	0	-1	0	0	0	0	-1
Z_2	1	0	0	0	1	0	0	0	1	1	0	0	0	-1	0	0	0	-1	0	0	-1
Z_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	-1	0	0	-1	0
Z_4	1	0	0	1	0	0	0	1	0	0	1	0	0	0	-1	0	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 65.

X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	
1	1	1	1	1	1	1	1	2	2	1	1	1	1	1	1	1	1	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{4/4}^2$
v^{10}	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{11}	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{12}	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	0	0	0	0	1	1	0	0	1	0	1	0	0	$H_{5/4}^3$
v^{14}	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	1	0	$H_{5/4}^3$
v^{15}	0	0	0	0	0	0	0	0	1	1	0	1	0	1	0	0	0	$H_{5/4}^3$
v^{16}	0	0	0	0	0	0	0	0	1	1	1	0	0	0	0	1	0	$H_{5/4}^3$
v^{17}	0	0	0	0	0	0	0	0	1	1	1	0	1	0	0	0	1	$H_{5/4}^3$
v^{18}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{19}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{20}	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{21}	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	0	$H_{4/3}^2$
v^{22}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{23}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{24}	0	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	1	$H_{4/3}^2$
v^{25}	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{26}	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{27}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{28}	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	0	1	$H_{4/3}^2$
v^{29}	0	0	0	0	0	1	0	0	0	1	1	0	0	0	0	0	1	$H_{4/3}^2$
v^{30}	0	0	0	0	0	1	0	1	0	1	0	0	0	0	0	0	1	$H_{5/4}^3$
v^{31}	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{32}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{33}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{34}	0	0	0	0	1	0	0	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{35}	0	0	0	0	1	0	0	0	1	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{36}	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{37}	0	0	0	0	1	0	1	0	1	0	0	0	0	0	0	0	1	$H_{5/4}^3$
v^{38}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{39}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{40}	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{41}	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{42}	0	0	0	1	0	1	0	0	0	1	0	0	0	0	0	0	1	$H_{5/4}^3$

v^{43}	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{44}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{45}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{46}	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	$H_{4/3}^2$
v^{47}	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{48}	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{49}	0	0	1	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	1	$H_{5/4}^3$
v^{50}	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{51}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{52}	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{53}	0	1	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{54}	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	1	1	$H_{5/4}^3$
v^{55}	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{56}	0	1	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	1	1	$H_{5/4}^3$
v^{57}	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{58}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{59}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{60}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	$H_{4/3}^2$
v^{61}	1	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{62}	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{63}	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	1	$H_{5/4}^3$
v^{64}	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{65}	1	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	1	$H_{5/4}^3$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	2	2	3	3	3	3
T_1	1	1	0	0	1	0	0	0	0	0	0	-1	0	0	0	-1	0	0	-1	0	0
T_2	1	0	1	1	0	0	0	0	0	0	0	0	-1	0	0	-1	-1	0	0	0	0
T_3	1	0	0	0	0	1	0	0	1	0	0	0	0	-1	0	-1	0	0	0	0	-1
T_4	1	0	0	0	0	0	1	1	0	0	0	0	0	0	-1	0	-1	0	0	-1	0
T_5	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-	Bipartite graph
v^1	1	1	1	1	1	1	1	1	2	2	1	1	1	1	2	2	3	3	3	3	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$

v^3	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^8	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^9	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	0	0	0	1	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	1	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{21}	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{22}	0	0	1	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{24}	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{25}	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{26}	0	1	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{27}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{28}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{30}	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{31}	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{32}	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$

SG 222

Space group $Pn\bar{3}n$, No. 222, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3X_4
dim	1	1	1	1	2	2	3	3	3	3	2	2	4

Δ_1	1	1	0	0	0	1	0	0	1	0	0	-1	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	-1	0	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	-1	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	-1	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 29.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3X_4	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	2	2	4	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	1	1	0	0	1	0	1	$H_{4/4}^2$
v^3	0	0	0	0	0	1	0	1	0	1	1	1	1	$H_{6/6}^5$
v^4	0	0	0	0	1	0	1	0	1	0	1	1	1	$H_{6/6}^5$
v^5	0	0	0	0	1	1	0	0	0	0	1	1	0	$H_{4/4}^2$
v^6	0	0	0	2	0	0	0	0	0	2	0	2	1	$H_{7/6}^3$
v^7	0	0	0	2	1	0	1	0	0	1	1	2	1	$H_{9/8}^1$
v^8	0	0	0	2	2	0	2	0	0	0	2	2	1	$H_{11/10}^1$
v^9	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	0	2	0	0	0	0	0	2	0	0	2	1	$H_{7/6}^3$
v^{11}	0	0	2	0	0	1	0	1	1	0	1	2	1	$H_{9/8}^1$
v^{12}	0	0	2	0	0	2	0	2	0	0	2	2	1	$H_{11/10}^1$
v^{13}	0	1	0	1	0	0	0	1	0	1	1	1	1	$H_{7/6}^3$
v^{14}	0	1	0	1	1	0	0	0	0	0	1	1	0	$H_{5/4}^3$
v^{15}	0	1	1	0	0	0	0	1	1	0	1	1	1	$H_{7/6}^3$
v^{16}	0	1	1	0	0	1	0	2	0	0	2	1	1	$H_{9/8}^1$
v^{17}	0	1	1	0	1	0	0	0	2	0	1	2	1	$H_{9/8}^1$
v^{18}	0	2	0	0	0	0	0	2	0	0	2	0	1	$H_{7/6}^3$
v^{19}	0	2	0	0	1	0	0	1	1	0	2	1	1	$H_{9/8}^1$
v^{20}	0	2	0	0	2	0	0	0	2	0	2	2	1	$H_{11/10}^1$
v^{21}	1	0	0	1	0	0	1	0	0	1	1	1	1	$H_{7/6}^3$
v^{22}	1	0	0	1	0	1	0	0	0	2	1	2	1	$H_{9/8}^1$
v^{23}	1	0	0	1	1	0	2	0	0	0	2	1	1	$H_{9/8}^1$
v^{24}	1	0	1	0	0	0	1	0	1	0	1	1	1	$H_{7/6}^3$
v^{25}	1	0	1	0	0	1	0	0	0	0	1	1	0	$H_{5/4}^3$
v^{26}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{27}	2	0	0	0	0	0	2	0	0	0	2	0	1	$H_{7/6}^3$
v^{28}	2	0	0	0	0	1	1	0	0	1	2	1	1	$H_{9/8}^1$
v^{29}	2	0	0	0	0	2	0	0	0	2	2	2	1	$H_{11/10}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
	dim	1	1	1	1	2	2	3	3	3	3	2	2	2	2
Σ_1	1	1	0	0	0	1	0	0	1	1	1	-1	-1	0	0
Σ_2	1	0	0	1	0	1	0	1	1	0	1	0	0	-1	-1
Σ_3	1	0	1	0	0	0	1	1	0	1	1	-1	0	0	-1
Σ_4	1	0	0	0	1	0	1	1	1	1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 64.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	0	1	$H_{5/6}^2, H_{5/6}^4$
v^2	0	0	0	0	0	0	0	0	1	1	2	0	1	0	$H_{5/6}^2$
v^3	0	0	0	0	0	0	0	1	1	0	0	2	0	1	$H_{5/6}^2$
v^4	0	0	0	0	0	0	0	1	1	0	1	1	1	0	$H_{5/6}^4, H_{5/6}^2$
v^5	0	0	0	0	0	0	1	0	0	1	0	1	0	2	$H_{5/6}^2$
v^6	0	0	0	0	0	0	1	0	0	1	1	0	1	1	$H_{5/6}^2, H_{5/6}^4$
v^7	0	0	0	0	0	0	1	1	0	0	0	1	1	1	$H_{5/6}^4, H_{5/6}^2$
v^8	0	0	0	0	0	0	1	1	0	0	1	0	2	0	$H_{5/6}^2$
v^9	0	0	0	0	0	1	0	0	0	2	1	1	0	2	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{10}	0	0	0	0	0	1	0	0	0	2	2	0	1	1	$H_{7/8}^{10}, H_{7/8}^{12}, H_{7/8}^3$
v^{11}	0	0	0	0	0	1	0	1	0	1	0	2	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{12}	0	0	0	0	0	1	0	1	0	1	1	1	1	1	$H_{7/8}^3, H_{7/8}^{12}, H_{7/8}^{10}$
v^{13}	0	0	0	0	0	1	0	1	0	1	2	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{14}	0	0	0	0	0	1	0	2	0	0	0	2	1	1	$H_{7/8}^{12}, H_{7/8}^{10}, H_{7/8}^3$
v^{15}	0	0	0	0	0	1	0	2	0	0	1	1	2	0	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{16}	0	0	0	0	1	0	0	0	2	0	1	2	0	1	$H_{7/8}^3, H_{7/8}^{12}, H_{7/8}^{10}$
v^{17}	0	0	0	0	1	0	0	0	2	0	2	1	1	0	$H_{7/8}^{12}, H_{7/8}^{10}, H_{7/8}^3$
v^{18}	0	0	0	0	1	0	1	0	1	0	0	2	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{19}	0	0	0	0	1	0	1	0	1	0	1	1	1	1	$H_{7/8}^{12}, H_{7/8}^{10}, H_{7/8}^3$
v^{20}	0	0	0	0	1	0	1	0	1	0	2	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{21}	0	0	0	0	1	0	2	0	0	0	0	1	1	2	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{22}	0	0	0	0	1	0	2	0	0	0	1	0	2	1	$H_{7/8}^{10}, H_{7/8}^3, H_{7/8}^{12}$
v^{23}	0	0	0	0	1	1	0	0	0	0	0	1	0	1	$H_{4/4}^2$
v^{24}	0	0	0	0	1	1	0	0	0	0	1	0	1	0	$H_{4/4}^2$
v^{25}	0	0	0	1	0	0	0	0	0	1	0	1	0	1	$H_{4/4}^4$
v^{26}	0	0	0	1	0	0	0	0	0	1	1	0	1	0	$H_{4/4}^4$
v^{27}	0	0	0	1	0	0	0	1	0	0	0	1	1	0	$H_{4/4}^4$
v^{28}	0	0	0	1	1	0	0	0	1	0	0	2	0	1	$H_{6/6}^8, H_{6/6}^7$

v^{29}	0	0	0	1	1	0	0	0	1	0	1	1	1	0	$H_{6/6}^7, H_{6/6}^8$
v^{30}	0	0	0	1	1	0	1	0	0	0	0	1	1	1	$H_{6/6}^8, H_{6/6}^7$
v^{31}	0	0	0	1	1	0	1	0	0	0	1	0	2	0	$H_{6/6}^8, H_{6/6}^7$
v^{32}	0	0	0	2	1	0	0	0	0	0	0	1	1	0	$H_{5/4}^3$
v^{33}	0	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{4/4}^4$
v^{34}	0	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{4/4}^4$
v^{35}	0	0	1	0	0	0	1	0	0	0	0	0	1	1	$H_{4/4}^4$
v^{36}	0	0	1	0	0	1	0	0	0	1	0	1	0	2	$H_{6/6}^8, H_{6/6}^7$
v^{37}	0	0	1	0	0	1	0	0	0	1	1	0	1	1	$H_{6/6}^8, H_{6/6}^7$
v^{38}	0	0	1	0	0	1	0	1	0	0	0	1	1	1	$H_{6/6}^7, H_{6/6}^8$
v^{39}	0	0	1	0	0	1	0	1	0	0	1	0	2	0	$H_{6/6}^8, H_{6/6}^7$
v^{40}	0	0	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{41}	0	0	2	0	0	1	0	0	0	0	0	0	1	1	$H_{5/4}^3$
v^{42}	0	1	0	0	0	0	0	0	0	1	1	0	0	1	$H_{4/4}^4$
v^{43}	0	1	0	0	0	0	0	1	0	0	0	1	0	1	$H_{4/4}^4$
v^{44}	0	1	0	0	0	0	0	1	0	0	1	0	1	0	$H_{4/4}^4$
v^{45}	0	1	0	0	1	0	0	0	1	0	1	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{46}	0	1	0	0	1	0	0	0	1	0	2	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{47}	0	1	0	0	1	0	1	0	0	0	0	1	0	2	$H_{6/6}^8, H_{6/6}^7$
v^{48}	0	1	0	0	1	0	1	0	0	0	1	0	1	1	$H_{6/6}^8, H_{6/6}^7$
v^{49}	0	1	0	1	1	0	0	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{50}	0	1	0	1	1	0	0	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{51}	0	1	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{52}	0	2	0	0	1	0	0	0	0	0	1	0	0	1	$H_{5/4}^3$
v^{53}	1	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{4/4}^4$
v^{54}	1	0	0	0	0	0	1	0	0	0	0	1	0	1	$H_{4/4}^4$
v^{55}	1	0	0	0	0	0	1	0	0	0	1	0	1	0	$H_{4/4}^4$
v^{56}	1	0	0	0	0	1	0	0	0	1	1	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{57}	1	0	0	0	0	1	0	0	0	1	2	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{58}	1	0	0	0	0	1	0	1	0	0	0	2	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{59}	1	0	0	0	0	1	0	1	0	0	1	1	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{60}	1	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{61}	1	0	1	0	0	1	0	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{62}	1	0	1	0	0	1	0	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{63}	1	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{64}	2	0	0	0	0	1	0	0	0	0	1	1	0	0	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1	R_2R_3	R_4
dim		1	1	1	1	2	2	3	3	3	3	2	4	6
Λ_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	-1
Λ_2	1	0	1	1	0	0	0	1	0	0	1	-1	0	-1
Λ_3	2	0	0	0	0	1	1	1	1	1	1	0	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 95.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1	R_2R_3	R_4	Bipartite graph
v^1	1	1	1	1	2	2	3	3	3	3	2	4	6	$H_{3/4}^1$
v^2	0	0	0	0	0	0	0	0	1	1	1	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	1	0	1	0	0	1	$H_{3/4}^1$
v^4	0	0	0	0	0	0	0	1	0	1	1	1	0	$H_{4/4}^2$
v^5	0	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/4}^1$
v^6	0	0	0	0	0	0	1	0	1	0	1	1	0	$H_{4/4}^2$
v^7	0	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/4}^1$
v^8	0	0	0	0	0	0	1	1	0	0	1	1	0	$H_{4/4}^2$
v^9	0	0	0	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	0	0	1	0	1	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{13}	0	0	0	1	0	1	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{14}	0	0	0	1	0	1	1	0	0	0	0	0	1	$H_{4/4}^1$
v^{15}	0	0	0	1	0	1	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{16}	0	0	0	1	1	0	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{17}	0	0	0	1	1	0	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{18}	0	0	0	1	1	0	1	0	0	0	0	0	1	$H_{4/4}^1$
v^{19}	0	0	0	1	1	0	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{20}	0	0	0	2	0	0	0	0	0	2	1	0	1	$H_{6/6}^9$
v^{21}	0	0	0	2	0	0	0	0	0	2	2	1	0	$H_{7/6}^3$
v^{22}	0	0	0	2	0	0	1	0	0	1	1	0	1	$H_{6/6}^9$
v^{23}	0	0	0	2	0	0	1	0	0	1	2	1	0	$H_{7/6}^3$
v^{24}	0	0	0	2	0	0	2	0	0	0	1	0	1	$H_{6/6}^9$
v^{25}	0	0	0	2	0	0	2	0	0	0	2	1	0	$H_{7/6}^3$
v^{26}	0	0	1	0	0	1	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{27}	0	0	1	0	0	1	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{28}	0	0	1	0	0	1	0	1	0	0	0	0	1	$H_{4/4}^1$
v^{29}	0	0	1	0	0	1	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{30}	0	0	1	0	1	0	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{31}	0	0	1	0	1	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{32}	0	0	1	0	1	0	0	1	0	0	0	0	1	$H_{4/4}^1$

v^{33}	0	0	1	0	1	0	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{34}	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{35}	0	0	1	1	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{36}	0	0	1	1	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{37}	0	0	1	1	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{38}	0	0	2	0	0	0	0	0	2	0	1	0	1	$H_{6/6}^9$
v^{39}	0	0	2	0	0	0	0	0	2	0	2	1	0	$H_{7/6}^3$
v^{40}	0	0	2	0	0	0	0	1	1	0	1	0	1	$H_{6/6}^9$
v^{41}	0	0	2	0	0	0	0	1	1	0	2	1	0	$H_{7/6}^3$
v^{42}	0	0	2	0	0	0	0	2	0	0	1	0	1	$H_{6/6}^9$
v^{43}	0	0	2	0	0	0	0	2	0	0	2	1	0	$H_{7/6}^3$
v^{44}	0	1	0	0	0	1	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{45}	0	1	0	0	0	1	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{46}	0	1	0	0	0	1	0	1	0	0	0	0	1	$H_{4/4}^1$
v^{47}	0	1	0	0	0	1	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{48}	0	1	0	0	1	0	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{49}	0	1	0	0	1	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{50}	0	1	0	0	1	0	0	1	0	0	0	0	1	$H_{4/4}^1$
v^{51}	0	1	0	0	1	0	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{52}	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{53}	0	1	0	1	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{54}	0	1	0	1	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{55}	0	1	0	1	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{56}	0	1	1	0	0	0	0	0	2	0	1	0	1	$H_{6/6}^9$
v^{57}	0	1	1	0	0	0	0	0	2	0	2	1	0	$H_{7/6}^3$
v^{58}	0	1	1	0	0	0	0	1	1	0	1	0	1	$H_{6/6}^9$
v^{59}	0	1	1	0	0	0	0	1	1	0	2	1	0	$H_{7/6}^3$
v^{60}	0	1	1	0	0	0	0	2	0	0	1	0	1	$H_{6/6}^9$
v^{61}	0	1	1	0	0	0	0	2	0	0	2	1	0	$H_{7/6}^3$
v^{62}	0	2	0	0	0	0	0	0	2	0	1	0	1	$H_{6/6}^9$
v^{63}	0	2	0	0	0	0	0	0	2	0	2	1	0	$H_{7/6}^3$
v^{64}	0	2	0	0	0	0	0	1	1	0	1	0	1	$H_{6/6}^9$
v^{65}	0	2	0	0	0	0	0	1	1	0	2	1	0	$H_{7/6}^3$
v^{66}	0	2	0	0	0	0	0	2	0	0	1	0	1	$H_{6/6}^9$
v^{67}	0	2	0	0	0	0	0	2	0	0	2	1	0	$H_{7/6}^3$
v^{68}	1	0	0	0	0	1	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{69}	1	0	0	0	0	1	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{70}	1	0	0	0	0	1	1	0	0	0	0	0	1	$H_{4/4}^1$
v^{71}	1	0	0	0	0	1	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{72}	1	0	0	0	1	0	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{73}	1	0	0	0	1	0	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{74}	1	0	0	0	1	0	1	0	0	0	0	0	1	$H_{4/4}^1$

v^{75}	1	0	0	0	1	0	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{76}	1	0	0	1	0	0	0	0	0	2	1	0	1	$H_{6/6}^9$
v^{77}	1	0	0	1	0	0	0	0	0	2	2	1	0	$H_{7/6}^3$
v^{78}	1	0	0	1	0	0	1	0	0	1	1	0	1	$H_{6/6}^9$
v^{79}	1	0	0	1	0	0	1	0	0	1	2	1	0	$H_{7/6}^3$
v^{80}	1	0	0	1	0	0	2	0	0	0	1	0	1	$H_{6/6}^9$
v^{81}	1	0	0	1	0	0	2	0	0	0	2	1	0	$H_{7/6}^3$
v^{82}	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{83}	1	0	1	0	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{84}	1	0	1	0	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{85}	1	0	1	0	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{86}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{87}	1	1	0	0	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{88}	1	1	0	0	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{89}	1	1	0	0	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{90}	2	0	0	0	0	0	0	0	0	2	1	0	1	$H_{6/6}^9$
v^{91}	2	0	0	0	0	0	0	0	0	2	2	1	0	$H_{7/6}^3$
v^{92}	2	0	0	0	0	0	1	0	0	1	1	0	1	$H_{6/6}^9$
v^{93}	2	0	0	0	0	0	1	0	0	1	2	1	0	$H_{7/6}^3$
v^{94}	2	0	0	0	0	0	2	0	0	0	1	0	1	$H_{6/6}^9$
v^{95}	2	0	0	0	0	0	2	0	0	0	2	1	0	$H_{7/6}^3$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_1	X_2	X_3X_4	R_1	R_2R_3	R_4	
dim	2	2	4	2	4	6	
S_1	2	1	1	2	-1	-2	-3

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 15.

	X_1	X_2	X_3X_4	R_1	R_2R_3	R_4	Bipartite graph
	2	2	4	2	4	6	
v^1	0	0	1	0	1	0	$H_{2/2}^1$
v^2	0	0	1	2	0	0	$H_{3/2}^1$
v^3	0	0	2	1	0	1	$H_{4/4}^4$
v^4	0	0	3	0	0	2	$H_{5/6}^2$
v^5	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	1	1	0	0	1	$H_{3/3}^1$
v^7	0	2	0	0	1	0	$H_{3/2}^1$
v^8	0	3	0	0	0	1	$H_{4/3}^1$

v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	0	1	$H_{3/3}^1$
v^{11}	1	1	0	0	1	0	$H_{3/2}^1$
v^{12}	1	2	0	0	0	1	$H_{4/3}^1$
v^{13}	2	0	0	0	1	0	$H_{3/2}^1$
v^{14}	2	1	0	0	0	1	$H_{4/3}^1$
v^{15}	3	0	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_1	X_2	X_3X_4	M_1	M_2	M_3	M_4
dim	2	2	4	2	2	2	2
Z_1	2	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	X_1	X_2	X_3X_4	M_1	M_2	M_3	M_4	Bipartite graph
	2	2	4	2	2	2	2	
v^1	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	0	1	0	1	0	1	$H_{3/2}^1$
v^5	0	0	1	0	1	1	0	$H_{3/2}^1$
v^6	0	0	1	0	2	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	1	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	1	2	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{16}	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{17}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	1	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_1	M_2	M_3	M_4	R_1	R_2R_3	R_4
	dim	2	2	2	2	2	4	6
T_1	1	0	0	0	1	0	0	-1
T_2	1	0	0	0	1	0	0	-1
T_3	1	0	1	0	0	0	0	-1
T_4	1	0	1	0	0	0	0	-1
T_5	2	1	0	1	0	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	M_1	M_2	M_3	M_4	R_1	R_2R_3	R_4	
	2	2	2	2	2	4	6	Bipartite graph
v^1	0	0	1	0	1	0	0	$H_{2/1}^1$
v^2	0	0	2	0	0	1	0	$H_{3/2}^1$
v^3	0	1	1	1	0	0	1	$H_{4/5}^1$
v^4	1	0	0	0	1	0	0	$H_{2/1}^1$
v^5	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	1	1	0	1	0	0	1	$H_{4/5}^1$
v^7	2	0	0	0	0	1	0	$H_{3/2}^1$

SG 223

Space group $Pm\bar{3}n$, No. 223, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	2	3	3	3	3	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	1	0	0	-1	0	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	0	-1	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	-1	0	0	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	-1	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4	
	1	1	1	1	2	2	3	3	3	3	2	2	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	1	0	1	1	0	0	2	$H_{5/4}^3$
v^2	0	0	0	0	0	0	0	1	0	1	1	1	0	1	$H_{5/4}^3$
v^3	0	0	0	0	0	0	0	1	0	1	1	2	0	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	0	1	0	0	0	1	2	$H_{5/4}^3$
v^5	0	0	0	0	0	0	1	0	1	0	0	1	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	0	1	0	1	0	0	2	1	0	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^8	0	0	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/2}^1$
v^9	0	0	0	1	0	0	1	0	0	0	0	0	1	1	$H_{4/3}^2$
v^{10}	0	0	0	1	0	0	1	0	0	0	0	1	1	0	$H_{4/3}^2$
v^{11}	0	0	1	0	0	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^{12}	0	0	1	0	0	0	0	1	0	0	1	1	0	0	$H_{4/3}^2$
v^{13}	0	1	0	0	0	0	0	0	1	0	0	0	1	1	$H_{4/3}^2$
v^{14}	0	1	0	0	0	0	0	0	1	0	0	1	1	0	$H_{4/3}^2$
v^{15}	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	0	0	0	0	0	0	0	1	1	0	0	1	$H_{4/3}^2$
v^{17}	1	0	0	0	0	0	0	0	0	1	1	1	0	0	$H_{4/3}^2$
v^{18}	1	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	
dim	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
Σ_1	1	1	0	0	0	1	0	0	1	1	1	0	-1	0	0	0	0	0	-1	-1	0
Σ_2	1	0	0	1	0	1	0	1	1	0	1	0	0	0	-1	0	-1	0	0	-1	0
Σ_3	1	0	1	0	0	0	1	1	0	1	1	-1	0	0	0	0	0	-1	0	0	-1
Σ_4	1	0	0	0	1	0	1	1	1	1	0	0	0	-1	0	-1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 108.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	$H_{3/3}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	1	0	0	$H_{4/3}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	1	0	0	$H_{4/3}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	1	0	0	0	$H_{4/3}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	1	0	0	0	$H_{4/3}^1$
v^6	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/3}^1$
v^7	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	1	0	0	0	$H_{4/3}^1$

v^8	0 0 0 0 0 0 0 0 0 1	1 0 0 1 0 0 0 1 0 0	$H_{4/3}^1$
v^9	0 0 0 0 0 0 0 0 0 1	1 1 0 0 0 1 0 0 0 0	$H_{4/3}^1$
v^{10}	0 0 0 0 0 0 0 0 0 1	1 1 0 1 0 0 0 0 0 0	$H_{4/3}^1$
v^{11}	0 0 0 0 0 0 0 0 1 0	0 0 0 0 0 0 0 1 0 1	$H_{3/3}^1$
v^{12}	0 0 0 0 0 0 0 0 1 0	0 0 0 0 1 0 1 1 0 0	$H_{4/3}^1$
v^{13}	0 0 0 0 0 0 0 0 1 0	0 0 1 0 0 0 1 1 0 0	$H_{4/3}^1$
v^{14}	0 0 0 0 0 0 0 0 1 0	0 1 0 0 0 0 0 0 0 1	$H_{3/3}^1$
v^{15}	0 0 0 0 0 0 0 0 1 0	0 1 0 0 1 0 1 0 0 0	$H_{4/3}^1$
v^{16}	0 0 0 0 0 0 0 0 1 0	0 1 1 0 0 0 1 0 0 0	$H_{4/3}^1$
v^{17}	0 0 0 0 0 0 0 0 1 0	1 0 0 0 1 0 0 1 0 0	$H_{4/3}^1$
v^{18}	0 0 0 0 0 0 0 0 1 0	1 0 1 0 0 0 0 1 0 0	$H_{4/3}^1$
v^{19}	0 0 0 0 0 0 0 0 1 0	1 1 0 0 1 0 0 0 0 0	$H_{4/3}^1$
v^{20}	0 0 0 0 0 0 0 0 1 0	1 1 1 0 0 0 0 0 0 0	$H_{4/3}^1$
v^{21}	0 0 0 0 0 0 0 1 0 0	0 0 0 0 1 0 0 0 1 0	$H_{3/3}^1$
v^{22}	0 0 0 0 0 0 0 1 0 0	0 0 0 0 1 1 0 1 0 0	$H_{4/3}^1$
v^{23}	0 0 0 0 0 0 0 1 0 0	0 0 0 1 1 0 0 1 0 0	$H_{4/3}^1$
v^{24}	0 0 0 0 0 0 0 1 0 0	0 0 1 0 0 0 0 0 1 0	$H_{3/3}^1$
v^{25}	0 0 0 0 0 0 0 1 0 0	0 0 1 0 0 1 0 1 0 0	$H_{4/3}^1$
v^{26}	0 0 0 0 0 0 0 1 0 0	0 0 1 1 0 0 0 1 0 0	$H_{4/3}^1$
v^{27}	0 0 0 0 0 0 0 1 0 0	0 1 0 0 1 1 0 0 0 0	$H_{4/3}^1$
v^{28}	0 0 0 0 0 0 0 1 0 0	0 1 0 1 1 0 0 0 0 0	$H_{4/3}^1$
v^{29}	0 0 0 0 0 0 0 1 0 0	0 1 1 0 0 1 0 0 0 0	$H_{4/3}^1$
v^{30}	0 0 0 0 0 0 0 1 0 0	0 1 1 1 0 0 0 0 0 0	$H_{4/3}^1$
v^{31}	0 0 0 0 0 0 0 1 0 1	0 0 0 0 0 0 0 0 2 1	$H_{5/6}^2$
v^{32}	0 0 0 0 0 0 0 1 0 1	0 0 0 0 0 1 0 1 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{33}	0 0 0 0 0 0 0 1 0 1	0 0 0 0 0 2 0 2 0 1	$H_{7/6}^2$
v^{34}	0 0 0 0 0 0 0 1 0 1	0 0 0 1 0 0 0 1 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{35}	0 0 0 0 0 0 0 1 0 1	0 0 0 1 0 1 0 2 0 1	$H_{7/6}^2$
v^{36}	0 0 0 0 0 0 0 1 0 1	0 0 0 2 0 0 0 2 0 1	$H_{7/6}^2$
v^{37}	0 0 0 0 0 0 0 1 0 1	0 1 0 0 0 1 0 0 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{38}	0 0 0 0 0 0 0 1 0 1	0 1 0 0 0 2 0 1 0 1	$H_{7/6}^2$
v^{39}	0 0 0 0 0 0 0 1 0 1	0 1 0 1 0 0 0 0 1 1	$H_{6/6}^A, H_{6/6}^2$
v^{40}	0 0 0 0 0 0 0 1 0 1	0 1 0 1 0 1 0 1 0 1	$H_{7/6}^2$
v^{41}	0 0 0 0 0 0 0 1 0 1	0 1 0 2 0 0 0 1 0 1	$H_{7/6}^2$
v^{42}	0 0 0 0 0 0 0 1 0 1	0 2 0 0 0 2 0 0 0 1	$H_{7/6}^2$
v^{43}	0 0 0 0 0 0 0 1 0 1	0 2 0 1 0 1 0 0 0 1	$H_{7/6}^2$
v^{44}	0 0 0 0 0 0 0 1 0 1	0 2 0 2 0 0 0 0 0 1	$H_{7/6}^2$
v^{45}	0 0 0 0 0 0 1 0 0 0	0 0 0 0 0 1 0 0 0 1	$H_{3/3}^1$
v^{46}	0 0 0 0 0 0 1 0 0 0	0 0 0 0 1 1 1 0 0 0	$H_{4/3}^1$
v^{47}	0 0 0 0 0 0 1 0 0 0	0 0 0 1 0 0 0 0 0 1	$H_{3/3}^1$
v^{48}	0 0 0 0 0 0 1 0 0 0	0 0 0 1 1 0 1 0 0 0	$H_{4/3}^1$
v^{49}	0 0 0 0 0 0 1 0 0 0	0 0 1 0 0 1 1 0 0 0	$H_{4/3}^1$

v ⁵⁰	0 0 0 0 0 0 1 0 0 0	0 0 1 1 0 0 1 0 0 0	$H_{4/3}^1$
v ⁵¹	0 0 0 0 0 0 1 0 0 0	1 0 0 0 1 1 0 0 0 0	$H_{4/3}^1$
v ⁵²	0 0 0 0 0 0 1 0 0 0	1 0 0 1 1 0 0 0 0 0	$H_{4/3}^1$
v ⁵³	0 0 0 0 0 0 1 0 0 0	1 0 1 0 0 1 0 0 0 0	$H_{4/3}^1$
v ⁵⁴	0 0 0 0 0 0 1 0 0 0	1 0 1 1 0 0 0 0 0 0	$H_{4/3}^1$
v ⁵⁵	0 0 0 0 0 0 1 0 1 0	0 0 0 0 0 0 0 0 1 2	$H_{5/6}^2$
v ⁵⁶	0 0 0 0 0 0 1 0 1 0	0 0 0 0 1 0 1 0 1 1	$H_{6/6}^4, H_{6/6}^2$
v ⁵⁷	0 0 0 0 0 0 1 0 1 0	0 0 0 0 2 0 2 0 1 0	$H_{7/6}^2$
v ⁵⁸	0 0 0 0 0 0 1 0 1 0	0 0 1 0 0 0 1 0 1 1	$H_{6/6}^4, H_{6/6}^2$
v ⁵⁹	0 0 0 0 0 0 1 0 1 0	0 0 1 0 1 0 2 0 1 0	$H_{7/6}^2$
v ⁶⁰	0 0 0 0 0 0 1 0 1 0	0 0 2 0 0 0 2 0 1 0	$H_{7/6}^2$
v ⁶¹	0 0 0 0 0 0 1 0 1 0	1 0 0 0 1 0 0 0 1 1	$H_{6/6}^4, H_{6/6}^2$
v ⁶²	0 0 0 0 0 0 1 0 1 0	1 0 0 0 2 0 1 0 1 0	$H_{7/6}^2$
v ⁶³	0 0 0 0 0 0 1 0 1 0	1 0 1 0 0 0 0 0 1 1	$H_{6/6}^4, H_{6/6}^2$
v ⁶⁴	0 0 0 0 0 0 1 0 1 0	1 0 1 0 1 0 1 0 1 0	$H_{7/6}^2$
v ⁶⁵	0 0 0 0 0 0 1 0 1 0	1 0 2 0 0 0 1 0 1 0	$H_{7/6}^2$
v ⁶⁶	0 0 0 0 0 0 1 0 1 0	2 0 0 0 2 0 0 0 1 0	$H_{7/6}^2$
v ⁶⁷	0 0 0 0 0 0 1 0 1 0	2 0 1 0 1 0 0 0 1 0	$H_{7/6}^2$
v ⁶⁸	0 0 0 0 0 0 1 0 1 0	2 0 2 0 0 0 0 0 1 0	$H_{7/6}^2$
v ⁶⁹	0 0 0 0 0 1 0 0 0 0	0 0 0 0 0 0 0 0 0 1	$H_{2/2}^1$
v ⁷⁰	0 0 0 0 0 1 0 0 0 0	0 0 0 0 1 0 1 0 0 0	$H_{3/2}^1$
v ⁷¹	0 0 0 0 0 1 0 0 0 0	0 0 1 0 0 0 1 0 0 0	$H_{3/2}^1$
v ⁷²	0 0 0 0 0 1 0 0 0 0	1 0 0 0 1 0 0 0 0 0	$H_{3/2}^1$
v ⁷³	0 0 0 0 0 1 0 0 0 0	1 0 1 0 0 0 0 0 0 0	$H_{3/2}^1$
v ⁷⁴	0 0 0 0 1 0 0 0 0 0	0 0 0 0 0 0 0 0 1 0	$H_{2/2}^1$
v ⁷⁵	0 0 0 0 1 0 0 0 0 0	0 0 0 0 0 1 0 1 0 0	$H_{3/2}^1$
v ⁷⁶	0 0 0 0 1 0 0 0 0 0	0 0 0 1 0 0 0 1 0 0	$H_{3/2}^1$
v ⁷⁷	0 0 0 0 1 0 0 0 0 0	0 1 0 0 0 1 0 0 0 0	$H_{3/2}^1$
v ⁷⁸	0 0 0 0 1 0 0 0 0 0	0 1 0 1 0 0 0 0 0 0	$H_{3/2}^1$
v ⁷⁹	0 0 0 1 0 0 0 0 0 0	0 0 0 0 1 0 0 0 0 0	$H_{2/1}^1$
v ⁸⁰	0 0 0 1 0 0 0 0 0 0	0 0 1 0 0 0 0 0 0 0	$H_{2/1}^1$
v ⁸¹	0 0 0 1 0 0 0 0 0 1	0 0 0 0 0 0 0 0 1 1	$H_{4/4}^4$
v ⁸²	0 0 0 1 0 0 0 0 0 1	0 0 0 0 0 1 0 1 0 1	$H_{5/4}^2$
v ⁸³	0 0 0 1 0 0 0 0 0 1	0 0 0 1 0 0 0 1 0 1	$H_{5/4}^2$
v ⁸⁴	0 0 0 1 0 0 0 0 0 1	0 1 0 0 0 1 0 0 0 1	$H_{5/4}^2$
v ⁸⁵	0 0 0 1 0 0 0 0 0 1	0 1 0 1 0 0 0 0 0 1	$H_{5/4}^2$
v ⁸⁶	0 0 1 0 0 0 0 0 0 0	0 0 0 0 0 1 0 0 0 0	$H_{2/1}^1$
v ⁸⁷	0 0 1 0 0 0 0 0 0 0	0 0 0 1 0 0 0 0 0 0	$H_{2/1}^1$
v ⁸⁸	0 0 1 0 0 0 0 0 1 0	0 0 0 0 0 0 0 0 1 1	$H_{4/4}^4$
v ⁸⁹	0 0 1 0 0 0 0 0 1 0	0 0 0 0 1 0 1 0 1 0	$H_{5/4}^2$
v ⁹⁰	0 0 1 0 0 0 0 0 1 0	0 0 1 0 0 0 1 0 1 0	$H_{5/4}^2$
v ⁹¹	0 0 1 0 0 0 0 0 1 0	1 0 0 0 1 0 0 0 1 0	$H_{5/4}^2$

v^{92}	0	0	1	0	0	0	0	0	0	1	0	1	0	1	0	0	0	1	0	$H_{5/4}^2$
v^{93}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{94}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{95}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	1	$H_{4/4}^4$
v^{96}	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^2$
v^{97}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	$H_{5/4}^2$
v^{98}	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	0	1	$H_{5/4}^2$
v^{99}	0	1	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	1	$H_{5/4}^2$
v^{100}	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{101}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{102}	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{103}	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	1	$H_{4/4}^4$
v^{104}	1	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	1	0	0	$H_{5/4}^2$
v^{105}	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	1	0	$H_{5/4}^2$
v^{106}	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	1	0	0	$H_{5/4}^2$
v^{107}	1	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	1	0	$H_{5/4}^2$
v^{108}	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1	R_2R_3	R_4
	dim	1	1	1	1	2	2	3	3	3	3	2	4	6
Λ_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	-1
Λ_2	1	0	1	1	0	0	0	1	0	0	1	-1	0	-1
Λ_3	2	0	0	0	0	1	1	1	1	1	1	0	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 95.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1	R_2R_3	R_4	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	2	4	6	
v^1	0	0	0	0	0	0	0	0	1	1	0	0	1	$H_{3/4}^1$
v^2	0	0	0	0	0	0	0	0	1	1	1	1	0	$H_{4/4}^2$
v^3	0	0	0	0	0	0	0	1	0	1	0	0	1	$H_{3/4}^1$
v^4	0	0	0	0	0	0	0	1	0	1	1	1	0	$H_{4/4}^2$
v^5	0	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/4}^1$
v^6	0	0	0	0	0	0	1	0	1	0	1	1	0	$H_{4/4}^2$
v^7	0	0	0	0	0	0	1	1	0	0	0	0	1	$H_{3/4}^1$
v^8	0	0	0	0	0	0	1	1	0	0	1	1	0	$H_{4/4}^2$
v^9	0	0	0	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$

v^{11}	0	0	0	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{12}	0	0	0	1	0	1	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{13}	0	0	0	1	0	1	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{14}	0	0	0	1	0	1	1	0	0	0	0	0	1	$H_{4/4}^1$
v^{15}	0	0	0	1	0	1	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{16}	0	0	0	1	1	0	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{17}	0	0	0	1	1	0	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{18}	0	0	0	1	1	0	1	0	0	0	0	0	1	$H_{4/4}^1$
v^{19}	0	0	0	1	1	0	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{20}	0	0	0	2	0	0	0	0	0	2	1	0	1	$H_{6/6}^9$
v^{21}	0	0	0	2	0	0	0	0	0	2	2	1	0	$H_{7/6}^3$
v^{22}	0	0	0	2	0	0	1	0	0	1	1	0	1	$H_{6/6}^9$
v^{23}	0	0	0	2	0	0	1	0	0	1	2	1	0	$H_{7/6}^3$
v^{24}	0	0	0	2	0	0	2	0	0	0	1	0	1	$H_{6/6}^9$
v^{25}	0	0	0	2	0	0	2	0	0	0	2	1	0	$H_{7/6}^3$
v^{26}	0	0	1	0	0	1	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{27}	0	0	1	0	0	1	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{28}	0	0	1	0	0	1	0	1	0	0	0	0	1	$H_{4/4}^1$
v^{29}	0	0	1	0	0	1	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{30}	0	0	1	0	1	0	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{31}	0	0	1	0	1	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{32}	0	0	1	0	1	0	0	1	0	0	0	0	1	$H_{4/4}^1$
v^{33}	0	0	1	0	1	0	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{34}	0	0	1	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{35}	0	0	1	1	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{36}	0	0	1	1	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{37}	0	0	1	1	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{38}	0	0	2	0	0	0	0	0	2	0	1	0	1	$H_{6/6}^9$
v^{39}	0	0	2	0	0	0	0	0	2	0	2	1	0	$H_{7/6}^3$
v^{40}	0	0	2	0	0	0	0	1	1	0	1	0	1	$H_{6/6}^9$
v^{41}	0	0	2	0	0	0	0	1	1	0	2	1	0	$H_{7/6}^3$
v^{42}	0	0	2	0	0	0	0	2	0	0	1	0	1	$H_{6/6}^9$
v^{43}	0	0	2	0	0	0	0	2	0	0	2	1	0	$H_{7/6}^3$
v^{44}	0	1	0	0	0	1	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{45}	0	1	0	0	0	1	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{46}	0	1	0	0	0	1	0	1	0	0	0	0	1	$H_{4/4}^1$
v^{47}	0	1	0	0	0	1	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{48}	0	1	0	0	1	0	0	0	1	0	0	0	1	$H_{4/4}^1$
v^{49}	0	1	0	0	1	0	0	0	1	0	1	1	0	$H_{5/4}^3$
v^{50}	0	1	0	0	1	0	0	1	0	0	0	0	1	$H_{4/4}^1$
v^{51}	0	1	0	0	1	0	0	1	0	0	1	1	0	$H_{5/4}^3$
v^{52}	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

v^{53}	0	1	0	1	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{54}	0	1	0	1	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{55}	0	1	0	1	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{56}	0	1	1	0	0	0	0	0	2	0	1	0	1	$H_{6/6}^9$
v^{57}	0	1	1	0	0	0	0	0	2	0	2	1	0	$H_{7/6}^3$
v^{58}	0	1	1	0	0	0	0	1	1	0	1	0	1	$H_{6/6}^9$
v^{59}	0	1	1	0	0	0	0	1	1	0	2	1	0	$H_{7/6}^3$
v^{60}	0	1	1	0	0	0	0	2	0	0	1	0	1	$H_{6/6}^9$
v^{61}	0	1	1	0	0	0	0	2	0	0	2	1	0	$H_{7/6}^3$
v^{62}	0	2	0	0	0	0	0	0	2	0	1	0	1	$H_{6/6}^9$
v^{63}	0	2	0	0	0	0	0	0	2	0	2	1	0	$H_{7/6}^3$
v^{64}	0	2	0	0	0	0	0	1	1	0	1	0	1	$H_{6/6}^9$
v^{65}	0	2	0	0	0	0	0	1	1	0	2	1	0	$H_{7/6}^3$
v^{66}	0	2	0	0	0	0	0	2	0	0	1	0	1	$H_{6/6}^9$
v^{67}	0	2	0	0	0	0	0	2	0	0	2	1	0	$H_{7/6}^3$
v^{68}	1	0	0	0	0	1	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{69}	1	0	0	0	0	1	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{70}	1	0	0	0	0	1	1	0	0	0	0	0	1	$H_{4/4}^1$
v^{71}	1	0	0	0	0	1	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{72}	1	0	0	0	1	0	0	0	0	1	0	0	1	$H_{4/4}^1$
v^{73}	1	0	0	0	1	0	0	0	0	1	1	1	0	$H_{5/4}^3$
v^{74}	1	0	0	0	1	0	1	0	0	0	0	0	1	$H_{4/4}^1$
v^{75}	1	0	0	0	1	0	1	0	0	0	1	1	0	$H_{5/4}^3$
v^{76}	1	0	0	1	0	0	0	0	0	2	1	0	1	$H_{6/6}^9$
v^{77}	1	0	0	1	0	0	0	0	0	2	2	1	0	$H_{7/6}^3$
v^{78}	1	0	0	1	0	0	1	0	0	1	1	0	1	$H_{6/6}^9$
v^{79}	1	0	0	1	0	0	1	0	0	1	2	1	0	$H_{7/6}^3$
v^{80}	1	0	0	1	0	0	2	0	0	0	1	0	1	$H_{6/6}^9$
v^{81}	1	0	0	1	0	0	2	0	0	0	2	1	0	$H_{7/6}^3$
v^{82}	1	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{83}	1	0	1	0	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{84}	1	0	1	0	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{85}	1	0	1	0	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{86}	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{87}	1	1	0	0	0	2	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{88}	1	1	0	0	1	1	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{89}	1	1	0	0	2	0	0	0	0	0	0	0	1	$H_{5/4}^1$
v^{90}	2	0	0	0	0	0	0	0	0	2	1	0	1	$H_{6/6}^9$
v^{91}	2	0	0	0	0	0	0	0	0	2	2	1	0	$H_{7/6}^3$
v^{92}	2	0	0	0	0	0	1	0	0	1	1	0	1	$H_{6/6}^9$
v^{93}	2	0	0	0	0	0	1	0	0	1	2	1	0	$H_{7/6}^3$
v^{94}	2	0	0	0	0	0	2	0	0	0	1	0	1	$H_{6/6}^9$

v^{95}	2	0	0	0	0	0	2	0	0	0	2	1	0	$H_{7/6}^3$
----------	---	---	---	---	---	---	---	---	---	---	---	---	---	-------------

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	R_1	R_2R_3	R_4
dim		2	2	2	2	2	4	6
S_1	2	1	1	1	1	-1	-2	-3

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 34.

	X_1	X_2	X_3	X_4	R_1	R_2R_3	R_4	Bipartite graph
v^1	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	0	2	0	1	0	$H_{3/2}^1$
v^3	0	0	0	3	0	0	1	$H_{4/3}^1$
v^4	0	0	1	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	1	0	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	1	$H_{4/3}^1$
v^7	0	0	2	0	0	1	0	$H_{3/2}^1$
v^8	0	0	2	1	0	0	1	$H_{4/3}^1$
v^9	0	0	3	0	0	0	1	$H_{4/3}^1$
v^{10}	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	1	0	1	0	1	0	$H_{3/2}^1$
v^{12}	0	1	0	2	0	0	1	$H_{4/3}^1$
v^{13}	0	1	1	0	0	1	0	$H_{3/2}^1$
v^{14}	0	1	1	1	0	0	1	$H_{4/3}^1$
v^{15}	0	1	2	0	0	0	1	$H_{4/3}^1$
v^{16}	0	2	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	2	0	1	0	0	1	$H_{4/3}^1$
v^{18}	0	2	1	0	0	0	1	$H_{4/3}^1$
v^{19}	0	3	0	0	0	0	1	$H_{4/3}^1$
v^{20}	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{21}	1	0	0	1	0	1	0	$H_{3/2}^1$
v^{22}	1	0	0	2	0	0	1	$H_{4/3}^1$
v^{23}	1	0	1	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	1	1	0	0	1	$H_{4/3}^1$
v^{25}	1	0	2	0	0	0	1	$H_{4/3}^1$
v^{26}	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{27}	1	1	0	1	0	0	1	$H_{4/3}^1$
v^{28}	1	1	1	0	0	0	1	$H_{4/3}^1$

v^{29}	1	2	0	0	0	0	1	$H_{4/3}^1$
v^{30}	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{31}	2	0	0	1	0	0	1	$H_{4/3}^1$
v^{32}	2	0	1	0	0	0	1	$H_{4/3}^1$
v^{33}	2	1	0	0	0	0	1	$H_{4/3}^1$
v^{34}	3	0	0	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_1	X_2	X_3	X_4	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	
dim	2	2	2	2	1	1	1	1	1	1	1	1	2	2	
Z_1	1	1	0	0	1	0	-1	0	0	0	-1	0	0	-1	0
Z_2	1	1	1	0	0	0	0	-1	0	0	0	-1	-1	0	
Z_3	1	0	0	1	1	-1	0	0	0	-1	0	0	0	-1	
Z_4	1	0	1	1	0	0	0	-1	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 27.

	X_1	X_2	X_3	X_4	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	Bipartite graph
	2	2	2	2	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	1	0	0	0	0	1	1	0	0	0	0	$H_{3/2}^1$
v^2	0	0	0	1	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	1	1	1	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^6	0	0	1	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^7	0	0	1	0	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^8	0	0	1	0	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^9	0	0	1	0	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{13}	0	1	0	0	0	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{14}	0	1	0	1	0	0	0	0	0	0	0	0	1	1	$H_{4/4}^2$
v^{15}	0	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{5/4}^3$
v^{16}	0	1	0	1	0	0	0	0	1	0	1	0	1	0	$H_{5/4}^3$
v^{17}	0	1	0	1	0	0	0	1	0	1	0	0	0	1	$H_{5/4}^3$
v^{18}	0	1	0	1	0	0	1	0	1	0	0	0	1	0	$H_{5/4}^3$
v^{19}	0	1	0	1	0	1	0	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{20}	0	1	0	1	0	1	0	1	0	0	0	0	0	1	$H_{5/4}^3$

v^{21}	0	1	0	1	1	0	0	0	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{22}	0	1	0	1	1	0	1	0	0	0	0	0	0	1	0	$H_{5/4}^3$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^{24}	1	0	0	0	0	0	0	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{25}	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{26}	1	0	0	0	0	1	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	0	0	1	0	1	0	0	0	0	0	0	0	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	R_1	R_2R_3	R_4
	dim	1	1	1	1	1	1	1	1	2	2	2	4	6
T_1	1	1	0	0	1	0	0	0	0	0	0	0	0	-1
T_2	1	0	1	1	0	0	0	0	0	0	0	0	0	-1
T_3	1	0	0	0	0	1	0	0	1	0	0	0	0	-1
T_4	1	0	0	0	0	0	1	1	0	0	0	0	0	-1
T_5	2	0	0	0	0	0	0	0	0	1	1	-1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 37.

	M_1^+	M_1^-	M_2^+	M_2^-	M_3^+	M_3^-	M_4^+	M_4^-	M_5^+	M_5^-	R_1	R_2R_3	R_4	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	4	6	
v^1	0	0	0	0	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^2	0	0	0	0	0	0	0	0	0	2	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^6	0	0	1	1	0	0	1	1	0	1	0	0	1	$H_{6/5}^2$
v^7	0	0	1	1	0	0	1	1	1	0	0	0	1	$H_{6/5}^2$
v^8	0	0	1	1	0	1	0	1	0	1	0	0	1	$H_{6/5}^2$
v^9	0	0	1	1	0	1	0	1	1	0	0	0	1	$H_{6/5}^2$
v^{10}	0	0	1	1	1	0	1	0	0	1	0	0	1	$H_{6/5}^2$
v^{11}	0	0	1	1	1	0	1	0	1	0	0	0	1	$H_{6/5}^2$
v^{12}	0	0	1	1	1	1	0	0	0	1	0	0	1	$H_{6/5}^2$
v^{13}	0	0	1	1	1	1	0	0	1	0	0	0	1	$H_{6/5}^2$
v^{14}	0	1	0	1	0	0	1	1	0	1	0	0	1	$H_{6/5}^2$
v^{15}	0	1	0	1	0	0	1	1	1	0	0	0	1	$H_{6/5}^2$
v^{16}	0	1	0	1	0	1	0	1	0	1	0	0	1	$H_{6/5}^2$
v^{17}	0	1	0	1	0	1	0	1	1	0	0	0	1	$H_{6/5}^2$
v^{18}	0	1	0	1	1	0	1	0	0	1	0	0	1	$H_{6/5}^2$

v^{19}	0	1	0	1	1	0	1	0	1	0	0	0	1	$H_{6/5}^2$
v^{20}	0	1	0	1	1	1	0	0	0	1	0	0	1	$H_{6/5}^2$
v^{21}	0	1	0	1	1	1	0	0	1	0	0	0	1	$H_{6/5}^2$
v^{22}	1	0	1	0	0	0	1	1	0	1	0	0	1	$H_{6/5}^2$
v^{23}	1	0	1	0	0	0	1	1	1	0	0	0	1	$H_{6/5}^2$
v^{24}	1	0	1	0	0	1	0	1	0	1	0	0	1	$H_{6/5}^2$
v^{25}	1	0	1	0	0	1	0	1	1	0	0	0	1	$H_{6/5}^2$
v^{26}	1	0	1	0	1	0	1	0	0	1	0	0	1	$H_{6/5}^2$
v^{27}	1	0	1	0	1	0	1	0	1	0	0	0	1	$H_{6/5}^2$
v^{28}	1	0	1	0	1	1	0	0	0	1	0	0	1	$H_{6/5}^2$
v^{29}	1	0	1	0	1	1	0	0	1	0	0	0	1	$H_{6/5}^2$
v^{30}	1	1	0	0	0	0	1	1	0	1	0	0	1	$H_{6/5}^2$
v^{31}	1	1	0	0	0	0	1	1	1	0	0	0	1	$H_{6/5}^2$
v^{32}	1	1	0	0	0	1	0	1	0	1	0	0	1	$H_{6/5}^2$
v^{33}	1	1	0	0	0	1	0	1	1	0	0	0	1	$H_{6/5}^2$
v^{34}	1	1	0	0	1	0	1	0	0	1	0	0	1	$H_{6/5}^2$
v^{35}	1	1	0	0	1	0	1	0	1	0	0	0	1	$H_{6/5}^2$
v^{36}	1	1	0	0	1	1	0	0	0	1	0	0	1	$H_{6/5}^2$
v^{37}	1	1	0	0	1	1	0	0	1	0	0	0	1	$H_{6/5}^2$

SG 224

Space group $Pn\bar{3}m$, No. 224, without SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4
	dim	1	1	1	1	2	2	3	3	3	3	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	1	0	0	0	0	0	-1
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	-1	0	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	-1	0	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1
Δ_5	2	0	0	0	0	0	0	1	1	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 33.

Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4	
1	1	1	1	2	2	3	3	3	3	2	2	2	2	Bipartite graph

v^1	0	0	0	0	0	0	0	1	1	0	0	0	2	1	$H_{5/4}^3$
v^2	0	0	0	0	0	0	0	1	1	0	1	0	1	1	$H_{5/4}^3$
v^3	0	0	0	0	0	0	0	1	1	0	2	0	0	1	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	0	0	1	0	1	2	0	$H_{5/4}^3$
v^5	0	0	0	0	0	0	1	0	0	1	1	1	1	0	$H_{5/4}^3$
v^6	0	0	0	0	0	0	1	0	0	1	2	1	0	0	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	0	1	2	1	$H_{7/6}^3$
v^8	0	0	0	0	0	1	0	1	0	1	1	1	1	1	$H_{7/6}^3$
v^9	0	0	0	0	0	1	0	1	0	1	2	1	0	1	$H_{7/6}^3$
v^{10}	0	0	0	0	1	0	1	0	1	0	0	1	2	1	$H_{7/6}^3$
v^{11}	0	0	0	0	1	0	1	0	1	0	1	1	1	1	$H_{7/6}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	2	1	0	1	$H_{7/6}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	0	1	0	1	$H_{4/4}^2$
v^{14}	0	0	0	1	0	0	0	1	0	0	0	0	1	1	$H_{4/3}^2$
v^{15}	0	0	0	1	0	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^{16}	0	0	0	1	1	0	1	0	0	0	0	1	1	1	$H_{6/5}^1$
v^{17}	0	0	0	1	1	0	1	0	0	0	1	1	0	1	$H_{6/5}^1$
v^{18}	0	0	1	0	0	0	1	0	0	0	0	1	1	0	$H_{4/3}^2$
v^{19}	0	0	1	0	0	0	1	0	0	0	1	1	0	0	$H_{4/3}^2$
v^{20}	0	0	1	0	0	1	0	1	0	0	0	1	1	1	$H_{6/5}^1$
v^{21}	0	0	1	0	0	1	0	1	0	0	1	1	0	1	$H_{6/5}^1$
v^{22}	0	1	0	0	0	0	0	0	0	1	0	1	1	0	$H_{4/3}^2$
v^{23}	0	1	0	0	0	0	0	0	0	1	1	1	0	0	$H_{4/3}^2$
v^{24}	0	1	0	0	1	0	0	0	1	0	0	1	1	1	$H_{6/5}^1$
v^{25}	0	1	0	0	1	0	0	0	1	0	1	1	0	1	$H_{6/5}^1$
v^{26}	0	1	0	1	1	0	0	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{27}	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{28}	1	0	0	0	0	0	0	0	1	0	0	0	1	1	$H_{4/3}^2$
v^{29}	1	0	0	0	0	0	0	0	1	0	1	0	0	1	$H_{4/3}^2$
v^{30}	1	0	0	0	0	1	0	0	0	1	0	1	1	1	$H_{6/5}^1$
v^{31}	1	0	0	0	0	1	0	0	0	1	1	1	0	1	$H_{6/5}^1$
v^{32}	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{33}	1	0	1	0	0	1	0	0	0	0	0	1	0	1	$H_{5/4}^3$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4
dim	1	1	1	1	2	2	3	3	3	3	2	2	2	2

Σ_1	1	1	0	0	0	1	0	0	1	1	1	0	0	-1	-1
Σ_2	1	0	0	1	0	1	0	1	1	0	1	-1	-1	0	0
Σ_3	1	0	1	0	0	0	1	1	0	1	1	0	-1	-1	0
Σ_4	1	0	0	0	1	0	1	1	1	1	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 64.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	M_1	M_2	M_3	M_4	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	1	1	1	$H_{5/6}^4, H_{5/6}^2$
v^2	0	0	0	0	0	0	0	0	1	1	1	0	2	0	$H_{5/6}^2$
v^3	0	0	0	0	0	0	0	1	1	0	0	1	0	2	$H_{5/6}^2$
v^4	0	0	0	0	0	0	0	1	1	0	1	0	1	1	$H_{5/6}^2, H_{5/6}^4$
v^5	0	0	0	0	0	0	1	0	0	1	0	2	0	1	$H_{5/6}^2$
v^6	0	0	0	0	0	0	1	0	0	1	1	1	1	0	$H_{5/6}^2, H_{5/6}^4$
v^7	0	0	0	0	0	0	1	1	0	0	1	1	0	1	$H_{5/6}^2, H_{5/6}^4$
v^8	0	0	0	0	0	0	1	1	0	0	2	0	1	0	$H_{5/6}^2$
v^9	0	0	0	0	0	1	0	0	0	2	0	2	1	1	$H_{7/8}^{12}, H_{7/8}^{10}, H_{7/8}^3$
v^{10}	0	0	0	0	0	1	0	0	0	2	1	1	2	0	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{11}	0	0	0	0	0	1	0	1	0	1	0	2	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{12}	0	0	0	0	0	1	0	1	0	1	1	1	1	1	$H_{7/8}^{12}, H_{7/8}^3, H_{7/8}^{10}$
v^{13}	0	0	0	0	0	1	0	1	0	1	2	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{14}	0	0	0	0	0	1	0	2	0	0	1	1	0	2	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{15}	0	0	0	0	0	1	0	2	0	0	2	0	1	1	$H_{7/8}^{10}, H_{7/8}^{12}, H_{7/8}^3$
v^{16}	0	0	0	0	1	0	0	0	2	0	0	1	1	2	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{17}	0	0	0	0	1	0	0	0	2	0	1	0	2	1	$H_{7/8}^{10}, H_{7/8}^3, H_{7/8}^{12}$
v^{18}	0	0	0	0	1	0	1	0	1	0	0	2	0	2	$H_{7/8}^{10}, H_{7/8}^3$
v^{19}	0	0	0	0	1	0	1	0	1	0	1	1	1	1	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{20}	0	0	0	0	1	0	1	0	1	0	2	0	2	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{21}	0	0	0	0	1	0	2	0	0	0	1	2	0	1	$H_{7/8}^3, H_{7/8}^{12}, H_{7/8}^{10}$
v^{22}	0	0	0	0	1	0	2	0	0	0	2	1	1	0	$H_{7/8}^{12}, H_{7/8}^{10}, H_{7/8}^3$
v^{23}	0	0	0	0	1	1	0	0	0	0	0	1	0	1	$H_{4/4}^2$
v^{24}	0	0	0	0	1	1	0	0	0	0	1	0	1	0	$H_{4/4}^2$
v^{25}	0	0	0	1	0	0	0	0	0	1	0	1	0	1	$H_{4/4}^4$
v^{26}	0	0	0	1	0	0	0	0	0	1	1	0	1	0	$H_{4/4}^4$
v^{27}	0	0	0	1	0	0	0	1	0	0	1	0	0	1	$H_{4/4}^4$
v^{28}	0	0	0	1	1	0	0	0	1	0	0	1	0	2	$H_{6/6}^8, H_{6/6}^7$
v^{29}	0	0	0	1	1	0	0	0	1	0	1	0	1	1	$H_{6/6}^8, H_{6/6}^7$
v^{30}	0	0	0	1	1	0	1	0	0	0	1	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{31}	0	0	0	1	1	0	1	0	0	0	2	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{32}	0	0	0	2	1	0	0	0	0	0	1	0	0	1	$H_{5/4}^3$
v^{33}	0	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{4/4}^4$
v^{34}	0	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{4/4}^4$

v^{35}	0	0	1	0	0	0	1	0	0	0	1	1	0	0	$H_{4/4}^A$
v^{36}	0	0	1	0	0	1	0	0	0	1	0	2	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{37}	0	0	1	0	0	1	0	0	0	1	1	1	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{38}	0	0	1	0	0	1	0	1	0	0	1	1	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{39}	0	0	1	0	0	1	0	1	0	0	2	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{40}	0	0	1	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{41}	0	0	2	0	0	1	0	0	0	0	1	1	0	0	$H_{5/4}^3$
v^{42}	0	1	0	0	0	0	0	0	0	1	0	1	1	0	$H_{4/4}^A$
v^{43}	0	1	0	0	0	0	0	1	0	0	0	1	0	1	$H_{4/4}^A$
v^{44}	0	1	0	0	0	0	0	1	0	0	1	0	1	0	$H_{4/4}^A$
v^{45}	0	1	0	0	1	0	0	0	1	0	0	1	1	1	$H_{6/6}^8, H_{6/6}^7$
v^{46}	0	1	0	0	1	0	0	0	1	0	1	0	2	0	$H_{6/6}^8, H_{6/6}^7$
v^{47}	0	1	0	0	1	0	1	0	0	0	0	2	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{48}	0	1	0	0	1	0	1	0	0	0	1	1	1	0	$H_{6/6}^7, H_{6/6}^8$
v^{49}	0	1	0	1	1	0	0	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{50}	0	1	0	1	1	0	0	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{51}	0	1	1	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{52}	0	2	0	0	1	0	0	0	0	0	0	1	1	0	$H_{5/4}^3$
v^{53}	1	0	0	0	0	0	0	0	1	0	0	0	1	1	$H_{4/4}^A$
v^{54}	1	0	0	0	0	0	1	0	0	0	0	1	0	1	$H_{4/4}^A$
v^{55}	1	0	0	0	0	0	1	0	0	0	1	0	1	0	$H_{4/4}^A$
v^{56}	1	0	0	0	0	1	0	0	0	1	0	1	1	1	$H_{6/6}^7, H_{6/6}^8$
v^{57}	1	0	0	0	0	1	0	0	0	1	1	0	2	0	$H_{6/6}^8, H_{6/6}^7$
v^{58}	1	0	0	0	0	1	0	1	0	0	0	1	0	2	$H_{6/6}^8, H_{6/6}^7$
v^{59}	1	0	0	0	0	1	0	1	0	0	1	0	1	1	$H_{6/6}^8, H_{6/6}^7$
v^{60}	1	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{61}	1	0	1	0	0	1	0	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{62}	1	0	1	0	0	1	0	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{63}	1	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{64}	2	0	0	0	0	1	0	0	0	0	0	0	1	1	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-
dim		1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	3	3	3	3
Λ_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	0	-1	0	0	0	-1	-1	0
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1	-1	0	0	0	-1	0	0	-1
Λ_3	2	0	0	0	0	1	1	1	1	1	1	0	0	0	0	-1	-1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 84.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-	Bipartite graph	
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	3	3	3	3		
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^{14}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/2}^1$
v^{15}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{19}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^{20}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/2}^1$
v^{21}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{22}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{23}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{24}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{25}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{26}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{28}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{30}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{32}	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{33}	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{34}	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{35}	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{36}	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{37}	0	0	0	1	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{38}	0	0	0	1	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{39}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{40}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

v^{41}	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{42}	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{43}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{44}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{45}	0	0	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{46}	0	0	1	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	$H_{4/3}^2$
v^{47}	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{48}	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{49}	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{50}	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	$H_{4/3}^2$
v^{51}	0	0	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{52}	0	0	1	0	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{53}	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{54}	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{55}	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{56}	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{57}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{58}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{59}	0	1	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{60}	0	1	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	$H_{4/3}^2$
v^{61}	0	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{62}	0	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{63}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	$H_{4/3}^2$
v^{64}	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	$H_{4/3}^2$
v^{65}	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	$H_{4/3}^2$
v^{66}	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{4/3}^2$
v^{67}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{68}	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{69}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{70}	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{71}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{72}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{73}	1	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	1	$H_{4/3}^2$
v^{74}	1	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	$H_{4/3}^2$
v^{75}	1	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{76}	1	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{77}	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{4/3}^2$
v^{78}	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	1	0	$H_{4/3}^2$
v^{79}	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{80}	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{81}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{82}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

v^{83}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$	
v^{84}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_1	X_2	X_3	X_4	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-
	dim	2	2	2	2	1	1	1	1	2	2	3	3	3	3
S_1	1	1	0	0	1	0	0	0	-1	0	-1	-1	-1	-1	0
S_2	1	0	0	1	1	-1	0	0	0	-1	0	0	-1	-1	-1
S_3	1	1	1	0	0	0	0	-1	0	-1	0	-1	-1	0	-1
S_4	1	0	1	1	0	0	-1	0	0	0	-1	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 64.

	X_1	X_2	X_3	X_4	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-	Bipartite graph
	2	2	2	2	1	1	1	1	2	2	3	3	3	3	
v^1	0	0	0	1	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^2	0	0	1	0	1	1	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^3	0	0	1	1	1	0	0	0	0	0	0	0	1	0	$H_{4/4}^4$
v^4	0	0	1	1	2	0	0	0	0	1	0	0	0	0	$H_{5/4}^3$
v^5	0	1	0	0	0	1	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	1	0	0	0	0	$H_{4/4}^2$
v^7	0	1	0	1	0	0	0	1	0	0	0	0	0	1	$H_{4/4}^4$
v^8	0	1	0	1	0	0	1	0	0	0	0	0	1	0	$H_{4/4}^4$
v^9	0	1	0	1	0	1	0	0	0	0	0	1	0	0	$H_{4/4}^4$
v^{10}	0	1	0	1	0	1	0	1	1	0	0	0	0	0	$H_{5/4}^3$
v^{11}	0	1	0	1	1	0	0	0	0	0	1	0	0	0	$H_{4/4}^4$
v^{12}	0	1	0	1	1	0	1	0	0	1	0	0	0	0	$H_{5/4}^3$
v^{13}	0	1	0	2	0	0	0	0	0	0	0	1	1	0	$H_{5/6}^2$
v^{14}	0	1	0	2	0	0	0	1	1	0	0	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{15}	0	1	0	2	1	0	0	0	0	1	0	1	0	0	$H_{6/6}^8, H_{6/6}^7$
v^{16}	0	1	1	0	0	1	0	0	0	0	0	0	0	1	$H_{4/4}^4$
v^{17}	0	1	1	0	0	2	0	0	1	0	0	0	0	0	$H_{5/4}^3$
v^{18}	0	1	1	1	0	0	0	0	0	0	0	0	1	1	$H_{5/6}^4, H_{5/6}^2$
v^{19}	0	1	1	1	0	1	0	0	1	0	0	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{20}	0	1	1	1	1	0	0	0	0	1	0	0	0	1	$H_{6/6}^7, H_{6/6}^8$
v^{21}	0	1	1	2	0	0	0	0	1	0	0	0	2	0	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{22}	0	2	0	1	0	0	0	0	0	0	1	0	0	1	$H_{5/6}^2$
v^{23}	0	2	0	1	0	0	1	0	0	1	0	0	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{24}	0	2	0	1	0	1	0	0	1	0	1	0	0	0	$H_{6/6}^8, H_{6/6}^7$

v^{25}	0	2	0	2	0	0	0	0	0	1	0	1	0	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{26}	0	2	0	2	0	0	0	0	1	0	1	0	1	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{27}	0	2	1	1	0	0	0	0	0	1	0	0	0	2	$H_{7/8}^{12}, H_{7/8}^{10}, H_{7/8}^3$
v^{28}	1	0	0	0	0	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{29}	1	0	0	1	0	0	0	1	0	0	0	1	0	0	$H_{4/4}^4$
v^{30}	1	0	0	1	0	0	0	2	1	0	0	0	0	0	$H_{5/4}^3$
v^{31}	1	0	1	0	0	0	0	0	1	1	0	0	0	0	$H_{4/4}^2$
v^{32}	1	0	1	0	0	0	0	1	0	0	0	0	0	1	$H_{4/4}^4$
v^{33}	1	0	1	0	0	0	1	0	0	0	0	0	1	0	$H_{4/4}^4$
v^{34}	1	0	1	0	0	1	0	0	0	0	0	1	0	0	$H_{4/4}^4$
v^{35}	1	0	1	0	0	1	0	1	1	0	0	0	0	0	$H_{5/4}^3$
v^{36}	1	0	1	0	1	0	0	0	0	0	1	0	0	0	$H_{4/4}^4$
v^{37}	1	0	1	0	1	0	1	0	0	1	0	0	0	0	$H_{5/4}^3$
v^{38}	1	0	1	1	0	0	0	0	0	0	0	1	1	0	$H_{5/6}^2, H_{5/6}^4$
v^{39}	1	0	1	1	0	0	0	1	1	0	0	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{40}	1	0	1	1	1	0	0	0	0	1	0	1	0	0	$H_{6/6}^8, H_{6/6}^7$
v^{41}	1	0	2	0	0	0	0	0	0	0	0	0	1	1	$H_{5/6}^2$
v^{42}	1	0	2	0	0	1	0	0	1	0	0	0	1	0	$H_{6/6}^8, H_{6/6}^7$
v^{43}	1	0	2	0	1	0	0	0	0	1	0	0	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{44}	1	0	2	1	0	0	0	0	1	0	0	0	2	0	$H_{7/8}^{10}, H_{7/8}^3, H_{7/8}^{12}$
v^{45}	1	1	0	0	0	0	1	0	0	0	1	0	0	0	$H_{4/4}^4$
v^{46}	1	1	0	0	0	0	2	0	0	1	0	0	0	0	$H_{5/4}^3$
v^{47}	1	1	0	1	0	0	0	0	0	0	1	1	0	0	$H_{5/6}^2, H_{5/6}^4$
v^{48}	1	1	0	1	0	0	0	1	1	0	1	0	0	0	$H_{6/6}^8, H_{6/6}^7$
v^{49}	1	1	0	1	0	0	1	0	0	1	0	1	0	0	$H_{6/6}^8, H_{6/6}^7$
v^{50}	1	1	0	2	0	0	0	0	0	1	0	2	0	0	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{51}	1	1	1	0	0	0	0	0	0	0	1	0	0	1	$H_{5/6}^2, H_{5/6}^4$
v^{52}	1	1	1	0	0	0	1	0	0	1	0	0	0	1	$H_{6/6}^8, H_{6/6}^7$
v^{53}	1	1	1	0	0	1	0	0	1	0	1	0	0	0	$H_{6/6}^7, H_{6/6}^8$
v^{54}	1	1	1	1	0	0	0	0	0	1	0	1	0	1	$H_{7/8}^{12}, H_{7/8}^3, H_{7/8}^{10}$
v^{55}	1	1	1	1	0	0	0	0	1	0	1	0	1	0	$H_{7/8}^3, H_{7/8}^{12}, H_{7/8}^{10}$
v^{56}	1	1	2	0	0	0	0	0	0	1	0	0	0	2	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{57}	1	2	0	1	0	0	0	0	1	0	2	0	0	0	$H_{7/8}^3, H_{7/8}^{10}, H_{7/8}^{12}$
v^{58}	2	0	1	0	0	0	0	0	0	0	1	1	0	0	$H_{5/6}^2$
v^{59}	2	0	1	0	0	0	0	1	1	0	1	0	0	0	$H_{6/6}^8, H_{6/6}^7$
v^{60}	2	0	1	0	0	0	1	0	0	1	0	1	0	0	$H_{6/6}^8, H_{6/6}^7$
v^{61}	2	0	1	1	0	0	0	0	0	1	0	2	0	0	$H_{7/8}^{10}, H_{7/8}^3, H_{7/8}^{12}$
v^{62}	2	0	2	0	0	0	0	0	0	1	0	1	0	1	$H_{7/8}^{10}, H_{7/8}^3$
v^{63}	2	0	2	0	0	0	0	0	1	0	1	0	1	0	$H_{7/8}^{10}, H_{7/8}^3$
v^{64}	2	1	1	0	0	0	0	0	1	0	2	0	0	0	$H_{7/8}^{12}, H_{7/8}^{10}, H_{7/8}^3$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_1	X_2	X_3	X_4	M_1	M_2	M_3	M_4
	dim	2	2	2	2	2	2	2	2
Z_1	2	1	1	1	1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	X_1	X_2	X_3	X_4	M_1	M_2	M_3	M_4	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_1	M_2	M_3	M_4	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-
	dim	2	2	2	2	1	1	1	1	2	2	3	3	3	3
T_1	1	0	0	0	1	0	0	0	-1	0	-1	0	0	-1	0
T_2	1	0	0	0	1	-1	0	0	0	-1	0	0	-1	0	0
T_3	1	0	1	0	0	0	0	-1	0	-1	0	0	0	0	-1
T_4	1	0	1	0	0	0	-1	0	0	0	-1	-1	0	0	0
T_5	2	1	0	1	0	0	0	0	0	0	0	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 33.

	M_1	M_2	M_3	M_4	R_1^+	R_1^-	R_2^+	R_2^-	R_3^+	R_3^-	R_4^+	R_4^-	R_5^+	R_5^-	Bipartite graph
	2	2	2	2	1	1	1	1	2	2	3	3	3	3	
v^1	0	0	0	1	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^2	0	0	1	1	0	0	0	1	0	0	0	1	0	0	$H_{4/3}^2$
v^3	0	0	1	1	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^4	0	0	2	1	0	0	0	0	0	0	0	1	1	0	$H_{5/4}^3$
v^5	0	1	0	0	0	1	1	0	0	0	0	0	0	0	$H_{3/2}^1$
v^6	0	1	0	1	0	0	0	0	1	1	0	0	0	0	$H_{4/4}^2$
v^7	0	1	0	1	0	1	0	1	1	0	0	0	0	0	$H_{5/4}^3$
v^8	0	1	0	1	1	0	1	0	0	1	0	0	0	0	$H_{5/4}^3$
v^9	0	1	1	0	0	0	1	0	0	0	1	0	0	0	$H_{4/3}^2$
v^{10}	0	1	1	0	0	1	0	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{11}	0	1	1	1	0	0	0	1	1	0	1	0	0	0	$H_{6/5}^1$
v^{12}	0	1	1	1	0	0	1	0	0	1	0	1	0	0	$H_{6/5}^1$
v^{13}	0	1	1	1	0	1	0	0	1	0	0	0	1	0	$H_{6/5}^1$
v^{14}	0	1	1	1	1	0	0	0	0	1	0	0	0	1	$H_{6/5}^1$
v^{15}	0	1	2	0	0	0	0	0	0	0	1	0	0	1	$H_{5/4}^3$
v^{16}	0	1	2	1	0	0	0	0	0	1	0	1	0	1	$H_{7/6}^3$
v^{17}	0	1	2	1	0	0	0	0	1	0	1	0	1	0	$H_{7/6}^3$
v^{18}	1	0	0	1	0	0	0	1	0	0	0	1	0	0	$H_{4/3}^2$
v^{19}	1	0	0	1	1	0	0	0	0	0	0	0	1	0	$H_{4/3}^2$
v^{20}	1	0	1	1	0	0	0	0	0	0	0	1	1	0	$H_{5/4}^3$
v^{21}	1	1	0	0	0	0	1	0	0	0	1	0	0	0	$H_{4/3}^2$
v^{22}	1	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{4/3}^2$
v^{23}	1	1	0	1	0	0	0	1	1	0	1	0	0	0	$H_{6/5}^1$
v^{24}	1	1	0	1	0	0	1	0	0	1	0	1	0	0	$H_{6/5}^1$
v^{25}	1	1	0	1	0	1	0	0	1	0	0	0	1	0	$H_{6/5}^1$
v^{26}	1	1	0	1	1	0	0	0	0	1	0	0	0	1	$H_{6/5}^1$
v^{27}	1	1	1	0	0	0	0	0	0	0	1	0	0	1	$H_{5/4}^3$
v^{28}	1	1	1	1	0	0	0	0	0	1	0	1	0	1	$H_{7/6}^3$
v^{29}	1	1	1	1	0	0	0	0	1	0	1	0	1	0	$H_{7/6}^3$
v^{30}	2	0	0	1	0	0	0	0	0	0	0	1	1	0	$H_{5/4}^3$
v^{31}	2	1	0	0	0	0	0	0	0	0	1	0	0	1	$H_{5/4}^3$
v^{32}	2	1	0	1	0	0	0	0	0	1	0	1	0	1	$H_{7/6}^3$
v^{33}	2	1	0	1	0	0	0	0	1	0	1	0	1	0	$H_{7/6}^3$

SG 225

Space group $Fm\bar{3}m$, No. 225, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-
	dim	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2
Δ_1	1	1	0	0	0	1	0	0	1	0	0	-1	0	0	-1	0	0	0	0	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	-1	-1	0	0	0	0	0	0	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	0	0	0	-1	0	0	-1	0	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	0	-1	-1	0	0	0	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	0	0	0	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{19}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{20}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{21}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{22}	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{23}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{24}	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{25}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{26}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

v^{28}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-
dim		1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2
Λ_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	0	-1	0	0
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1	-1	0	0	0
Λ_3	2	0	0	0	0	1	1	1	1	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 28.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{18}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{19}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{21}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

v^{23}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{24}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{25}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{26}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{28}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	W_1	W_2	W_3	W_4	W_5
	dim	1	1	1	1	1	1	1	1	2	2	1	1	1	1	2
Z_1	1	1	0	0	0	1	0	0	0	0	1	-1	0	0	-1	0
Z_2	1	0	0	0	1	0	0	0	1	1	0	0	0	0	0	-1
Z_3	1	0	1	0	0	0	1	0	0	1	0	0	-1	-1	0	0
Z_4	1	0	0	1	0	0	0	1	0	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{15}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	W_1	W_2	W_3	W_4	W_5	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	1	1	1	1	2	
v^1	0	0	0	0	0	0	0	0	1	1	0	0	1	1	1	$H_{5/4}^3$
v^2	0	0	0	0	0	0	0	0	1	1	0	1	0	1	1	$H_{5/4}^3$
v^3	0	0	0	0	0	0	0	0	1	1	1	0	1	0	1	$H_{5/4}^3$
v^4	0	0	0	0	0	0	0	0	1	1	1	1	0	0	1	$H_{5/4}^3$
v^5	0	0	0	0	0	0	0	1	0	1	0	0	0	1	1	$H_{4/3}^2$
v^6	0	0	0	0	0	0	0	1	0	1	1	0	0	0	1	$H_{4/3}^2$
v^7	0	0	0	0	0	0	1	0	1	0	0	0	1	0	1	$H_{4/3}^2$
v^8	0	0	0	0	0	0	1	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^9	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	0	1	0	0	0	1	1	$H_{4/3}^2$
v^{15}	0	0	0	1	0	0	0	0	0	1	1	0	0	0	1	$H_{4/3}^2$
v^{16}	0	0	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	0	0	1	0	0	0	1	0	1	$H_{4/3}^2$
v^{18}	0	0	1	0	0	0	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^{19}	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	0	1	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

v^{21}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	W_1	W_2	W_3	W_4	W_5
	dim	1	1	1	1	2	2	1	1	1	1	2
Q_1	1	1	1	0	0	1	1	-1	0	-1	0	-1
Q_2	1	0	0	1	1	1	1	0	-1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{11}$ and the corresponding bipartite graph, number of Hilbert basis: 22.

	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	W_1	W_2	W_3	W_4	W_5	Bipartite graph
	1	1	1	1	2	2	1	1	1	1	2	
v^1	0	0	0	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	1	1	0	0	0	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^7	0	0	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^8	0	0	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{14}	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{21}	1	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$

SG 226

Space group $Fm\bar{3}c$, No. 226, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	
dim		1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
Δ_1	1	1	0	0	0	1	0	0	1	0	0	0	-1	-1	0	0	0	0	0	0	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	-1	0	0	-1	0	0	0	0	0	0	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	0	0	0	0	-1	-1	0	0	0	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	0	-1	0	0	-1	0	0	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	0	0	0	0	0	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	2	2	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{18}	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{19}	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{21}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{22}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$
v^{23}	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	$H_{3/2}^1$

v^{24}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{3/2}^1$
v^{25}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{26}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{31}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{32}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1L_2	L_3
	dim	1	1	1	1	2	2	3	3	3	3	4	2
Λ_1	1	1	0	0	1	0	0	0	1	1	0	0	-1
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1
Λ_3	2	0	0	0	0	1	1	1	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 45.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1L_2	L_3	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	4	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	1	0	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	0	0	1	0	1	0	1	1	$H_{4/4}^2$
v^4	0	0	0	0	0	0	1	1	0	0	1	1	$H_{4/4}^2$
v^5	0	0	0	0	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^9	0	0	0	1	0	1	1	0	0	0	1	1	$H_{5/4}^3$
v^{10}	0	0	0	1	1	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{11}	0	0	0	1	1	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{12}	0	0	0	2	0	0	0	0	0	2	1	2	$H_{7/6}^3$
v^{13}	0	0	0	2	0	0	1	0	0	1	1	2	$H_{7/6}^3$
v^{14}	0	0	0	2	0	0	2	0	0	0	1	2	$H_{7/6}^3$
v^{15}	0	0	1	0	0	1	0	0	1	0	1	1	$H_{5/4}^3$
v^{16}	0	0	1	0	0	1	0	1	0	0	1	1	$H_{5/4}^3$
v^{17}	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{18}	0	0	1	0	1	0	0	1	0	0	1	1	$H_{5/4}^3$

v^{19}	0	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	0	2	0	0	0	0	0	2	0	1	2	$H_{7/6}^3$	
v^{21}	0	0	2	0	0	0	0	1	1	0	1	2	$H_{7/6}^3$	
v^{22}	0	0	2	0	0	0	0	2	0	0	1	2	$H_{7/6}^3$	
v^{23}	0	1	0	0	0	1	0	0	1	0	1	1	$H_{5/4}^3$	
v^{24}	0	1	0	0	0	1	0	1	0	0	1	1	$H_{5/4}^3$	
v^{25}	0	1	0	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$	
v^{26}	0	1	0	0	1	0	0	1	0	0	1	1	$H_{5/4}^3$	
v^{27}	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$	
v^{28}	0	1	1	0	0	0	0	0	2	0	1	2	$H_{7/6}^3$	
v^{29}	0	1	1	0	0	0	0	1	1	0	1	2	$H_{7/6}^3$	
v^{30}	0	1	1	0	0	0	0	2	0	0	1	2	$H_{7/6}^3$	
v^{31}	0	2	0	0	0	0	0	0	2	0	1	2	$H_{7/6}^3$	
v^{32}	0	2	0	0	0	0	0	1	1	0	1	2	$H_{7/6}^3$	
v^{33}	0	2	0	0	0	0	0	2	0	0	1	2	$H_{7/6}^3$	
v^{34}	1	0	0	0	0	1	0	0	0	1	1	1	$H_{5/4}^3$	
v^{35}	1	0	0	0	0	1	1	0	0	0	1	1	$H_{5/4}^3$	
v^{36}	1	0	0	0	1	0	0	0	0	1	1	1	$H_{5/4}^3$	
v^{37}	1	0	0	0	1	0	1	0	0	0	1	1	$H_{5/4}^3$	
v^{38}	1	0	0	1	0	0	0	0	0	2	1	2	$H_{7/6}^3$	
v^{39}	1	0	0	1	0	0	1	0	0	1	1	2	$H_{7/6}^3$	
v^{40}	1	0	0	1	0	0	2	0	0	0	1	2	$H_{7/6}^3$	
v^{41}	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$	
v^{42}	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$	
v^{43}	2	0	0	0	0	0	0	0	0	2	1	2	$H_{7/6}^3$	
v^{44}	2	0	0	0	0	0	1	0	0	1	1	2	$H_{7/6}^3$	
v^{45}	2	0	0	0	0	0	2	0	0	0	1	2	$H_{7/6}^3$	

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	W_1W_4	W_2W_3	W_5
dim	1	1	1	1	1	1	1	1	2	2	2	2	2
Z_1	1	0	1	0	0	0	1	0	0	1	0	0	-1
Z_2	1	0	0	1	0	0	0	1	0	0	1	-2	0
Z_3	1	1	0	0	0	1	0	0	0	0	1	0	-1
Z_4	1	0	0	0	1	0	0	0	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 37.

	X_1^+	X_1^-	X_2^+	X_2^-	X_3^+	X_3^-	X_4^+	X_4^-	X_5^+	X_5^-	W_1W_4	W_2W_3	W_5	Bipartite graph
	1	1	1	1	1	1	1	1	2	2	2	2	2	
v^1	0	0	0	0	0	0	0	0	2	2	1	1	2	$H_{8/8}^1$
v^2	0	0	0	0	0	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	1	1	1	1	1	1	1	$H_{7/6}^3$
v^4	0	0	0	0	0	0	2	0	0	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	1	2	1	1	2	$H_{9/8}^1$
v^6	0	0	0	0	0	1	1	0	0	1	1	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	2	0	0	0	2	1	0	2	$H_{7/6}^3$
v^8	0	0	0	0	1	0	0	1	1	0	0	1	1	$H_{5/4}^3$
v^9	0	0	0	0	1	0	1	0	2	1	1	1	2	$H_{9/8}^1$
v^{10}	0	0	0	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	0	0	0	2	0	0	0	2	0	0	1	2	$H_{7/6}^3$
v^{12}	0	0	0	1	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	1	0	1	1	1	1	1	$H_{7/6}^3$
v^{14}	0	0	0	1	0	1	0	0	1	2	1	1	2	$H_{9/8}^1$
v^{15}	0	0	0	1	1	0	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{16}	0	0	0	2	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	0	0	1	0	0	0	0	1	1	1	1	1	1	$H_{7/6}^3$
v^{18}	0	0	1	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{19}	0	0	1	0	0	1	0	0	0	1	1	0	1	$H_{5/4}^3$
v^{20}	0	0	1	0	1	0	0	0	2	1	1	1	2	$H_{9/8}^1$
v^{21}	0	0	1	1	0	0	0	0	1	1	1	1	1	$H_{7/6}^3$
v^{22}	0	0	2	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{23}	0	1	0	0	0	0	0	1	1	2	1	1	2	$H_{9/8}^1$
v^{24}	0	1	0	0	0	0	1	0	0	1	1	0	1	$H_{5/4}^3$
v^{25}	0	1	0	0	0	1	0	0	0	2	1	0	2	$H_{7/6}^3$
v^{26}	0	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{27}	0	1	0	1	0	0	0	0	1	2	1	1	2	$H_{9/8}^1$
v^{28}	0	1	1	0	0	0	0	0	0	1	1	0	1	$H_{5/4}^3$
v^{29}	0	2	0	0	0	0	0	0	0	2	1	0	2	$H_{7/6}^3$
v^{30}	1	0	0	0	0	0	0	1	1	0	0	1	1	$H_{5/4}^3$
v^{31}	1	0	0	0	0	0	1	0	2	1	1	1	2	$H_{9/8}^1$
v^{32}	1	0	0	0	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{33}	1	0	0	0	1	0	0	0	2	0	0	1	2	$H_{7/6}^3$
v^{34}	1	0	0	1	0	0	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{35}	1	0	1	0	0	0	0	0	2	1	1	1	2	$H_{9/8}^1$
v^{36}	1	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{37}	2	0	0	0	0	0	0	0	2	0	0	1	2	$H_{7/6}^3$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep	$L_1 L_2$	L_3	$W_1 W_4$	$W_2 W_3$	W_5
dim	4	2	2	2	2
$Q_1 Q_2$	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	$L_1 L_2$	L_3	$W_1 W_4$	$W_2 W_3$	W_5	Bipartite graph
	4	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	2	$H_{3/2}^1$
v^5	1	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	2	0	$H_{3/2}^1$
v^7	1	0	1	0	1	$H_{3/2}^1$
v^8	1	0	1	1	0	$H_{3/2}^1$
v^9	1	0	2	0	0	$H_{3/2}^1$

SG 227

Space group $Fd\bar{3}m$, No. 227, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4
dim		1	1	1	1	2	2	3	3	3	3	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	1	0	0	-1	0	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	0	-1	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0
Δ_4	1	0	0	0	1	0	1	0	0	1	0	-1	0	0	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 33.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	2	2	2	2	
v^1	0	0	0	0	0	0	0	1	1	0	1	0	0	2	$H_{5/4}^3$
v^2	0	0	0	0	0	0	0	1	1	0	1	1	0	1	$H_{5/4}^3$
v^3	0	0	0	0	0	0	0	1	1	0	1	2	0	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	0	0	1	0	0	1	2	$H_{5/4}^3$
v^5	0	0	0	0	0	0	1	0	0	1	0	1	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	0	1	0	0	1	0	2	1	0	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	1	0	1	2	$H_{7/6}^3$
v^8	0	0	0	0	0	1	0	1	0	1	1	1	1	1	$H_{7/6}^3$
v^9	0	0	0	0	0	1	0	1	0	1	1	2	1	0	$H_{7/6}^3$
v^{10}	0	0	0	0	1	0	1	0	1	0	1	0	1	2	$H_{7/6}^3$
v^{11}	0	0	0	0	1	0	1	0	1	0	1	1	1	1	$H_{7/6}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	1	2	1	0	$H_{7/6}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	1	0	1	0	$H_{4/4}^2$
v^{14}	0	0	0	1	0	0	0	1	0	0	1	0	0	1	$H_{4/3}^2$
v^{15}	0	0	0	1	0	0	0	1	0	0	1	1	0	0	$H_{4/3}^2$
v^{16}	0	0	0	1	1	0	1	0	0	0	1	0	1	1	$H_{6/5}^1$
v^{17}	0	0	0	1	1	0	1	0	0	0	1	1	1	0	$H_{6/5}^1$
v^{18}	0	0	1	0	0	0	1	0	0	0	0	0	1	1	$H_{4/3}^2$
v^{19}	0	0	1	0	0	0	1	0	0	0	0	1	1	0	$H_{4/3}^2$
v^{20}	0	0	1	0	0	1	0	1	0	0	1	0	1	1	$H_{6/5}^1$
v^{21}	0	0	1	0	0	1	0	1	0	0	1	1	1	0	$H_{6/5}^1$
v^{22}	0	1	0	0	0	0	0	0	0	1	0	0	1	1	$H_{4/3}^2$
v^{23}	0	1	0	0	0	0	0	0	0	1	0	1	1	0	$H_{4/3}^2$
v^{24}	0	1	0	0	1	0	0	0	1	0	1	0	1	1	$H_{6/5}^1$
v^{25}	0	1	0	0	1	0	0	0	1	0	1	1	1	0	$H_{6/5}^1$
v^{26}	0	1	0	1	1	0	0	0	0	0	1	0	1	0	$H_{5/4}^3$
v^{27}	0	1	1	0	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{28}	1	0	0	0	0	0	0	0	1	0	1	0	0	1	$H_{4/3}^2$
v^{29}	1	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{4/3}^2$
v^{30}	1	0	0	0	0	1	0	0	0	1	1	0	1	1	$H_{6/5}^1$
v^{31}	1	0	0	0	0	1	0	0	0	1	1	1	1	0	$H_{6/5}^1$
v^{32}	1	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{33}	1	0	1	0	0	1	0	0	0	0	1	0	1	0	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-
dim		1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2
Λ_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	0	-1	0	0
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1	-1	0	0	0
Λ_3	2	0	0	0	0	1	1	1	1	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{16}$ and the corresponding bipartite graph, number of Hilbert basis: 28.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	Bipartite graph
v^1	1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	$H_{3/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^9	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^{10}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{11}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{14}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{16}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{17}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{18}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{19}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{20}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{21}	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{23}	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{24}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{25}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{26}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{28}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z: $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_1	X_2	X_3	X_4	W_1	W_2
dim	2	2	2	2	2	2
Z_1	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_1	X_2	X_3	X_4	W_1	W_2	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	W_1	W_2
dim	1	1	1	1	2	2	2	2
Q_1	1	1	0	0	1	1	-1	-1
Q_2	1	0	0	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	L_1^+	L_1^-	L_2^+	L_2^-	L_3^+	L_3^-	W_1	W_2	Bipartite graph
	1	1	1	1	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/2}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/2}^1$
v^5	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^7	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^9	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	0	1	0	0	0	1	0	$H_{3/2}^1$

SG 228

Space group $Fd\bar{3}c$, No. 228, without SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4
dim		1	1	1	1	2	2	3	3	3	3	2	2	2	2
Δ_1	1	1	0	0	0	1	0	0	1	0	0	0	-1	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	0	0	-1
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	0	0	-1
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	-1	0	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{14}$ and the corresponding bipartite graph, number of Hilbert basis: 33.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	X_1	X_2	X_3	X_4	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	2	2	2	2	
v^1	0	0	0	0	0	0	0	1	1	0	0	1	2	0	$H_{5/4}^3$
v^2	0	0	0	0	0	0	0	1	1	0	1	1	1	0	$H_{5/4}^3$
v^3	0	0	0	0	0	0	0	1	1	0	2	1	0	0	$H_{5/4}^3$
v^4	0	0	0	0	0	0	1	0	0	1	0	0	2	1	$H_{5/4}^3$
v^5	0	0	0	0	0	0	1	0	0	1	1	0	1	1	$H_{5/4}^3$
v^6	0	0	0	0	0	0	1	0	0	1	2	0	0	1	$H_{5/4}^3$
v^7	0	0	0	0	0	1	0	1	0	1	0	1	2	1	$H_{7/6}^3$
v^8	0	0	0	0	0	1	0	1	0	1	1	1	1	1	$H_{7/6}^3$
v^9	0	0	0	0	0	1	0	1	0	1	2	1	0	1	$H_{7/6}^3$
v^{10}	0	0	0	0	1	0	1	0	1	0	0	1	2	1	$H_{7/6}^3$
v^{11}	0	0	0	0	1	0	1	0	1	0	1	1	1	1	$H_{7/6}^3$
v^{12}	0	0	0	0	1	0	1	0	1	0	2	1	0	1	$H_{7/6}^3$
v^{13}	0	0	0	0	1	1	0	0	0	0	0	1	0	1	$H_{4/4}^2$
v^{14}	0	0	0	1	0	0	0	1	0	0	0	1	1	0	$H_{4/3}^2$
v^{15}	0	0	0	1	0	0	0	1	0	0	1	1	0	0	$H_{4/3}^2$
v^{16}	0	0	0	1	1	0	1	0	0	0	0	1	1	1	$H_{6/5}^1$
v^{17}	0	0	0	1	1	0	1	0	0	0	1	1	0	1	$H_{6/5}^1$
v^{18}	0	0	1	0	0	0	1	0	0	0	0	0	1	1	$H_{4/3}^2$
v^{19}	0	0	1	0	0	0	1	0	0	0	1	0	0	1	$H_{4/3}^2$
v^{20}	0	0	1	0	0	1	0	1	0	0	0	1	1	1	$H_{6/5}^1$
v^{21}	0	0	1	0	0	1	0	1	0	0	1	1	0	1	$H_{6/5}^1$

v^{22}	0	1	0	0	0	0	0	0	0	1	0	0	1	1	$H_{4/3}^2$
v^{23}	0	1	0	0	0	0	0	0	0	1	1	0	0	1	$H_{4/3}^2$
v^{24}	0	1	0	0	1	0	0	0	1	0	0	1	1	1	$H_{6/5}^1$
v^{25}	0	1	0	0	1	0	0	0	1	0	1	1	0	1	$H_{6/5}^1$
v^{26}	0	1	0	1	1	0	0	0	0	0	0	1	0	1	$H_{5/4}^3$
v^{27}	0	1	1	0	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{28}	1	0	0	0	0	0	0	0	1	0	0	1	1	0	$H_{4/3}^2$
v^{29}	1	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{4/3}^2$
v^{30}	1	0	0	0	0	1	0	0	0	1	0	1	1	1	$H_{6/5}^1$
v^{31}	1	0	0	0	0	1	0	0	0	1	1	1	0	1	$H_{6/5}^1$
v^{32}	1	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{3/2}^1$
v^{33}	1	0	1	0	0	1	0	0	0	0	0	1	0	1	$H_{5/4}^3$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1L_2	L_3
	dim	1	1	1	1	2	2	3	3	3	3	4	2
Λ_1	1	1	0	0	1	0	0	0	1	1	0	0	-1
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1
Λ_3	2	0	0	0	0	1	1	1	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 45.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	L_1L_2	L_3	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	4	2	
v^1	0	0	0	0	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^2	0	0	0	0	0	0	0	1	0	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	0	0	1	0	1	0	1	1	$H_{4/4}^2$
v^4	0	0	0	0	0	0	1	1	0	0	1	1	$H_{4/4}^2$
v^5	0	0	0	0	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	0	0	0	1	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^9	0	0	0	1	0	1	1	0	0	0	1	1	$H_{5/4}^3$
v^{10}	0	0	0	1	1	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{11}	0	0	0	1	1	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{12}	0	0	0	2	0	0	0	0	0	2	1	2	$H_{7/6}^3$
v^{13}	0	0	0	2	0	0	1	0	0	1	1	2	$H_{7/6}^3$
v^{14}	0	0	0	2	0	0	2	0	0	0	1	2	$H_{7/6}^3$
v^{15}	0	0	1	0	0	1	0	0	1	0	1	1	$H_{5/4}^3$

v^{16}	0	0	1	0	0	1	0	1	0	0	1	1	$H_{5/4}^3$
v^{17}	0	0	1	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{18}	0	0	1	0	1	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{19}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	0	2	0	0	0	0	0	2	0	1	2	$H_{7/6}^3$
v^{21}	0	0	2	0	0	0	0	1	1	0	1	2	$H_{7/6}^3$
v^{22}	0	0	2	0	0	0	0	2	0	0	1	2	$H_{7/6}^3$
v^{23}	0	1	0	0	0	1	0	0	1	0	1	1	$H_{5/4}^3$
v^{24}	0	1	0	0	0	1	0	1	0	0	1	1	$H_{5/4}^3$
v^{25}	0	1	0	0	1	0	0	0	1	0	1	1	$H_{5/4}^3$
v^{26}	0	1	0	0	1	0	0	1	0	0	1	1	$H_{5/4}^3$
v^{27}	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{28}	0	1	1	0	0	0	0	0	2	0	1	2	$H_{7/6}^3$
v^{29}	0	1	1	0	0	0	0	1	1	0	1	2	$H_{7/6}^3$
v^{30}	0	1	1	0	0	0	0	2	0	0	1	2	$H_{7/6}^3$
v^{31}	0	2	0	0	0	0	0	0	2	0	1	2	$H_{7/6}^3$
v^{32}	0	2	0	0	0	0	0	1	1	0	1	2	$H_{7/6}^3$
v^{33}	0	2	0	0	0	0	0	2	0	0	1	2	$H_{7/6}^3$
v^{34}	1	0	0	0	0	1	0	0	0	1	1	1	$H_{5/4}^3$
v^{35}	1	0	0	0	0	1	1	0	0	0	1	1	$H_{5/4}^3$
v^{36}	1	0	0	0	1	0	0	0	0	1	1	1	$H_{5/4}^3$
v^{37}	1	0	0	0	1	0	1	0	0	0	1	1	$H_{5/4}^3$
v^{38}	1	0	0	1	0	0	0	0	0	2	1	2	$H_{7/6}^3$
v^{39}	1	0	0	1	0	0	1	0	0	1	1	2	$H_{7/6}^3$
v^{40}	1	0	0	1	0	0	2	0	0	0	1	2	$H_{7/6}^3$
v^{41}	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{42}	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{43}	2	0	0	0	0	0	0	0	0	2	1	2	$H_{7/6}^3$
v^{44}	2	0	0	0	0	0	1	0	0	1	1	2	$H_{7/6}^3$
v^{45}	2	0	0	0	0	0	2	0	0	0	1	2	$H_{7/6}^3$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_1	X_2	X_3	X_4	W_1W_2
dim	2	2	2	2	4
Z_1	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	X_1	X_2	X_3	X_4	W_1W_2	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	L_1L_2	L_3	W_1W_2
dim	4	2	4
Q_1Q_2	2	2	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_1L_2	L_3	W_1W_2	
	4	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	0	1	$H_{2/2}^1$

SG 229

Space group $Im\bar{3}m$, No. 229, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	N_1^+	N_1^-	N_2^+	N_2^-	N_3^+	N_3^-	N_4^+	N_4^-	
dim	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	
Σ_1	1	1	0	0	0	1	0	0	1	1	1	-1	0	0	0	0	0	0	-1
Σ_2	1	0	0	1	0	1	0	1	1	0	1	0	0	0	-1	-1	0	0	0
Σ_3	1	0	1	0	0	0	1	1	0	1	1	0	-1	0	0	0	0	-1	0
Σ_4	1	0	0	0	1	0	1	1	1	1	0	0	0	-1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 48.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	N_1^+	N_1^-	N_2^+	N_2^-	N_3^+	N_3^-	N_4^+	N_4^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	1	$H_{4/3}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	1	$H_{4/3}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	1	$H_{4/3}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	1	$H_{4/3}^1$
v^5	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	1	0	$H_{4/3}^1$
v^6	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	0	1	0	$H_{4/3}^1$
v^7	0	0	0	0	0	0	0	0	0	1	1	1	0	0	1	0	0	0	$H_{4/3}^1$
v^8	0	0	0	0	0	0	0	0	0	1	1	1	0	1	0	0	0	0	$H_{4/3}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	1	$H_{4/3}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	1	$H_{4/3}^1$
v^{11}	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	1	$H_{4/3}^1$
v^{12}	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	1	$H_{4/3}^1$
v^{13}	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	1	0	$H_{4/3}^1$
v^{14}	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	1	0	$H_{4/3}^1$
v^{15}	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	1	0	0	$H_{4/3}^1$
v^{16}	0	0	0	0	0	0	0	0	1	0	1	1	1	0	0	0	0	0	$H_{4/3}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	1	$H_{4/3}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	1	$H_{4/3}^1$
v^{19}	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	1	$H_{4/3}^1$
v^{20}	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	1	$H_{4/3}^1$
v^{21}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	1	0	0	$H_{4/3}^1$
v^{22}	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	1	0	0	$H_{4/3}^1$
v^{23}	0	0	0	0	0	0	0	1	0	0	1	0	1	0	1	0	0	0	$H_{4/3}^1$
v^{24}	0	0	0	0	0	0	0	1	0	0	1	0	1	1	0	0	0	0	$H_{4/3}^1$
v^{25}	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	1	1	0	$H_{4/3}^1$
v^{26}	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	1	0	$H_{4/3}^1$
v^{27}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{4/3}^1$
v^{28}	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	0	1	0	$H_{4/3}^1$
v^{29}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	1	0	0	$H_{4/3}^1$
v^{30}	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	1	0	0	$H_{4/3}^1$
v^{31}	0	0	0	0	0	0	1	0	0	0	0	1	1	0	1	0	0	0	$H_{4/3}^1$
v^{32}	0	0	0	0	0	0	1	0	0	0	0	1	1	1	0	0	0	0	$H_{4/3}^1$
v^{33}	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{34}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{35}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{36}	0	0	0	0	0	1	0	0	0	0	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^{37}	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{38}	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$

v^{39}	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{40}	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{41}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{42}	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{43}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{44}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{45}	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{46}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{47}	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{48}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	H_1^+	H_1^-	H_2^+	H_2^-	H_3^+	H_3^-	H_4^+	H_4^-	H_5^+	H_5^-
dim		1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	3	3	3	3
Δ_1	1	1	0	0	0	1	0	0	1	0	0	-1	0	0	0	-1	0	0	-1	0	0
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	-1	0	0	0	-1	-1	0	0	0
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	0	-1	0	0	0	0	-1
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	0	-1	0	-1	0	0	-1	0
Δ_5	2	0	0	0	0	0	0	1	1	1	1	0	0	0	0	0	0	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{20}$ and the corresponding bipartite graph, number of Hilbert basis: 88.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	H_1^+	H_1^-	H_2^+	H_2^-	H_3^+	H_3^-	H_4^+	H_4^-	H_5^+	H_5^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	2	2	3	3	3	3	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/2}^1$
v^4	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/2}^1$
v^5	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/2}^1$
v^6	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	0	0	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	0	0	0	1	0	0	1	1	0	0	1	0	0	0	$H_{5/4}^3$
v^8	0	0	0	0	0	1	0	0	0	1	0	1	1	0	0	0	0	0	1	0	$H_{5/4}^3$
v^9	0	0	0	0	0	1	0	0	0	2	0	0	2	0	0	0	1	0	1	0	$H_{7/6}^3$
v^{10}	0	0	0	0	0	1	0	0	1	0	0	0	0	2	0	0	1	0	0	0	$H_{5/4}^3$
v^{11}	0	0	0	0	0	1	0	1	0	0	1	0	0	1	0	0	1	0	0	0	$H_{5/4}^3$
v^{12}	0	0	0	0	0	1	0	1	0	0	1	1	0	0	0	0	0	0	1	0	$H_{5/4}^3$
v^{13}	0	0	0	0	0	1	0	1	0	1	0	0	0	0	1	0	1	0	1	0	$H_{6/6}^5$
v^{14}	0	0	0	0	0	1	0	1	0	1	1	0	1	0	0	0	1	0	1	0	$H_{7/6}^3$
v^{15}	0	0	0	0	0	1	0	2	0	0	2	0	0	0	0	0	1	0	1	0	$H_{7/6}^3$

v^{16}	0 0 0 0 0 1 1 0 0 0	0 2 0 0 0 0 0 0 1 0	$H_{5/4}^3$
v^{17}	0 0 0 0 0 2 0 1 0 1	0 0 0 2 1 0 2 0 0 0	$H_{9/8}^1$
v^{18}	0 0 0 0 0 2 0 1 0 1	0 2 0 0 1 0 0 0 2 0	$H_{9/8}^1$
v^{19}	0 0 0 0 1 0 0 0 0 0	0 0 0 0 1 0 0 0 0 0	$H_{2/2}^1$
v^{20}	0 0 0 0 1 0 0 0 0 0	1 0 1 0 0 0 0 0 0 0	$H_{3/2}^1$
v^{21}	0 0 0 0 1 0 0 0 0 1	0 0 2 0 0 0 0 1 0 0	$H_{5/4}^3$
v^{22}	0 0 0 0 1 0 0 0 1 0	0 0 1 1 0 0 0 1 0 0	$H_{5/4}^3$
v^{23}	0 0 0 0 1 0 0 0 1 0	1 0 0 1 0 0 0 0 0 1	$H_{5/4}^3$
v^{24}	0 0 0 0 1 0 0 0 2 0	0 0 0 2 0 0 0 1 0 1	$H_{7/6}^3$
v^{25}	0 0 0 0 1 0 0 1 0 0	2 0 0 0 0 0 0 0 0 1	$H_{5/4}^3$
v^{26}	0 0 0 0 1 0 1 0 0 0	0 1 1 0 0 0 0 1 0 0	$H_{5/4}^3$
v^{27}	0 0 0 0 1 0 1 0 0 0	1 1 0 0 0 0 0 0 0 1	$H_{5/4}^3$
v^{28}	0 0 0 0 1 0 1 0 1 0	0 0 0 0 0 1 0 1 0 1	$H_{6/6}^5$
v^{29}	0 0 0 0 1 0 1 0 1 0	0 1 0 1 0 0 0 1 0 1	$H_{7/6}^3$
v^{30}	0 0 0 0 1 0 2 0 0 0	0 2 0 0 0 0 0 1 0 1	$H_{7/6}^3$
v^{31}	0 0 0 0 2 0 1 0 1 0	0 0 2 0 0 1 0 2 0 0	$H_{9/8}^1$
v^{32}	0 0 0 0 2 0 1 0 1 0	2 0 0 0 0 1 0 0 0 2	$H_{9/8}^1$
v^{33}	0 0 0 1 0 0 0 0 0 0	0 0 0 1 0 0 0 0 0 0	$H_{2/1}^1$
v^{34}	0 0 0 1 0 0 0 0 0 1	0 0 1 0 0 0 0 0 1 0	$H_{4/3}^2$
v^{35}	0 0 0 1 0 0 0 1 0 0	1 0 0 0 0 0 0 0 1 0	$H_{4/3}^2$
v^{36}	0 0 0 1 0 0 1 0 0 0	0 1 0 0 0 0 0 0 1 0	$H_{4/3}^2$
v^{37}	0 0 0 1 0 1 0 1 0 1	0 1 0 0 1 0 0 0 2 0	$H_{8/7}^1$
v^{38}	0 0 0 1 1 0 1 0 0 0	0 0 1 0 0 1 0 1 0 0	$H_{6/5}^1$
v^{39}	0 0 0 1 1 0 1 0 0 0	1 0 0 0 0 1 0 0 0 1	$H_{6/5}^1$
v^{40}	0 0 0 1 1 0 2 0 0 0	0 1 0 0 0 1 0 1 0 1	$H_{8/7}^1$
v^{41}	0 0 0 2 0 0 0 1 0 1	0 0 0 0 1 0 0 0 2 0	$H_{7/6}^3$
v^{42}	0 0 0 2 0 0 1 0 0 0	0 0 0 0 0 1 0 0 1 0	$H_{5/4}^3$
v^{43}	0 0 0 2 1 0 2 0 0 0	0 0 0 0 0 2 0 1 0 1	$H_{9/8}^1$
v^{44}	0 0 1 0 0 0 0 0 0 0	0 0 1 0 0 0 0 0 0 0	$H_{2/1}^1$
v^{45}	0 0 1 0 0 0 0 0 1 0	0 0 0 1 0 0 0 0 0 1	$H_{4/3}^2$
v^{46}	0 0 1 0 0 0 0 1 0 0	1 0 0 0 0 0 0 0 0 1	$H_{4/3}^2$
v^{47}	0 0 1 0 0 0 1 0 0 0	0 1 0 0 0 0 0 0 0 1	$H_{4/3}^2$
v^{48}	0 0 1 0 0 1 0 1 0 0	0 0 0 1 1 0 1 0 0 0	$H_{6/5}^1$
v^{49}	0 0 1 0 0 1 0 1 0 0	0 1 0 0 1 0 0 0 1 0	$H_{6/5}^1$
v^{50}	0 0 1 0 0 1 0 2 0 0	1 0 0 0 1 0 1 0 1 0	$H_{8/7}^1$
v^{51}	0 0 1 0 1 0 1 0 1 0	1 0 0 0 0 1 0 0 0 2	$H_{8/7}^1$
v^{52}	0 0 1 1 0 0 0 1 0 0	0 0 0 0 1 0 0 0 1 0	$H_{5/4}^3$
v^{53}	0 0 1 1 0 0 1 0 0 0	0 0 0 0 0 1 0 0 0 1	$H_{5/4}^3$
v^{54}	0 0 2 0 0 0 0 1 0 0	0 0 0 0 1 0 0 0 0 1	$H_{5/4}^3$
v^{55}	0 0 2 0 0 0 1 0 1 0	0 0 0 0 0 1 0 0 0 2	$H_{7/6}^3$
v^{56}	0 0 2 0 0 1 0 2 0 0	0 0 0 0 2 0 1 0 1 0	$H_{9/8}^1$
v^{57}	0 1 0 0 0 0 0 0 0 0	0 1 0 0 0 0 0 0 0 0	$H_{2/1}^1$

v ⁵⁸	0	1	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	H _{4/3} ²
v ⁵⁹	0	1	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	H _{4/3} ²
v ⁶⁰	0	1	0	0	0	0	0	1	0	0	1	0	0	0	1	0	0	0	H _{4/3} ²
v ⁶¹	0	1	0	0	0	1	0	1	0	1	0	0	1	1	0	2	0	0	H _{8/7} ¹
v ⁶²	0	1	0	0	1	0	0	0	1	0	0	0	1	0	1	0	0	0	H _{6/5} ¹
v ⁶³	0	1	0	0	1	0	0	0	1	0	1	0	0	0	1	0	0	0	H _{6/5} ¹
v ⁶⁴	0	1	0	0	1	0	0	0	2	0	0	0	1	0	1	0	1	0	H _{8/7} ¹
v ⁶⁵	0	1	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	H _{3/2} ¹
v ⁶⁶	0	1	0	1	0	0	0	1	0	1	0	0	0	1	0	1	0	1	H _{7/6} ³
v ⁶⁷	0	1	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	H _{5/4} ³
v ⁶⁸	0	1	1	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	H _{5/4} ³
v ⁶⁹	0	2	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	H _{5/4} ³
v ⁷⁰	0	2	0	0	0	0	0	1	0	1	0	0	0	1	0	2	0	0	H _{7/6} ³
v ⁷¹	0	2	0	0	1	0	0	0	2	0	0	0	0	0	2	0	1	0	H _{9/8} ¹
v ⁷²	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	H _{2/1} ¹
v ⁷³	1	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	H _{4/3} ²
v ⁷⁴	1	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	1	0	H _{4/3} ²
v ⁷⁵	1	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	H _{4/3} ²
v ⁷⁶	1	0	0	0	0	1	0	0	0	1	0	0	1	1	0	1	0	0	H _{6/5} ¹
v ⁷⁷	1	0	0	0	0	1	0	0	0	1	0	0	1	0	0	0	1	0	H _{6/5} ¹
v ⁷⁸	1	0	0	0	0	1	0	0	0	2	0	0	1	0	1	0	1	0	H _{8/7} ¹
v ⁷⁹	1	0	0	0	1	0	1	0	1	0	0	0	1	0	2	0	0	0	H _{8/7} ¹
v ⁸⁰	1	0	0	1	0	0	0	0	0	1	0	0	0	1	0	0	1	0	H _{5/4} ³
v ⁸¹	1	0	0	1	0	0	1	0	0	0	0	0	1	0	1	0	0	0	H _{5/4} ³
v ⁸²	1	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	H _{3/2} ¹
v ⁸³	1	0	1	0	0	0	1	0	1	0	0	0	1	0	1	0	1	0	H _{7/6} ³
v ⁸⁴	1	1	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	H _{5/4} ³
v ⁸⁵	1	1	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	0	H _{5/4} ³
v ⁸⁶	2	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0	0	H _{5/4} ³
v ⁸⁷	2	0	0	0	0	0	1	0	1	0	0	0	1	0	2	0	0	0	H _{7/6} ³
v ⁸⁸	2	0	0	0	0	1	0	0	0	2	0	0	1	0	1	0	1	0	H _{9/8} ¹

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	P_1	P_2	P_3	P_4	P_5
	dim	1	1	1	1	2	2	3	3	3	3	1	1	2	3	3
Λ_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	0	0	-1
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1	0	-1	0
Λ_3	2	0	0	0	0	1	1	1	1	1	1	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{15}$ and the corresponding bipartite graph, number of Hilbert basis: 30.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	P_1	P_2	P_3	P_4	P_5	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	2	3	3	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^6	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^8	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{11}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	1	0	0	0	0	0	1	0	1	0	0	1	$H_{4/3}^2$
v^{13}	0	0	0	1	0	0	1	0	0	0	0	1	0	0	1	$H_{4/3}^2$
v^{14}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	0	0	1	0	0	0	0	0	1	0	1	0	0	1	0	$H_{4/3}^2$
v^{18}	0	0	1	0	0	0	0	1	0	0	1	0	0	1	0	$H_{4/3}^2$
v^{19}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{20}	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	1	0	1	0	0	1	0	$H_{4/3}^2$
v^{23}	0	1	0	0	0	0	0	1	0	0	1	0	0	1	0	$H_{4/3}^2$
v^{24}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	0	1	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{26}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	0	0	0	0	1	0	1	0	0	1	$H_{4/3}^2$
v^{28}	1	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{4/3}^2$
v^{29}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{30}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$
A matrix

corep		H_1^+	H_1^-	H_2^+	H_2^-	H_3^+	H_3^-	H_4^+	H_4^-	H_5^+	H_5^-	N_1^+	N_1^-	N_2^+	N_2^-	N_3^+	N_3^-	N_4^+	N_4^-
dim		1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1
G_1	1	1	0	0	0	1	0	0	1	1	1	-1	0	0	-1	0	0	0	0
G_2	1	0	0	0	1	0	1	1	1	1	0	0	0	0	0	-1	-1	0	0
G_3	1	0	1	0	0	0	1	1	0	1	1	0	-1	-1	0	0	0	0	0
G_4	1	0	0	1	0	1	0	1	1	0	1	0	0	0	0	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{18}$ and the corresponding bipartite graph, number of Hilbert basis: 48.

	H_1^+	H_1^-	H_2^+	H_2^-	H_3^+	H_3^-	H_4^+	H_4^-	H_5^+	H_5^-	N_1^+	N_1^-	N_2^+	N_2^-	N_3^+	N_3^-	N_4^+	N_4^-	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	1	1	0	0	0	1	$H_{4/3}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	0	1	1	1	0	0	0	$H_{4/3}^1$
v^3	0	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	1	$H_{4/3}^1$
v^4	0	0	0	0	0	0	0	0	0	1	0	1	0	1	1	0	0	0	$H_{4/3}^1$
v^5	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	1	$H_{4/3}^1$
v^6	0	0	0	0	0	0	0	0	0	1	1	0	1	0	1	0	0	0	$H_{4/3}^1$
v^7	0	0	0	0	0	0	0	0	0	1	1	1	0	0	0	0	0	1	$H_{4/3}^1$
v^8	0	0	0	0	0	0	0	0	0	1	1	1	0	0	1	0	0	0	$H_{4/3}^1$
v^9	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	1	0	$H_{4/3}^1$
v^{10}	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	1	0	0	$H_{4/3}^1$
v^{11}	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	1	0	$H_{4/3}^1$
v^{12}	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	1	0	0	$H_{4/3}^1$
v^{13}	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	1	0	$H_{4/3}^1$
v^{14}	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	1	0	0	$H_{4/3}^1$
v^{15}	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	1	0	$H_{4/3}^1$
v^{16}	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	1	0	0	$H_{4/3}^1$
v^{17}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	1	$H_{4/3}^1$
v^{18}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	1	$H_{4/3}^1$
v^{19}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	0	1	0	$H_{4/3}^1$
v^{20}	0	0	0	0	0	0	0	1	0	0	0	0	0	1	1	1	0	0	$H_{4/3}^1$
v^{21}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	1	1	$H_{4/3}^1$
v^{22}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	1	$H_{4/3}^1$
v^{23}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	1	0	$H_{4/3}^1$
v^{24}	0	0	0	0	0	0	0	1	0	0	1	0	0	0	1	1	0	0	$H_{4/3}^1$
v^{25}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	1	$H_{4/3}^1$
v^{26}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	1	$H_{4/3}^1$
v^{27}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	1	0	$H_{4/3}^1$
v^{28}	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	1	0	0	$H_{4/3}^1$
v^{29}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	1	1	$H_{4/3}^1$
v^{30}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	0	1	$H_{4/3}^1$
v^{31}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	1	0	$H_{4/3}^1$

v^{32}	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	$H_{4/3}^1$
v^{33}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{34}	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{35}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{36}	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{37}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{38}	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{39}	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{3/2}^1$
v^{40}	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{41}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{42}	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{43}	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{44}	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{45}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{46}	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{47}	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{48}	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_1^+	H_1^-	H_2^+	H_2^-	H_3^+	H_3^-	H_4^+	H_4^-	H_5^+	H_5^-	P_1	P_2	P_3	P_4	P_5
	dim	1	1	1	1	2	2	3	3	3	3	1	1	2	3	3
F_1	1	1	0	0	1	0	0	0	1	1	0	-1	0	0	0	-1
F_2	1	0	1	1	0	0	0	1	0	0	1	0	-1	0	-1	0
F_3	2	0	0	0	0	1	1	1	1	1	1	0	0	-1	-1	-1

ker $M \cap \mathbb{Z}_{\geq 0}^{15}$ and the corresponding bipartite graph, number of Hilbert basis: 30.

	H_1^+	H_1^-	H_2^+	H_2^-	H_3^+	H_3^-	H_4^+	H_4^-	H_5^+	H_5^-	P_1	P_2	P_3	P_4	P_5	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	1	1	2	3	3	
v^1	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^6	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	$H_{2/2}^1$
v^8	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^9	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$

v^{11}	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	0	1	0	0	0	0	0	1	0	1	0	0	1	$H_{4/3}^2$
v^{13}	0	0	0	1	0	0	1	0	0	0	0	1	0	0	1	$H_{4/3}^2$
v^{14}	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{15}	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	0	0	1	0	0	0	0	0	1	0	1	0	0	1	0	$H_{4/3}^2$
v^{18}	0	0	1	0	0	0	0	1	0	0	1	0	0	1	0	$H_{4/3}^2$
v^{19}	0	0	1	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{20}	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	0	0	1	0	1	0	0	1	0	$H_{4/3}^2$
v^{23}	0	1	0	0	0	0	0	1	0	0	1	0	0	1	0	$H_{4/3}^2$
v^{24}	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	0	1	0	0	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{26}	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	0	0	0	0	1	0	1	0	0	1	$H_{4/3}^2$
v^{28}	1	0	0	0	0	0	1	0	0	0	0	1	0	0	1	$H_{4/3}^2$
v^{29}	1	0	0	0	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{30}	1	0	0	0	1	0	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_1	P_2	P_3	P_4	P_5	N_1^+	N_1^-	N_2^+	N_2^-	N_3^+	N_3^-	N_4^+	N_4^-
	dim	1	1	2	3	3	1	1	1	1	1	1	1	1
D_1	1	1	0	1	0	1	-1	0	0	0	0	-1	0	0
D_2	1	0	0	0	1	1	0	0	-1	0	0	0	0	-1
D_3	1	0	1	1	1	0	0	-1	0	0	-1	0	0	0
D_4	1	0	0	0	1	1	0	0	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 24.

	P_1	P_2	P_3	P_4	P_5	N_1^+	N_1^-	N_2^+	N_2^-	N_3^+	N_3^-	N_4^+	N_4^-	Bipartite graph
	1	1	2	3	3	1	1	1	1	1	1	1	1	
v^1	0	0	0	0	1	0	0	0	0	0	1	1	1	$H_{4/3}^1$
v^2	0	0	0	0	1	0	0	0	1	0	1	0	1	$H_{4/3}^1$
v^3	0	0	0	0	1	0	0	1	0	0	1	1	0	$H_{4/3}^1$
v^4	0	0	0	0	1	0	0	1	1	0	1	0	0	$H_{4/3}^1$
v^5	0	0	0	0	1	1	0	0	0	0	0	1	1	$H_{4/3}^1$
v^6	0	0	0	0	1	1	0	0	1	0	0	0	1	$H_{4/3}^1$

v^7	0	0	0	0	1	1	0	1	0	0	0	1	0	$H_{4/3}^1$
v^8	0	0	0	0	1	1	0	1	1	0	0	0	0	$H_{4/3}^1$
v^9	0	0	0	1	0	0	0	0	0	1	0	1	1	$H_{4/3}^1$
v^{10}	0	0	0	1	0	0	0	0	1	1	0	0	1	$H_{4/3}^1$
v^{11}	0	0	0	1	0	0	0	1	0	1	0	1	0	$H_{4/3}^1$
v^{12}	0	0	0	1	0	0	0	1	1	1	0	0	0	$H_{4/3}^1$
v^{13}	0	0	0	1	0	0	1	0	0	0	0	1	1	$H_{4/3}^1$
v^{14}	0	0	0	1	0	0	1	0	1	0	0	0	1	$H_{4/3}^1$
v^{15}	0	0	0	1	0	0	1	1	0	0	0	1	0	$H_{4/3}^1$
v^{16}	0	0	0	1	0	0	1	1	1	0	0	0	0	$H_{4/3}^1$
v^{17}	0	0	1	0	0	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{18}	0	0	1	0	0	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^{19}	0	0	1	0	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^{20}	0	0	1	0	0	1	1	0	0	0	0	0	0	$H_{3/2}^1$
v^{21}	0	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	1	0	0	0	0	0	0	$H_{2/1}^1$
v^{23}	1	0	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	1	0	0	0	0	0	0	0	$H_{2/1}^1$

SG 230

Space group $Ia\bar{3}d$, No. 230, without SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	N_1	N_2
	dim	1	1	1	1	2	2	3	3	3	3	2	2
Σ_1	1	1	0	0	0	1	0	0	1	1	1	-1	0
Σ_2	1	0	0	1	0	1	0	1	1	0	1	0	-1
Σ_3	1	0	1	0	0	0	1	1	0	1	1	-1	0
Σ_4	1	0	0	0	1	0	1	1	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	N_1	N_2	
	1	1	1	1	2	2	3	3	3	3	2	2	Bipartite graph
v^1	0	0	0	0	0	0	0	0	1	1	2	1	$H_{5/6}^2$
v^2	0	0	0	0	0	0	1	1	0	0	1	2	$H_{5/6}^2$
v^3	0	0	0	0	0	1	0	1	0	1	2	2	$H_{7/8}^{10}, H_{7/8}^3$
v^4	0	0	0	0	1	0	1	0	1	0	2	2	$H_{7/8}^{10}, H_{7/8}^3$

v^5	0	0	0	0	1	1	0	0	0	0	1	1	$H_{4/4}^2$
v^6	0	0	0	1	0	0	0	0	0	1	1	1	$H_{4/4}^4$
v^7	0	0	0	1	1	0	1	0	0	0	1	2	$H_{6/6}^8, H_{6/6}^7$
v^8	0	0	1	0	0	0	0	0	1	0	1	1	$H_{4/4}^4$
v^9	0	0	1	0	0	1	0	1	0	0	1	2	$H_{6/6}^8, H_{6/6}^7$
v^{10}	0	0	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	0	1	0	0	1	1	$H_{4/4}^4$
v^{12}	0	1	0	0	1	0	0	0	1	0	2	1	$H_{6/6}^8, H_{6/6}^7$
v^{13}	0	1	0	1	1	0	0	0	0	0	1	1	$H_{5/4}^3$
v^{14}	1	0	0	0	0	0	1	0	0	0	1	1	$H_{4/4}^4$
v^{15}	1	0	0	0	0	1	0	0	0	1	2	1	$H_{6/6}^8, H_{6/6}^7$
v^{16}	1	0	1	0	0	1	0	0	0	0	1	1	$H_{5/4}^3$
v^{17}	1	1	0	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	H_1	H_2	H_3	H_4
dim		1	1	1	1	2	2	3	3	3	3	2	4	6	
Δ_1	1	1	0	0	0	1	0	0	1	0	0	0	0	-1	
Δ_2	1	0	1	0	0	0	1	1	0	0	0	0	0	-1	
Δ_3	1	0	0	1	0	1	0	0	0	0	1	0	0	-1	
Δ_4	1	0	0	0	1	0	1	0	0	1	0	0	0	-1	
Δ_5	2	0	0	0	0	0	0	1	1	1	1	-1	-2	-1	

$\ker M \cap \mathbb{Z}_{\geq 0}^{13}$ and the corresponding bipartite graph, number of Hilbert basis: 46.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	H_1	H_2	H_3	H_4	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	2	4	6		
v^1	0	0	0	0	0	0	1	1	1	1	1	1	1		$H_{7/8}^{13}$
v^2	0	0	0	0	0	0	1	1	1	1	3	0	1		$H_{8/8}^2$
v^3	0	0	0	0	0	0	2	2	2	2	0	3	2		$H_{13/16}^1, H_{13/16}^5, H_{13/16}^2, H_{13/16}^4, H_{13/16}^3$
v^4	0	0	0	0	0	1	0	1	0	1	1	0	1		$H_{5/6}^6$
v^5	0	0	0	0	0	1	1	2	1	2	0	2	2		$H_{11/14}^1, H_{11/14}^6, H_{11/14}^5, H_{11/14}^3, H_{11/14}^2, H_{11/14}^4$
v^6	0	0	0	0	0	2	0	2	0	2	0	1	2		$H_{9/12}^{12}, H_{9/12}^{11}$
v^7	0	0	0	0	1	0	1	0	1	0	1	0	1		$H_{5/6}^6$
v^8	0	0	0	0	1	0	2	1	2	1	0	2	2		$H_{11/14}^6, H_{11/14}^5, H_{11/14}^1, H_{11/14}^2, H_{11/14}^3, H_{11/14}^4$
v^9	0	0	0	0	1	1	1	1	1	1	0	1	2		$H_{9/12}^{11}, H_{9/12}^{13}, H_{9/12}^{12}, H_{9/12}^{14}$
v^{10}	0	0	0	0	1	2	0	1	0	1	0	0	2		$H_{7/10}^1$
v^{11}	0	0	0	0	2	0	2	0	2	0	0	1	2		$H_{9/12}^{12}, H_{9/12}^{11}$
v^{12}	0	0	0	0	2	1	1	0	1	0	0	0	2		$H_{7/10}^1$

v ¹³	0	0	0	1	0	0	1	1	0	1	0	1	1	$H_{6/7}^1$
v ¹⁴	0	0	0	1	0	0	1	1	0	1	2	0	1	$H_{7/7}^1$
v ¹⁵	0	0	0	1	1	0	1	0	0	0	0	0	1	$H_{4/5}^1$
v ¹⁶	0	0	1	0	0	0	1	1	1	0	0	1	1	$H_{6/7}^1$
v ¹⁷	0	0	1	0	0	0	1	1	1	0	2	0	1	$H_{7/7}^1$
v ¹⁸	0	0	1	0	0	1	0	1	0	0	0	0	1	$H_{4/5}^1$
v ¹⁹	0	0	1	1	0	0	1	1	0	0	1	0	1	$H_{6/6}^{10}$
v ²⁰	0	0	2	2	0	0	2	2	0	0	0	1	2	$H_{11/12}^3, H_{11/12}^4$
v ²¹	0	1	0	0	0	0	0	1	1	1	0	1	1	$H_{6/7}^1$
v ²²	0	1	0	0	0	0	0	1	1	1	2	0	1	$H_{7/7}^1$
v ²³	0	1	0	0	1	0	0	0	1	0	0	0	1	$H_{4/5}^1$
v ²⁴	0	1	0	1	0	0	0	1	0	1	1	0	1	$H_{6/6}^{10}$
v ²⁵	0	1	0	1	0	1	0	2	0	2	0	1	2	$H_{10/12}^{12}, H_{10/12}^8, H_{10/12}^{11}, H_{10/12}^{10}, H_{10/12}^9$
v ²⁶	0	1	0	1	1	1	0	1	0	1	0	0	2	$H_{8/10}^{12}, H_{8/10}^{11}$
v ²⁷	0	1	1	0	0	0	0	1	1	0	1	0	1	$H_{6/6}^{10}$
v ²⁸	0	1	1	1	0	0	0	1	0	0	0	0	1	$H_{5/5}^4$
v ²⁹	0	2	0	2	0	0	0	2	0	2	0	1	2	$H_{11/12}^3, H_{11/12}^4$
v ³⁰	0	2	0	2	1	0	0	1	0	1	0	0	2	$H_{9/10}^1$
v ³¹	0	2	2	0	0	0	0	2	2	0	0	1	2	$H_{11/12}^3, H_{11/12}^4$
v ³²	1	0	0	0	0	0	1	0	1	1	0	1	1	$H_{6/7}^1$
v ³³	1	0	0	0	0	0	1	0	1	1	2	0	1	$H_{7/7}^1$
v ³⁴	1	0	0	0	0	1	0	0	0	1	0	0	1	$H_{4/5}^1$
v ³⁵	1	0	0	1	0	0	1	0	0	1	1	0	1	$H_{6/6}^{10}$
v ³⁶	1	0	1	0	0	0	1	0	1	0	1	0	1	$H_{6/6}^{10}$
v ³⁷	1	0	1	0	1	0	2	0	2	0	0	1	2	$H_{10/12}^{12}, H_{10/12}^8, H_{10/12}^{11}, H_{10/12}^{10}, H_{10/12}^9$
v ³⁸	1	0	1	0	1	1	1	0	1	0	0	0	2	$H_{8/10}^{12}, H_{8/10}^{11}$
v ³⁹	1	0	1	1	0	0	1	0	0	0	0	0	1	$H_{5/5}^4$
v ⁴⁰	1	1	0	0	0	0	0	0	1	1	1	0	1	$H_{6/6}^{10}$
v ⁴¹	1	1	0	1	0	0	0	0	0	1	0	0	1	$H_{5/5}^4$
v ⁴²	1	1	1	0	0	0	0	0	1	0	0	0	1	$H_{5/5}^4$
v ⁴³	2	0	0	2	0	0	2	0	0	2	0	1	2	$H_{11/12}^3, H_{11/12}^4$
v ⁴⁴	2	0	2	0	0	0	2	0	2	0	0	1	2	$H_{11/12}^3, H_{11/12}^4$
v ⁴⁵	2	0	2	0	0	1	1	0	1	0	0	0	2	$H_{9/10}^1$
v ⁴⁶	2	2	0	0	0	0	0	0	2	2	0	1	2	$H_{11/12}^3, H_{11/12}^4$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$P_1 P_2$	P_3
dim	1	1	1	1	2	2	3	3	3	3	4	4

Λ_1	1	1	0	0	1	0	0	0	1	1	0	0	-1
Λ_2	1	0	1	1	0	0	0	1	0	0	1	0	-1
Λ_3	2	0	0	0	0	1	1	1	1	1	1	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 36.

	Γ_1^+	Γ_1^-	Γ_2^+	Γ_2^-	Γ_3^+	Γ_3^-	Γ_4^+	Γ_4^-	Γ_5^+	Γ_5^-	$P_1 P_2$	P_3	Bipartite graph
	1	1	1	1	2	2	3	3	3	3	4	4	
v^1	0	0	0	0	0	0	0	0	2	2	1	2	$H_{7/8}^3, H_{7/8}^{10}$
v^2	0	0	0	0	0	0	0	1	1	2	1	2	$H_{7/8}^3, H_{7/8}^{10}$
v^3	0	0	0	0	0	0	0	2	0	2	1	2	$H_{7/8}^3, H_{7/8}^{10}$
v^4	0	0	0	0	0	0	1	0	2	1	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^5	0	0	0	0	0	0	1	1	1	1	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^6	0	0	0	0	0	0	1	2	0	1	1	2	$H_{7/8}^{10}, H_{7/8}^3$
v^7	0	0	0	0	0	0	2	0	2	0	1	2	$H_{7/8}^3, H_{7/8}^{10}$
v^8	0	0	0	0	0	0	2	1	1	0	1	2	$H_{7/8}^3, H_{7/8}^{10}$
v^9	0	0	0	0	0	0	2	2	0	0	1	2	$H_{7/8}^3, H_{7/8}^{10}$
v^{10}	0	0	0	0	0	1	0	0	1	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{11}	0	0	0	0	0	1	0	1	0	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{12}	0	0	0	0	0	1	1	0	1	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{13}	0	0	0	0	0	1	1	1	0	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{14}	0	0	0	0	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	0	0	0	1	0	0	0	1	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{16}	0	0	0	0	1	0	0	1	0	1	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{17}	0	0	0	0	1	0	1	0	1	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{18}	0	0	0	0	1	0	1	1	0	0	1	1	$H_{5/5}^3, H_{5/5}^2$
v^{19}	0	0	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{20}	0	0	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	1	0	1	$H_{3/3}^1$
v^{22}	0	0	0	1	0	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{23}	0	0	1	0	0	0	0	0	1	0	0	1	$H_{3/3}^1$
v^{24}	0	0	1	0	0	0	0	1	0	0	0	1	$H_{3/3}^1$
v^{25}	0	0	1	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{26}	0	0	1	1	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{27}	0	1	0	0	0	0	0	0	1	0	0	1	$H_{3/3}^1$
v^{28}	0	1	0	0	0	0	0	1	0	0	0	1	$H_{3/3}^1$
v^{29}	0	1	0	1	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{30}	0	1	0	1	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{31}	1	0	0	0	0	0	0	0	0	1	0	1	$H_{3/3}^1$
v^{32}	1	0	0	0	0	0	1	0	0	0	0	1	$H_{3/3}^1$
v^{33}	1	0	1	0	0	1	0	0	0	0	0	1	$H_{4/3}^1$
v^{34}	1	0	1	0	1	0	0	0	0	0	0	1	$H_{4/3}^1$
v^{35}	1	1	0	0	0	1	0	0	0	0	0	1	$H_{4/3}^1$

$$\overline{v^{36} \mid 1 \ 1 \ 0 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ \mid 0 \ 1 \ \mid H_{4/3}^1}$$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep	H_1	H_2H_3	H_4	N_1	N_2
dim	2	4	6	2	2
G_1	2	1	2	3	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	H_1	H_2H_3	H_4	N_1	N_2	
	2	4	6	2	2	Bipartite graph
v^1	0	0	1	0	3	$H_{4/3}^1$
v^2	0	0	1	1	2	$H_{4/3}^1$
v^3	0	0	1	2	1	$H_{4/3}^1$
v^4	0	0	1	3	0	$H_{4/3}^1$
v^5	0	1	0	0	2	$H_{3/2}^1$
v^6	0	1	0	1	1	$H_{3/2}^1$
v^7	0	1	0	2	0	$H_{3/2}^1$
v^8	1	0	0	0	1	$H_{2/1}^1$
v^9	1	0	0	1	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	H_1	H_2H_3	H_4	P_1P_2	P_3
dim	2	4	6	4	4
F_1	1	1	0	1	0 -1
F_2	1	1	0	1	0 -1
F_3	2	0	2	2	-2 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	H_1	H_2H_3	H_4	P_1P_2	P_3	
	2	4	6	4	4	Bipartite graph
v^1	0	0	2	1	2	$H_{5/8}^2, H_{5/8}^1$
v^2	0	1	0	1	0	$H_{2/2}^1$
v^3	1	0	1	0	2	$H_{4/6}^1$
v^4	2	1	0	0	2	$H_{5/6}^2$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D : $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		$P_1 P_2$	P_3	N_1	N_2
	dim	4	4	2	2
D_1	2	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$P_1 P_2$	P_3	N_1	N_2	
	4	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	2	$H_{3/2}^1$
v^5	1	0	1	1	$H_{3/2}^1$
v^6	1	0	2	0	$H_{3/2}^1$

S7. HILBERT BASIS AND BIPARTITE GRAPH IN 230 SPACE GROUPS WITH SOC

SG 1

Space group $P1$, No. 1, with SOC

SG 2

Space group $P\bar{1}$, No. 2, with SOC

SG 3

Space group $P2$, No. 3, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_3 \Gamma_4$	$Z_3 Z_4$
	dim	2	2
Λ_3	1	1	-1
Λ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_3\Gamma_4$	Z_3Z_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		B_3B_4	D_3D_4
dim		2	2
V_3	1	1	-1
V_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	B_3B_4	D_3D_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_3Y_4	C_3C_4
dim		2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	C_3C_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U : $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_3A_4	E_3E_4
	dim	2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_3A_4	E_3E_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 4

Space group $P2_1$, No. 4, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_3\Gamma_4$	Z_3Z_3	Z_4Z_4
	dim	2	2	2
Λ_3	1	1	-2	0
Λ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_3\Gamma_4$	Z_3Z_3	Z_4Z_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		B_3B_4	D_3D_3	D_4D_4
	dim	2	2	2
V_3	1	1	-2	0
V_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	B_3B_4	D_3D_3	D_4D_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_3Y_4	C_3C_3	C_4C_4
	dim	2	2	2
W_3	1	1	-2	0
W_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	C_3C_3	C_4C_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U : $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_3A_4	E_3E_3	E_4E_4
	dim	2	2	2
U_3	1	1	-2	0
U_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_3A_4	E_3E_3	E_4E_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

SG 5

Space group $C2$, No. 5, with SOC

Λ

High symmetry line $\Lambda(0, -u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		$\Gamma_3\Gamma_4$	Z_3Z_4
dim		2	2
Λ_3	1	1	-1
Λ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		$\Gamma_3\Gamma_4$	Z_3Z_4	
		2	2	Bipartite graph
v^1	1	1	1	$H_{2/2}^1$

U

High symmetry line $U(-\frac{1}{2}, -u, u)$, high symmetry points at both ends of U : $A(-\frac{1}{2}, 0, 0), M(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_3A_4	M_3M_4
dim		2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		A_3A_4	M_3M_4	
		2	2	Bipartite graph
v^1	1	1	1	$H_{2/2}^1$

SG 6

Space group Pm , No. 6, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_3\Gamma_4$	Z_3Z_4
dim		2	2
Λ_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_3\Gamma_4$	Z_3Z_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		B_3B_4	D_3D_4
dim		2	2
V_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	B_3B_4	D_3D_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_3Y_4	C_3C_4
dim		2	2
W_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	C_3C_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U : $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_3A_4	E_3E_4
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_3A_4	E_3E_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 7

Space group Pc , No. 7, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_3\Gamma_4$	Z_3Z_4
dim		2	2
Λ_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_3\Gamma_4$	Z_3Z_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		B_3B_3	B_4B_4	D_3D_3	D_4D_4
dim		2	2	2	2
V_2V_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	B_3B_3	B_4B_4	D_3D_3	D_4D_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_3Y_4	C_3C_4
	dim	2	2
W_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	C_3C_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U: $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_3A_3	A_4A_4	E_3E_3	E_4E_4
	dim	2	2	2	2
U_2U_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_3A_3	A_4A_4	E_3E_3	E_4E_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 8

Space group Cm , No. 8, with SOC

Λ

High symmetry line $\Lambda(0, -u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		$\Gamma_3\Gamma_4$	Z_3Z_4
dim		2	2
Λ_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_3\Gamma_4$	Z_3Z_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(-\frac{1}{2}, -u, u)$, high symmetry points at both ends of U: $A(-\frac{1}{2}, 0, 0), M(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_3A_4	M_3M_4
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_3A_4	M_3M_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 9

Space group Cc , No. 9, with SOC

Λ

High symmetry line $\Lambda(0, -u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		$\Gamma_3\Gamma_4$	Z_3Z_4
dim		2	2
Λ_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_3\Gamma_4$	Z_3Z_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(-\frac{1}{2}, -u, u)$, high symmetry points at both ends of U : $A(-\frac{1}{2}, 0, 0), M(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_3A_3	A_4A_4	M_3M_3	M_4M_4
dim	2	2	2	2
U_2U_2	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_3A_3	A_4A_4	M_3M_3	M_4M_4	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 10

Space group $P2/m$, No. 10, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$
dim	2	2	2	2
$\Lambda_3\Lambda_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_3^+\Gamma_4^+$	$\Gamma_3^-\Gamma_4^-$	$Z_3^+Z_4^+$	$Z_3^-Z_4^-$	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	$B_3^+ B_4^+$	$B_3^- B_4^-$	$D_3^+ D_4^+$	$D_3^- D_4^-$
dim	2	2	2	2
$V_3 V_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$B_3^+ B_4^+$	$B_3^- B_4^-$	$D_3^+ D_4^+$	$D_3^- D_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Y_3^+ Y_4^+$	$Y_3^- Y_4^-$	$C_3^+ C_4^+$	$C_3^- C_4^-$
dim	2	2	2	2
$W_3 W_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$Y_3^+ Y_4^+$	$Y_3^- Y_4^-$	$C_3^+ C_4^+$	$C_3^- C_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U: $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_3^+ A_4^+$	$A_3^- A_4^-$	$E_3^+ E_4^+$	$E_3^- E_4^-$
dim	2	2	2	2
$U_3 U_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$A_3^+ A_4^+$	$A_3^- A_4^-$	$E_3^+ E_4^+$	$E_3^- E_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 11

Space group $P2_1/m$, No. 11, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$

A matrix

corep	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_2 Z_2$
dim	2	2	4
$\Lambda_3 \Lambda_4$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_2 Z_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V : $B(-\frac{1}{2}, 0, 0)$, $D(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	$B_3^+ B_4^+$	$B_3^- B_4^-$	$D_2 D_2$
dim	2	2	4
$V_3 V_4$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$B_3^+ B_4^+$	$B_3^- B_4^-$	$D_2 D_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Y_3^+ Y_4^+$	$Y_3^- Y_4^-$	$C_2 C_2$
dim	2	2	4
$W_3 W_4$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$Y_3^+ Y_4^+$	$Y_3^- Y_4^-$	$C_2 C_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U: $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_3^+ A_4^+$	$A_3^- A_4^-$	$E_2 E_2$
dim	2	2	4
$U_3 U_4$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$A_3^+ A_4^+$	$A_3^- A_4^-$	$E_2 E_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 12

Space group $C2/m$, No. 12, with SOC

Λ

High symmetry line $\Lambda(0, -u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$
dim	2	2	2	2
$\Lambda_3 \Lambda_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(-\frac{1}{2}, -u, u)$, high symmetry points at both ends of U: $A(-\frac{1}{2}, 0, 0), M(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_3^+ A_4^+$	$A_3^- A_4^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$
dim	2	2	2	2
$U_3 U_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$A_3^+ A_4^+$	$A_3^- A_4^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 13

Space group $P2/c$, No. 13, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$
dim	2	2	2	2
$\Lambda_3 \Lambda_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V: $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	$B_2 B_2$	$D_2 D_2$
dim	4	4
$V_3 V_3$	2	1
$V_4 V_4$	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$B_2 B_2$	$D_2 D_2$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Y_3^+ Y_4^+$	$Y_3^- Y_4^-$	$C_3^+ C_4^+$	$C_3^- C_4^-$
dim	2	2	2	2
$W_3 W_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$Y_3^+ Y_4^+$	$Y_3^- Y_4^-$	$C_3^+ C_4^+$	$C_3^- C_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U : $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$A_2 A_2$	$E_2 E_2$
	dim	4	4
$U_3 U_3$	2	1	-1
$U_4 U_4$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$A_2 A_2$	$E_2 E_2$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 14

Space group $P2_1/c$, No. 14, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_2 Z_2$
	dim	2	2	4
$\Lambda_3 \Lambda_4$	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_2 Z_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

V

High symmetry line $V(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of V: $B(-\frac{1}{2}, 0, 0), D(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	B_2B_2	$D_3^+D_3^+$	$D_3^-D_3^-$	$D_4^+D_4^+$	$D_4^-D_4^-$
dim	4	2	2	2	2
V_3V_3	2	1	-1	-1	0
V_4V_4	2	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	B_2B_2	$D_3^+D_3^+$	$D_3^-D_3^-$	$D_4^+D_4^+$	$D_4^-D_4^-$	Bipartite graph
	4	2	2	2	2	
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $Y(0, \frac{1}{2}, 0), C(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Y_3^+Y_4^+$	$Y_3^-Y_4^-$	C_2C_2
dim	2	2	4
W_3W_4	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$Y_3^+Y_4^+$	$Y_3^-Y_4^-$	C_2C_2	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

U

High symmetry line $U(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of U: $A(\frac{1}{2}, \frac{1}{2}, 0), E(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_2A_2	$E_3^+ E_3^+$	$E_3^- E_3^-$	$E_4^+ E_4^+$	$E_4^- E_4^-$
dim	4	2	2	2	2
U_3U_3	2	1	-1	-1	0
U_4U_4	2	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_2A_2	$E_3^+ E_3^+$	$E_3^- E_3^-$	$E_4^+ E_4^+$	$E_4^- E_4^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

SG 15

Space group $C2/c$, No. 15, with SOC

Λ

High symmetry line $\Lambda(0, -u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$
dim	2	2	2	2
$\Lambda_3\Lambda_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_3^+ \Gamma_4^+$	$\Gamma_3^- \Gamma_4^-$	$Z_3^+ Z_4^+$	$Z_3^- Z_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(-\frac{1}{2}, -u, u)$, high symmetry points at both ends of U : $A(-\frac{1}{2}, 0, 0), M(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_2A_2	M_2M_2
dim	4	4
U_3U_3	2	1
U_4U_4	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_2A_2	M_2M_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 16

Space group $P222$, No. 16, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_5	Y_5
dim	2	2
Δ_3	1	-1
Δ_4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_5	X_5
dim	2	2
Σ_3	1	-1
Σ_4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	Z_5
	dim	2	2
Λ_3	1	1	-1
Λ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_5	S_5
	dim	2	2
C_3	1	1	-1
C_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H : $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_5	T_5
	dim	2	2
H_3	1	1	-1
H_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	S_5
	dim	2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	U_5
	dim	2	2
G_3	1	1	-1
G_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_5	T_5
dim		2	2
B_3	1	1	-1
B_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	U_5
dim		2	2
A_3	1	1	-1
A_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	U_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_5	R_5
dim		2	2
P_3	1	1	-1
P_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_5	R_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_5	R_5
dim		2	2
E_3	1	1	-1
E_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_5	R_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_5	R_5
dim		2	2
Q_3	1	1	-1
Q_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_5	R_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 17

Space group $P222_1$, No. 17, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5	Y_5
dim		2	2
Δ_3	1	1	-1
Δ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	X_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	Z_2Z_3	Z_4Z_5
dim		2	2	2
Λ_3	1	1	0	-2
Λ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_2Z_3	Z_4Z_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_5	S_5
	dim	2	2
C_3	1	1	-1
C_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_5	T_2T_3	T_4T_5
	dim	2	2	2
H_3	1	1	0	-2
H_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_2T_3	T_4T_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	S_5
	dim	2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_5	U_2U_3	U_4U_5
	dim	2	2	2
G_3	1	1	0	-2
G_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_2U_3	U_4U_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5
	dim	2	2	2	2
B_3B_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_2Z_3	Z_4Z_5	U_2U_3	U_4U_5
dim	2	2	2	2
A_3A_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_2Z_3	Z_4Z_5	U_2U_3	U_4U_5	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_2U_3	U_4U_5	R_2R_3	R_4R_5
dim	2	2	2	2
P_3P_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_2U_3	U_4U_5	R_2R_3	R_4R_5	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_2T_3	T_4T_5	R_2R_3	R_4R_5
dim	2	2	2	2
E_3E_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_2T_3	T_4T_5	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_5	R_2R_3	R_4R_5
	dim	2	2	2
Q_3	1	1	0	-2
Q_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_5	R_2R_3	R_4R_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

SG 18

Space group $P2_12_12$, No. 18, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
	dim	2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_5	X_2X_3	X_4X_5
dim	2	2	2
Σ_3	1	1	-2
Σ_4	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_2X_3	X_4X_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5	Z_5
dim	2	2
Λ_3	1	-1
Λ_4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_2Y_3	Y_4Y_5	S_5S_5
dim	2	2	4
C_3C_4	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_2Y_3	Y_4Y_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5
dim	2	2	2	2
H_3H_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_2X_3	X_4X_5	S_5S_5
dim	2	2	4
D_3D_4	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_2X_3	X_4X_5	U_2U_3	U_4U_5
dim	2	2	2	2
G_3G_4	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_2X_3	X_4X_5	U_2U_3	U_4U_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_5	T_2T_3	T_4T_5
dim	2	2	2
B_3	1	1	-2 0
B_4	1	1	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_2T_3	T_4T_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	U_2U_3	U_4U_5
dim	2	2	2
A_3	1	1	0 -2
A_4	1	1	-2 0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	U_2U_3	U_4U_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_2U_3	U_4U_5	R_5R_5
	dim	2	2	4
P_3P_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_2U_3	U_4U_5	R_5R_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_2T_3	T_4T_5	R_5R_5
	dim	2	2	4
E_3E_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	T_2T_3	T_4T_5	R_5R_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$S_5 S_5$	$R_5 R_5$
dim		4	4
$Q_3 Q_3$	2	1	-1
$Q_4 Q_4$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$S_5 S_5$	$R_5 R_5$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 19

Space group $P2_12_12_1$, No. 19, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5	$Y_2 Y_3$	$Y_4 Y_5$
dim		2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	$Y_2 Y_3$	$Y_4 Y_5$	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	$X_2 X_3$	$X_4 X_5$
dim		2	2	2
Σ_3	1	1	0	-2
Σ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_2X_3	X_4X_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	Z_2Z_3	Z_4Z_5
	dim	2	2	2
Λ_3	1	1	0	-2
Λ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_2Z_3	Z_4Z_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_2Y_3	Y_4Y_5	S_5S_5
	dim	2	2	4
C_3C_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_2Y_3	Y_4Y_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H : $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_2Y_3	Y_4Y_5	T_5T_5
dim	2	2	4
H_3H_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_2Y_3	Y_4Y_5	T_5T_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_2X_3	X_4X_5	S_5S_5
dim	2	2	4
D_3D_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_2X_3	X_4X_5	U_5U_5
dim	2	2	4
G_3G_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	U_5U_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_2Z_3	Z_4Z_5	T_5T_5
dim	2	2	4
B_3B_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_2Z_3	Z_4Z_5	T_5T_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_2Z_3	Z_4Z_5	U_5U_5
dim	2	2	4
A_3A_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_2Z_3	Z_4Z_5	U_5U_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_5U_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	
dim	4	2	2	2	2	
P_3P_3	2	1	-1	0	-1	0
P_4P_4	2	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_5U_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	Bipartite graph
	4	2	2	2	2	
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_5T_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	
dim	4	2	2	2	2	
E_3E_3	2	1	0	0	-1	-1
E_4E_4	2	1	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_5T_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	Bipartite graph
	4	2	2	2	2	
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_5S_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5
	dim	4	2	2	2	2
Q_3Q_3	2	1	-1	0	0	-1
Q_4Q_4	2	1	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_5S_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	Bipartite graph
	4	2	2	2	2	
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

SG 20

Space group $C222_1$, No. 20, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_2Z_3	Z_4Z_5
	dim	2	2	2
Λ_3	1	1	0	-2
Λ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_2Z_3	Z_4Z_5	Bipartite graph
	2	2	2	
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	Y_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_5	T_2T_3	T_4T_5
	dim	2	2	2
H_3	1	1	0	-2
H_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_2T_3	T_4T_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5
	dim	2	2	2	2
A_3A_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_3R_3	R_4R_4
dim		2	2	2
D_3	1	1	-2	0
D_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_3	R_4R_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

SG 21

Space group $C222$, No. 21, with SOC

 Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	Z_5
dim		2	2
Λ_3	1	1	-1
Λ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

 Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	Y_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_5	T_5
	dim	2	2
H_3	1	1	-1
H_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	T_5
	dim	2	2
A_3	1	1	-1
A_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_3R_4
dim		2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		S_3S_4	R_3R_4	
		2	2	Bipartite graph
v^1	1	1		$H_{2/2}^1$

SG 22

Space group $F222$, No. 22, with SOC

 Λ

High symmetry line $\Lambda(u, u, 0)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	Z_5
dim		2	2
Λ_3	1	1	-1
Λ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		Γ_5	Z_5	
		2	2	Bipartite graph
v^1	1	1		$H_{2/2}^1$

 Σ

High symmetry line $\Sigma(u, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, -u, -u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(0, -\frac{1}{2}, -\frac{1}{2})$

A matrix

corep		Γ_5	Y_5
	dim	2	2
Δ_3	1	1	-1
Δ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 23

Space group *I222*, No. 23, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		R_3R_4	W_5
	dim	2	2
Q_3	1	1	-1
Q_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	W_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		S_3S_4	W_5
	dim	2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	W_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		T_3T_4	W_5
	dim	2	2
P_3	1	1	-1
P_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	W_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 24

Space group $I2_12_12_1$, No. 24, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q : $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		R_3R_4	W_2	W_3	W_4	W_5
	dim	2	1	1	1	1
Q_3	1	1	0	-1	0	-1
Q_4	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_3R_4	W_2	W_3	W_4	W_5	
	2	1	1	1	1	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	S_3S_4	W_2	W_3	W_4	W_5	
dim	2	1	1	1	1	
D_3	1	1	0	-1	-1	0
D_4	1	1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_3S_4	W_2	W_3	W_4	W_5	
	2	1	1	1	1	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	T_3T_4	W_2	W_3	W_4	W_5	
dim	2	1	1	1	1	
P_3	1	1	0	0	-1	-1
P_4	1	1	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_3T_4	W_2	W_3	W_4	W_5	
	2	1	1	1	1	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

SG 25

Space group $Pmm2$, No. 25, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5	Y_5
dim		2	2
Δ_3	1	1	-1
Δ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		Γ_5	Y_5	
		2	2	Bipartite graph
v^1	1	1		$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	X_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		Γ_5	X_5	
		2	2	Bipartite graph
v^1	1	1		$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_5
dim		2	2
Λ_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_5	S_5
	dim	2	2
C_3	1	1	-1
C_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_5	T_5
	dim	2	2
H_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	S_5
dim		2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_5	U_5
dim		2	2
G_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_5	T_5
dim		2	2
B_3	1	1	-1
B_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	U_5
dim		2	2
A_3	1	1	-1
A_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	U_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_5	R_5
dim		2	2
P_3	1	1	-1
P_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_5	R_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_5	R_5
dim		2	2
E_3	1	1	-1
E_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_5	R_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_5	R_5
	dim	2	2
Q_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_5	R_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 26

Space group $Pmc2_1$, No. 26, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5	Y_5
	dim	2	2
Δ_3	1	1	-1
Δ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	X_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	$Z_5 Z_5$
dim		2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	$Z_5 Z_5$	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_5	S_5
dim		2	2
C_3	1	1	-1
C_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_5	T_5T_5
dim	2	4
H_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

Y_5	T_5T_5	
2	4	Bipartite graph
v^1	2	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_5	S_5
dim	2	2
D_3	1	-1
D_4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

X_5	S_5	
2	2	Bipartite graph
v^1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_5	U_5U_5
dim	2	4
G_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

X_5	U_5U_5	
2	4	Bipartite graph
v^1	2	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_5Z_5	T_5T_5
dim	4	4
B_3B_3	2	1
B_4B_4	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	T_5T_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	U_5U_5
dim	4	4
A_3A_4	2	2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	U_5U_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_5U_5	R_5R_5
dim	4	4
P_3P_3	2	1
P_4P_4	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_5U_5	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_5T_5	R_5R_5
dim	4	4
E_3E_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_5T_5	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_5	R_5R_5
dim	2	4
Q_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_5	R_5R_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

SG 27

Space group $Pcc2$, No. 27, with SOC

 Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_5	Y_5
dim	2	2
Δ_3	1	-1
Δ_4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	$Z_5 Z_5$
	dim	2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	$Z_5 Z_5$	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_5	S_5
	dim	2	2
C_3	1	1	-1
C_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_5	T_5T_5
	dim	2	4
H_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5T_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_5	S_5
	dim	2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_5	U_5U_5
dim	2	4
G_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_5U_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_5Z_5	T_5T_5
dim	4	4
B_3B_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	T_5T_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	U_5U_5
dim	4	4
A_3A_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	U_5U_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_5U_5	R_5R_5
dim	4	4
P_3P_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_5U_5	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_5T_5	R_5R_5
dim	4	4
E_3E_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_5T_5	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_5	R_5R_5
dim	2	4
Q_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_5	R_5R_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

SG 28

Space group $Pma2$, No. 28, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
	dim	2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_5
	dim	2	2
Λ_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_2Y_3	Y_4Y_5	S_2S_3	S_4S_5
dim	2	2	2	2
C_3C_4	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_2Y_3	Y_4Y_5	S_2S_3	S_4S_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5
dim	2	2	2	2
H_2	1	1	0	-1 0
H_3	1	0	1	0 -1
H_4	1	1	0	-1 0
H_5	1	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	S_2S_3	S_4S_5
	dim	2	2	2
D_3	1	1	-2	0
D_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_2S_3	S_4S_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	U_5
	dim	2	2
G_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_5	T_2T_3	T_4T_5
	dim	2	2	2
B_3	1	1	-2	0
B_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_2T_3	T_4T_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	U_5
dim		2	2
A_3	1	1	-1
A_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	U_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_5	R_2R_3	R_4R_5
dim		2	2	2
P_3	1	1	-2	0
P_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_5	R_2R_3	R_4R_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_2T_3	T_4T_5	R_2R_3	R_4R_5
dim		2	2	2	2
E_3E_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_2T_3	T_4T_5	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_2S_3	S_4S_5	R_2R_3	R_4R_5
	dim	2	2	2	2
Q_2	1	1	0	-1	0
Q_3	1	0	1	0	-1
Q_4	1	1	0	-1	0
Q_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_2S_3	S_4S_5	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 29

Space group $Pca2_1$, No. 29, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
	dim	2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_5Z_5
	dim	2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5Z_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_2Y_3	Y_4Y_5	S_2S_3	S_4S_5
dim	2	2		2	2
C_3C_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_2Y_3	Y_4Y_5	S_2S_3	S_4S_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_2Y_3	Y_4Y_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5
dim	2	2		2	2	2	2
H_2	1	1	0	0	0	-2	0
H_3	1	0	1	0	-2	0	0
H_4	1	1	0	0	0	0	-2
H_5	1	0	1	-2	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_2Y_3	Y_4Y_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	2	1	1	0	0	$H_{4/4}^2$
v^2	2	0	0	0	1	1	$H_{4/4}^2$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	S_2S_3	S_4S_5
dim	2	2	2	
D_3	1	1	-2	0
D_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_2S_3	S_4S_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_5	U_5U_5
dim	2	4
G_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_5U_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_5Z_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5
dim	4	2	2	2	2
B_3B_3	2	1	0	0	-1
B_4B_4	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5Z_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	U_5U_5
dim	4	4
A_3A_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	U_5U_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_5U_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	
dim	4	2	2	2	2	
P_3P_3	2	1	0	0	-1	-1
P_4P_4	2	1	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_5U_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_2T_2	T_3T_3	T_4T_4	T_5T_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5
dim	2	2	2	2	2	2	2	2
E_3E_3	2	1	0	1	0	-1	0	-1
E_4E_4	2	0	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	T_2T_2	T_3T_3	T_4T_4	T_5T_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_2S_3	S_4S_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5
	dim	2	2	2	2	2	2
Q_2	1	1	0	0	0	-2	0
Q_3	1	0	1	0	-2	0	0
Q_4	1	1	0	0	0	0	-2
Q_5	1	0	1	-2	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_2S_3	S_4S_5	R_2R_2	R_3R_3	R_4R_4	R_5R_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	2	1	1	0	0	$H_{4/4}^2$
v^2	2	0	0	0	1	1	$H_{4/4}^2$

SG 30

Space group $Pnc2$, No. 30, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
	dim	2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_5Z_5
	dim	2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5Z_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_2Y_3	Y_4Y_5	S_2S_3	S_4S_5
dim	2	2	2	2
C_3C_4	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_2Y_3	Y_4Y_5	S_2S_3	S_4S_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5
dim	2	2	2	2
H_2	1	1	0	0 -1
H_3	1	0	1	-1 0
H_4	1	1	0	-1 0
H_5	1	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_5	S_2S_3	S_4S_5
dim	2	2	2
D_3	1	1	-2 0
D_4	1	1	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_2S_3	S_4S_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_5	U_5U_5
dim	2	4
G_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_5U_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_5Z_5	T_2T_3	T_4T_5
dim	4	2	2
B_3B_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5Z_5	T_2T_3	T_4T_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_5Z_5	U_5U_5
dim	4	4
A_3A_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	U_5U_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_5U_5	R_2R_3	R_4R_5
dim	4	2	2
P_3P_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_5U_5	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_2T_3	T_4T_5	R_2R_3	R_4R_5
dim	2	2	2	2
E_3	1	0	2	0
E_4	1	2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_2T_3	T_4T_5	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_2S_3	S_4S_5	R_2R_3	R_4R_5
dim		2	2	2	2
Q_2	1	1	0	0	-1
Q_3	1	0	1	-1	0
Q_4	1	1	0	-1	0
Q_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_2S_3	S_4S_5	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 31

Space group $Pmn2_1$, No. 31, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
dim		2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	X_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		Γ_5	X_5	
		2	2	Bipartite graph
v^1	1	1		$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	$Z_5 Z_5$
dim		2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		Γ_5	$Z_5 Z_5$	
		2	4	Bipartite graph
v^1	2	1		$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		$Y_2 Y_3$	$Y_4 Y_5$	$S_2 S_3$	$S_4 S_5$
dim		2	2	2	2
$C_3 C_4$	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

		$Y_2 Y_3$	$Y_4 Y_5$	$S_2 S_3$	$S_4 S_5$	
		2	2	2	2	Bipartite graph
v^1	0	1	0	1		$H_{2/1}^1$
v^2	0	1	1	0		$H_{2/1}^1$
v^3	1	0	0	1		$H_{2/1}^1$
v^4	1	0	1	0		$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5
	dim	2	2	2	2
H_2	1	1	0	-1	0
H_3	1	0	1	-1	0
H_4	1	1	0	0	-1
H_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	S_2S_3	S_4S_5
	dim	2	2	2
D_3	1	1	-2	0
D_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	S_2S_3	S_4S_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	U_5U_5
	dim	2	4
G_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	U_5U_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_5Z_5	T_2T_3	T_4T_5
dim	4	2	2
B_3B_4	2	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5Z_5	T_2T_3	T_4T_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	U_5U_5
dim	4	4
A_3A_3	2	1 -1
A_4A_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	U_5U_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_5U_5	R_2R_3	R_4R_5
	dim	4	2	2
P_3P_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_5U_5	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_2T_3	T_4T_5	R_2R_3	R_4R_5
	dim	2	2	2	2
E_3E_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_2T_3	T_4T_5	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_2S_3	S_4S_5	R_2R_3	R_4R_5
	dim	2	2	2	2
Q_2	1	1	0	-1	0
Q_3	1	0	1	-1	0
Q_4	1	1	0	0	-1
Q_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_2S_3	S_4S_5	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 32

Space group $Pba2$, No. 32, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
dim		2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	X_2X_3	X_4X_5
dim		2	2	2
Σ_3	1	1	0	-2
Σ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_2X_3	X_4X_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_5
	dim	2	2
Λ_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_2Y_3	Y_4Y_5	S_5S_5
	dim	2	2	4
C_3C_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_2Y_3	Y_4Y_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H : $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5
	dim	2	2	2	2
H_2	1	1	0	-1	0
H_3	1	0	1	0	-1
H_4	1	1	0	-1	0
H_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_2X_3	X_4X_5	S_5S_5
dim	2	2	4
D_3D_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_2X_3	X_4X_5	U_2U_3	U_4U_5
dim	2	2	2	2
G_2	1	0	1	0
G_3	1	1	0	-1
G_4	1	1	0	0
G_5	1	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_2X_3	X_4X_5	U_2U_3	U_4U_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_5	T_2T_3	T_4T_5
dim		2	2	2
B_3	1	1	-2	0
B_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_2T_3	T_4T_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	U_2U_3	U_4U_5
dim		2	2	2
A_3	1	1	0	-2
A_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	U_2U_3	U_4U_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_2U_3	U_4U_5	R_5R_5
dim		2	2	4
P_3P_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_2U_3	U_4U_5	R_5R_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_2T_3	T_4T_5	R_5R_5
dim	2	2	4
E_3E_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	T_2T_3	T_4T_5	R_5R_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_5S_5	R_5R_5
dim	4	4
Q_5	2	2
		-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_5S_5	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 33

Space group $Pna2_1$, No. 33, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
	dim	2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	X_2X_3	X_4X_5
	dim	2	2	2
Σ_3	1	1	0	-2
Σ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_2X_3	X_4X_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	Z_5Z_5
	dim	2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5Z_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_2Y_3	Y_4Y_5	S_5S_5
dim	2	2	4
C_3C_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_2Y_3	Y_4Y_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_2Y_3	Y_4Y_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5
dim	2	2	2	2	2	2
H_2	1	1	0	0	-2	0
H_3	1	0	1	0	-2	0
H_4	1	1	0	0	0	-2
H_5	1	0	1	-2	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_2Y_3	Y_4Y_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	2	1	1	0	0	$H_{4/4}^2$
v^2	2	0	0	0	1	1	$H_{4/4}^2$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_2X_3	X_4X_5	S_5S_5
dim	2	2	4
D_3D_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_2X_3	X_4X_5	U_2U_3	U_4U_5
	dim	2	2	2	2
G_2	1	0	1	0	-1
G_3	1	1	0	0	-1
G_4	1	1	0	-1	0
G_5	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_2X_3	X_4X_5	U_2U_3	U_4U_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_5Z_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5
	dim	4	2	2	2	2
B_3B_3	2	1	0	0	-1	-1
B_4B_4	2	1	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5Z_5	T_2T_2	T_3T_3	T_4T_4	T_5T_5	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	U_2U_3	U_4U_5
dim	4	2	2
A_3A_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5Z_5	U_2U_3	U_4U_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_2U_3	U_4U_5	R_5R_5
dim	2	2	4
P_3P_4	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_2U_3	U_4U_5	R_5R_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_2T_2	T_3T_3	T_4T_4	T_5T_5	R_5R_5
dim	2	2	2	2	4
E_3E_3	2	1	0	1	0
E_4E_4	2	0	1	0	1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_2T_2	T_3T_3	T_4T_4	T_5T_5	R_5R_5	
	2	2	2	2	4	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_5S_5	R_5R_5
	dim	4	4
Q_5	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_5S_5	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 34

Space group $Pnn2$, No. 34, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
	dim	2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_5	X_2X_3	X_4X_5
dim	2	2	2
Σ_3	1	1	-2
Σ_4	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_2X_3	X_4X_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5	Z_5Z_5
dim	2	4
Λ_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5Z_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_2Y_3	Y_4Y_5	S_5S_5
dim	2	2	4
C_3C_4	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_2Y_3	Y_4Y_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5
	dim	2	2	2	2
H_2	1	1	0	0	-1
H_3	1	0	1	-1	0
H_4	1	1	0	-1	0
H_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_2Y_3	Y_4Y_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_2X_3	X_4X_5	S_5S_5
	dim	2	2	4
D_3D_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	S_5S_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_2X_3	X_4X_5	U_2U_3	U_4U_5
	dim	2	2	2	2
G_2	1	0	1	-1	0
G_3	1	1	0	0	-1
G_4	1	1	0	-1	0
G_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_2X_3	X_4X_5	U_2U_3	U_4U_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_5Z_5	T_2T_3	T_4T_5
	dim	4	2	2
B_3B_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5Z_5	T_2T_3	T_4T_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_5	U_2U_3	U_4U_5
	dim	4	2	2
A_3A_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5Z_5	U_2U_3	U_4U_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_2U_3	U_4U_5	R_5
	dim	2	2	2
P_3	1	0	2	-1
P_4	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_2U_3	U_4U_5	R_5	
	2	2	2	Bipartite graph
v^1	1	1	2	$H_{4/4}^2$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_2T_3	T_4T_5	R_5
	dim	2	2	2
E_3	1	0	2	-1
E_4	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_2T_3	T_4T_5	R_5	
	2	2	2	Bipartite graph
v^1	1	1	2	$H_{4/4}^2$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$S_5 S_5$	R_5
	dim	4	2
Q_5	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$S_5 S_5$	R_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/2}^1$

SG 35

Space group $Cmm2$, No. 35, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_5
	dim	2	2
Λ_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	Y_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_5	T_5
dim		2	2
H_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	T_5
dim		2	2
A_3	1	1	-1
A_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_3R_4
dim		2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 36

Space group $Cmc2_1$, No. 36, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5	$Z_5 Z_5$
dim	2	4
Λ_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

Γ_5	$Z_5 Z_5$	
2	4	Bipartite graph
v^1	2	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_5	Y_5
dim	2	2
Σ_3	1	-1
Σ_4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

Γ_5	Y_5	
2	2	Bipartite graph
v^1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Y_5	$T_5 T_5$
dim	2	4
H_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5T_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_5	T_5T_5
dim		4	4
A_3A_3	2	1	-1
A_4A_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	T_5T_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_3R_3	R_4R_4
dim		2	2	2
D_3	1	1	-2	0
D_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_3	R_4R_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

SG 37

Space group $Ccc2$, No. 37, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	$Z_5 Z_5$
dim		2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	$Z_5 Z_5$	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	Y_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_5	$T_5 T_5$
dim		2	4
H_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	$T_5 T_5$	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_5	T_5T_5
dim		4	4
A_3A_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5Z_5	T_5T_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_3R_4
dim		2	2
D_3	1	1	-1
D_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 38

Space group $Amm2$, No. 38, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	Z_5
dim		2	2
Λ_3	1	1	-1
Λ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	Y_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_5	T_5
	dim	2	2
H_3	1	1	-1
H_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A : $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	T_5
dim		2	2
A_3	1	1	-1
A_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_3S_4	R_3R_4
dim		2	2
D_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 39

Space group $Abm2$, No. 39, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_5
dim		2	2
Λ_3	1	1	-1
Λ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	Y_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_5	T_5
	dim	2	2
H_3	1	1	-1
H_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A : $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	T_5
	dim	2	2
A_3	1	1	-1
A_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	T_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_3	S_4S_4	R_3R_3	R_4R_4
dim		2	2	2	2
D_2D_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_3S_3	S_4S_4	R_3R_3	R_4R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 40

Space group $Ama2$, No. 40, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	Z_2Z_3	Z_4Z_5
dim		2	2	2
Λ_3	1	1	0	-2
Λ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_2Z_3	Z_4Z_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	Y_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_5	T_2T_3	T_4T_5
dim		2	2	2
H_3	1	1	0	-2
H_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_2T_3	T_4T_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A : $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5
dim		2	2	2	2
A_3A_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_3S_4	R_3R_4
	dim	2	2
D_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 41

Space group *Aba2*, No. 41, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5	Z_2Z_3	Z_4Z_5
	dim	2	2	2
Λ_3	1	1	0	-2
Λ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_2Z_3	Z_4Z_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	Y_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_5	T_2T_3	T_4T_5
dim		2	2	2
H_3	1	1	-2	0
H_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_5	T_2T_3	T_4T_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A : $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5
dim		2	2	2	2
A_3A_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_2Z_3	Z_4Z_5	T_2T_3	T_4T_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D : $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_3S_3	S_4S_4	R_3R_3	R_4R_4
dim	2	2	2	2
D_2D_2	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_3S_3	S_4S_4	R_3R_3	R_4R_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 42

Space group $Fmm2$, No. 42, with SOC

Λ

High symmetry line $\Lambda(u, u, 0)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_5	Z_5
dim	2	2
Λ_5	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Σ High symmetry line $\Sigma(u, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$ **A** matrix

corep		Γ_5	X_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

 $\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

Γ_5	X_5	
2	2	Bipartite graph
v^1	1	$H_{2/2}^1$

 Δ High symmetry line $\Delta(0, -u, -u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(0, -\frac{1}{2}, -\frac{1}{2})$ **A** matrix

corep		Γ_5	Y_5
dim		2	2
Δ_3	1	1	-1
Δ_4	1	1	-1

 $\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

Γ_5	Y_5	
2	2	Bipartite graph
v^1	1	$H_{2/2}^1$

SG 43Space group *Fdd2*, No. 43, with SOC Λ High symmetry line $\Lambda(u, u, 0)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(\frac{1}{2}, \frac{1}{2}, 0)$ **A** matrix

corep		Γ_5	$Z_5 Z_5$
dim		2	4
Λ_5	2	1	-2

 $\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Z_5Z_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_5	X_2X_3	X_4X_5
dim		2	2	2
Σ_3	1	1	0	-2
Σ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_2X_3	X_4X_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Δ

High symmetry line $\Delta(0, -u, -u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(0, -\frac{1}{2}, -\frac{1}{2})$

A matrix

corep		Γ_5	Y_2Y_3	Y_4Y_5
dim		2	2	2
Δ_3	1	1	-2	0
Δ_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	Y_2Y_3	Y_4Y_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

SG 44

Space group *Imm2*, No. 44, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_5	X_5
dim		2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q : $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep		R_3R_4	W_3	W_4
dim		2	1	1
Q_2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	R_3R_4	W_3	W_4	
	2	1	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D : $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep		S_3S_4	W_3	W_4
dim		2	1	1
D_2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	S_3S_4	W_3	W_4	
	2	1	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P : $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		T_3T_4	W_3	W_4
	dim	2	1	1
P_3	1	1	-1	0
P_4	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	W_3	W_4	
	2	1	1	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 45

Space group $Iba2$, No. 45, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	R_3R_3	R_4R_4	W_3W_3	W_4W_4
dim	2	2	2	2
Q_2Q_2	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_3R_3	R_4R_4	W_3W_3	W_4W_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	S_3S_3	S_4S_4	W_3W_3	W_4W_4
dim	2	2	2	2
D_2D_2	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_3S_3	S_4S_4	W_3W_3	W_4W_4	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		T_3T_4	W_3W_3	W_4W_4
	dim	2	2	2
P_3	1	1	-2	0
P_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	W_3W_3	W_4W_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

SG 46

Space group *Ima2*, No. 46, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_5	X_5
	dim	2	2
Σ_3	1	1	-1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q : $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep		R_3R_4	W_3W_4
	dim	2	2
Q_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	W_3W_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	S_3S_3	S_4S_4	W_3W_4
dim	2	2	2
D_2D_2	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_3S_3	S_4S_4	W_3W_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	T_3T_4	W_3W_4
dim	2	2
P_3	1	1
P_4	1	1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	W_3W_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 47

Space group *Pmmm*, No. 47, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2
Δ_5	2	1	1	-1
				-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	X_5^+	X_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
dim	2	2	2	2
Λ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_5^+	Y_5^-	S_5^+	S_5^-
	dim	2	2	2	2
C_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_5^+	Y_5^-	S_5^+	S_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_5^+	Y_5^-	T_5^+	T_5^-
	dim	2	2	2	2
H_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_5^+	Y_5^-	T_5^+	T_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5^+	X_5^-	S_5^+	S_5^-
	dim	2	2	2	2
D_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_5^+	X_5^-	S_5^+	S_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_5^+	X_5^-	U_5^+	U_5^-
dim	2	2	2	2
G_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_5^+	X_5^-	U_5^+	U_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_5^+	Z_5^-	T_5^+	T_5^-
dim	2	2	2	2
B_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5^+	Z_5^-	T_5^+	T_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5^+	Z_5^-	U_5^+	U_5^-
	dim	2	2	2	2
A_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5^+	Z_5^-	U_5^+	U_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_5^+	U_5^-	R_5^+	R_5^-
	dim	2	2	2	2
P_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_5^+	U_5^-	R_5^+	R_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_5^+	T_5^-	R_5^+	R_5^-
	dim	2	2	2	2
E_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_5^+	T_5^-	R_5^+	R_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_5^+	S_5^-	R_5^+	R_5^-
dim	2	2	2	2
Q_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_5^+	S_5^-	R_5^+	R_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 48

Space group $Pnnn$, No. 48, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	Y_3Y_4
dim	2	2	4
Δ_5	2	1	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	X_3X_4
dim	2	2	4
Σ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_3Y_4	S_3S_4
dim	4	4
C_2C_5	2	-1
C_3C_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_3Y_4	T_3T_4
dim	4	4
H_2H_5	2	-1
H_3H_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	S_3S_4
dim	4	4
D_2D_5	2	-1
D_3D_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	U_3U_4
dim		4	4
G_2G_5	2	1	-1
G_3G_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_3Z_4	T_3T_4
dim		4	4
B_2B_5	2	1	-1
B_3B_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_3Z_4	U_3U_4
dim		4	4
A_2A_4	2	1	-1
A_3A_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_3U_4	R_5^+	R_5^-
dim		4	2	2
P_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	U_3U_4	R_5^+	R_5^-	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_3T_4	R_5^+	R_5^-
dim		4	2	2
E_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	T_3T_4	R_5^+	R_5^-	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_5^+	R_5^-
	dim	4	2	2
Q_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	S_3S_4	R_5^+	R_5^-	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

SG 49

Space group $Pccm$, No. 49, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
	dim	2	2	2	2
Δ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	X_5^+	X_5^-
	dim	2	2	2	2
Σ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	Z_3Z_4
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_5^+	Y_5^-	S_5^+	S_5^-
	dim	2	2	2	2
C_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_5^+	Y_5^-	S_5^+	S_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_5^+	Y_5^-	T_3T_4	
dim	2	2	4	
H_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_5^+	X_5^-	S_5^+	S_5^-	
dim	2	2	2	2	
D_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_5^+	X_5^-	S_5^+	S_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5^+	X_5^-	U_3U_4
dim		2	2	4
G_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_5^+	X_5^-	U_3U_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_3Z_4	T_3T_4
dim		4	4
B_2B_5	2	1	-1
B_3B_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_3Z_4	U_3U_4
dim		4	4
A_2A_4	2	1	-1
A_3A_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_3U_4	R_3R_4
dim		4	4
P_2P_5	2	1	-1
P_3P_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_3U_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_3T_4	R_3R_4
dim		4	4
E_2E_4	2	1	-1
E_3E_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_5^+	S_5^-	R_3R_4
	dim	2	2	4
Q_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	S_5^+	S_5^-	R_3R_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 50

Space group *Pban*, No. 50, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	Y_3Y_4
	dim	2	2	4
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	X_3X_4
	dim	2	2	4
Σ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
dim	2	2	2	2
Λ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_3Y_4	S_3S_4
dim	4	4
C_2C_5	2	1 -1
C_3C_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_3Y_4	T_3T_4
dim	4	4
H_2H_5	2	-1
H_3H_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	S_3S_4
dim	4	4
D_2D_5	2	-1
D_3D_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	U_3U_4
dim	4	4
G_2G_5	2	-1
G_3G_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_5^+	Z_5^-	T_3T_4
dim	2	2	4
B_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5^+	Z_5^-	U_3U_4
dim	2	2	4
A_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	U_3U_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_3U_4	R_3R_4
dim		4	4
P_2P_5	2	1	-1
P_3P_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_3U_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_3T_4	R_3R_4
dim		4	4
E_2E_5	2	1	-1
E_3E_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_3R_4
dim		4	4
Q_5	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 51

Space group $Pmma$, No. 51, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2
Δ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	X_5^+	X_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	Z_3Z_4
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_5^+	Y_5^-	S_5^+	S_5^-
	dim	2	2	2	2
C_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_5^+	Y_5^-	S_5^+	S_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_5^+	Y_5^-	T_3T_4
	dim	2	2	4
H_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_5^+	X_5^-	S_5^+	S_5^-
dim	2	2	2	2
D_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_5^+	X_5^-	S_5^+	S_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_5^+	X_5^-	U_3U_4
dim	2	2	4
G_5	2	1	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_5^+	X_5^-	U_3U_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_3Z_4	T_3T_4
dim	4	4
B_2B_3	2	1
B_4B_5	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	U_3U_4
dim	4	4
A_5A_5	4	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_3U_4	R_3R_4
dim	4	4
P_2P_3	2	1
P_4P_5	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_3U_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of *E*: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_3T_4	R_3R_4
dim	4	4
E_5E_5	4	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of *Q*: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_5^+	S_5^-	R_3R_4
dim	2	2	4
Q_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	S_5^+	S_5^-	R_3R_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 52

Space group *Pnna*, No. 52, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	Y_3Y_4
dim	2	2	4
Δ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	X_3X_4
dim	2	2	4
Σ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_3Y_4	S_3S_4
dim	4	4
C_2C_5	2	1
C_3C_4	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_3Y_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$
dim	4	2	2	2	2
H_2H_5	2	1	-1	0	-1
H_3H_4	2	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_3Y_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	S_3S_4
dim	4	4
D_2D_5	2	1
D_3D_4	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_3X_4	U_3U_3	U_4U_4
dim		4	4	4
G_2G_5	2	1	0	-2
G_3G_4	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_3Z_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$
dim		4	2	2	2	2
B_2B_3	2	1	-1	-1	0	0
B_4B_5	2	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_3Z_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	U_3U_3	U_4U_4
dim	4	4	4
A_5A_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_3Z_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_3U_3	U_4U_4	R_3R_4
dim	4	4	4
P_5P_5	4	1	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_3U_3	U_4U_4	R_3R_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	R_3R_4
dim	2	2	2	2	4
E_2E_4	2	0	0	1	1 -1
E_3E_5	2	1	1	0	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	R_3R_4	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		S_3S_4	R_3R_4
	dim	4	4
Q_5	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 53

Space group $Pmna$, No. 53, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep		Γ_5^+	Γ_5^-	Y_3Y_4
	dim	2	2	4
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	X_5^+	X_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_3Y_4	S_3S_4
dim	4	4
C_2C_5	2	1
		-1
C_3C_4	2	1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_3Y_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$
dim	4	2	2	2	2
H_2H_5	2	1	0	0	-1
H_3H_4	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_3Y_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_5^+	X_5^-	S_3S_4
dim	2	2	4
D_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_5^+	X_5^-	S_3S_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5^+	X_5^-	U_3U_4
dim		2	2	4
G_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_5^+	X_5^-	U_3U_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_3Z_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$
dim		4	2	2	2	2
B_2B_3	2	1	0	-1	-1	0
B_4B_5	2	1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_3Z_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	Bipartite graph
	4	2	2	2	2	
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_3Z_4	U_3U_4
dim		4	4
A_5A_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		U_3U_4	$R_2^+R_3^+$	$R_2^-R_3^-$	$R_4^+R_5^+$	$R_4^-R_5^-$
dim		4	2	2	2	2
P_2P_3	2	1	0	-1	-1	0
P_4P_5	2	1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_3U_4	$R_2^+R_3^+$	$R_2^-R_3^-$	$R_4^+R_5^+$	$R_4^-R_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	$R_2^+R_3^+$	$R_2^-R_3^-$	$R_4^+R_5^+$	$R_4^-R_5^-$
dim		2	2	2	2	2	2	2	2
E_2E_5	2	0	1	1	0	0	-1	-1	0
E_3E_4	2	1	0	0	1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	$R_2^+R_3^+$	$R_2^-R_3^-$	$R_4^+R_5^+$	$R_4^-R_5^-$	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	0	0	1	0	0	$H_{2/1}^1$

v^7	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_3S_4	$R_2^+R_3^+$	$R_2^-R_3^-$	$R_4^+R_5^+$	$R_4^-R_5^-$
dim	4	2	2	2	2
Q_2Q_5	2	1	0	0	-1
Q_3Q_4	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_3S_4	$R_2^+R_3^+$	$R_2^-R_3^-$	$R_4^+R_5^+$	$R_4^-R_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

SG 54

Space group $Pcca$, No. 54, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2
Δ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	X_3X_4
	dim	2	2	4
Σ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5^+	Γ_5^-	Z_3Z_4
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_5^+	Y_5^-	S_3S_4
	dim	2	2	4
C_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	S_3S_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_5^+	Y_5^-	T_3T_4
dim	2	2	4
H_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_3X_4	S_3S_4
dim	4	4
D_2D_5	2	1
D_3D_4	2	1
		-1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	U_3U_3	U_4U_4
dim	4	4	4
G_2G_5	2	1	0 -2
G_3G_4	2	1	-2 0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_3Z_4	T_3T_4
dim	4	4
B_2B_3	2	1 -1
B_4B_5	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	U_3U_3	U_4U_4
dim	4	4	4
A_5A_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_3Z_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	U_3U_3	U_4U_4	R_3R_3	R_4R_4
dim	4	4	4	4
P_5P_5	4	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_3U_3	U_4U_4	R_3R_3	R_4R_4	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	T_3T_4	R_3R_3	R_4R_4
dim	4	4	4
E_5E_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_3T_4	R_3R_3	R_4R_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_4	R_3R_3	R_4R_4
	dim	4	4	4
Q_2Q_5	2	1	0	-2
Q_3Q_4	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_3	R_4R_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

SG 55

Space group *Pbam*, No. 55, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	Y_3Y_4
	dim	2	2	4
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	X_3X_4
	dim	2	2	4
Σ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
	dim	2	2	2	2
Λ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_3Y_4	$S_5^+S_5^+$	$S_5^-S_5^-$
	dim	4	4	4
C_5C_5	4	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_3Y_4	$S_5^+S_5^+$	$S_5^-S_5^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_3Y_4	T_3T_4
dim	4	4
H_2H_3	2	-1
H_4H_5	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	$S_5^+S_5^+$	$S_5^-S_5^-$
dim	4	4	4
D_5D_5	4	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	$S_5^+S_5^+$	$S_5^-S_5^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	U_3U_4
	dim	4	4
G_2G_3	2	1	-1
G_4G_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Z_5^+	Z_5^-	T_3T_4
	dim	2	2	4
B_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_5^+	Z_5^-	U_3U_4
	dim	2	2	4
A_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	U_3U_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_3U_4	$R_5^+ R_5^+$	$R_5^- R_5^-$
dim	4	4	4
P_5P_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	U_3U_4	$R_5^+ R_5^+$	$R_5^- R_5^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_3T_4	$R_5^+ R_5^+$	$R_5^- R_5^-$
dim	4	4	4
E_5E_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_3T_4	$R_5^+ R_5^+$	$R_5^- R_5^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$S_5^+ S_5^+$	$S_5^- S_5^-$	$R_5^+ R_5^+$	$R_5^- R_5^-$
dim	4	4	4	4
Q_5Q_5	4	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$S_5^+ S_5^+$	$S_5^- S_5^-$	$R_5^+ R_5^+$	$R_5^- R_5^-$	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 56

Space group $Pccn$, No. 56, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	$Y_3 Y_4$
dim	2	2	4
Δ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	$Y_3 Y_4$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	$X_3 X_4$
dim	2	2	4
Σ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	$X_3 X_4$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_3Y_4	S_3S_4
dim	4	4
C_2C_4	2	1
C_3C_5	2	1
		-1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H : $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_3Y_4	T_3T_3	T_4T_4
dim	4	4	4
H_5H_5	4	1	-1
		-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_3Y_4	T_3T_3	T_4T_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_3X_4	S_3S_4
	dim	4	4
D_2D_5	2	1	-1
D_3D_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	U_3U_3	U_4U_4
	dim	4	4	4
G_5G_5	4	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_3Z_4	T_3T_3	T_4T_4
dim		4	4	4
B_2B_5	2	1	0	-2
B_3B_4	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_3	T_4T_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_3Z_4	U_3U_3	U_4U_4
dim		4	4	4
A_2A_4	2	1	0	-2
A_3A_5	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_3U_3	U_4U_4	$R_5^+R_5^+$	$R_5^-R_5^-$
dim		4	4	4	4
P_5P_5	4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	U_3U_3	U_4U_4	$R_5^+R_5^+$	$R_5^-R_5^-$	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_3T_3	T_4T_4	$R_5^+ R_5^+$	$R_5^- R_5^-$
dim	4	4	4	4
E_5E_5	4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_3T_3	T_4T_4	$R_5^+ R_5^+$	$R_5^- R_5^-$	Bipartite graph
	4	4	4	4	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_3S_4	$R_5^+ R_5^+$	$R_5^- R_5^-$
dim	4	4	4
Q_5Q_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_3S_4	$R_5^+ R_5^+$	$R_5^- R_5^-$	Bipartite graph
	4	4	4	
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

SG 57

Space group *Pbcm*, No. 57, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix



corep		Γ_5^+	Γ_5^-	Y_3Y_4
	dim	2	2	4
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5^+	Γ_5^-	X_3X_4
	dim	2	2	4
Σ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
	dim	2	2	2	2
Λ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_3Y_4	S_3S_3	S_4S_4
dim	4	4	4
C_5C_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_3Y_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_3Y_4	T_3T_4
dim	4	4
H_2H_3	2	1 -1
H_4H_5	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	S_3S_3	S_4S_4
dim	4	4	4
D_2D_5	2	1	-2 0
D_3D_4	2	1	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	U_3U_4
dim	4	4
G_5G_5	4	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_5^+	Z_5^-	T_3T_4
dim	2	2	4
B_5	2	1 1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_5^+	Z_5^-	U_3U_4
	dim	2	2	4
A_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	U_3U_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		U_3U_4	R_3R_3	R_4R_4
	dim	4	4	4
P_2P_5	2	1	-2	0
P_3P_4	2	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_3U_4	R_3R_3	R_4R_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_3T_4	R_3R_3	R_4R_4
dim	4	4	4
E_5E_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_3T_4	R_3R_3	R_4R_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_3S_3	S_4S_4	R_3R_3	R_4R_4
dim	4	4	4	4
Q_2Q_2	2	0	1	0 -1
Q_3Q_3	2	1	0	-1 0
Q_4Q_4	2	0	1	0 -1
Q_5Q_5	2	1	0	-1 0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	S_3S_3	S_4S_4	R_3R_3	R_4R_4	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 58

Space group $Pn\bar{m}$, No. 58, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix



corep		Γ_5^+	Γ_5^-	Y_3Y_4
	dim	2	2	4
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5^+	Γ_5^-	X_3X_4
	dim	2	2	4
Σ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	Z_3Z_4
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Y_3Y_4	$S_5^+S_5^+$	$S_5^-S_5^-$
dim	4	4	4
C_5C_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_3Y_4	$S_5^+S_5^+$	$S_5^-S_5^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Y_3Y_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$
dim	4	2	2	2	2
H_2H_3	2	1	-1	0	0
H_4H_5	2	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_3Y_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	$S_5^+S_5^+$	$S_5^-S_5^-$
dim	4	4	4
D_5D_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	$S_5^+S_5^+$	$S_5^-S_5^-$	Bipartite graph
	4	4	4	
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$
dim	4	2	2	2	2
G_2G_3	2	1	0	0	-1
G_4G_5	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_3X_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	Bipartite graph
	4	2	2	2	2	
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_3Z_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$
dim	4	2	2	2	2
B_2B_5	2	1	0	-1	0
B_3B_4	2	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_3Z_4	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_3Z_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$
dim	4	2	2	2	2
A_2A_4	2	1	0	-1	-1
A_3A_5	2	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_3Z_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	R_3R_4
dim	2	2	2	2	4
P_2P_4	2	1	1	0	0
P_3P_5	2	0	0	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	R_3R_4	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	R_3R_4	
dim	2	2	2	2	4	
E_2E_5	2	0	0	1	1	-1
E_3E_4	2	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$T_2^+T_3^+$	$T_2^-T_3^-$	$T_4^+T_5^+$	$T_4^-T_5^-$	R_3R_4	Bipartite graph
	2	2	2	2	4	
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$S_5^+S_5^+$	$S_5^-S_5^-$	R_3R_4	
dim	4	4	4	
Q_5Q_5	4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$S_5^+S_5^+$	$S_5^-S_5^-$	R_3R_4	Bipartite graph
	4	4	4	
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 59

Space group $Pm\bar{m}n$, No. 59, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix



corep		Γ_5^+	Γ_5^-	Y_3Y_4
	dim	2	2	4
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5^+	Γ_5^-	X_3X_4
	dim	2	2	4
Σ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
	dim	2	2	2	2
Λ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Y_3Y_4	S_3S_4
	dim	4	4
C_2C_4	2	1	-1
C_3C_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Y_3Y_4	T_3T_4
	dim	4	4
H_5H_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_3X_4	S_3S_4
	dim	4	4
D_2D_5	2	1	-1
D_3D_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	U_3U_4
	dim	4	4
G_5G_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_5^+	Z_5^-	T_3T_4
	dim	2	2	4
B_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5^+	Z_5^-	U_3U_4
dim	2	2	4
A_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5^+	Z_5^-	U_3U_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_3U_4	R_3R_4
dim	4	4
P_2P_5	2	1
P_3P_4	2	1
		-1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	U_3U_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_3T_4	R_3R_4
dim	4	4
E_2E_4	2	1
E_3E_5	2	1
		-1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_3T_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_3S_4	R_3R_4	
dim	4	4	
Q_5Q_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_4	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 60

Space group $Pbcn$, No. 60, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	Y_3Y_4	
dim	2	2	4	
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	X_3X_4
dim	2	2	4
Σ_5	2	1 1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1 1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_3Y_4	S_3S_3	S_4S_4
dim	4	4	4
C_2C_4	2	1	-2 0
C_3C_5	2	1	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_3Y_4	T_3T_3	T_4T_4
dim	4	4	4
H_5H_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_3Y_4	T_3T_3	T_4T_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	S_3S_3	S_4S_4
dim	4	4	4
D_5D_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	
dim	4	2	2	2	2	
G_2G_3	2	1	0	0	-1	-1
G_4G_5	2	1	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_3X_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	Bipartite graph
	4	2	2	2	2	
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Z_3Z_4	T_3T_3	T_4T_4	
dim	4	4	4	
B_2B_5	2	1	0	-2
B_3B_4	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_3	T_4T_4	Bipartite graph
	4	4	4	
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_3Z_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	
dim	4	2	2	2	2	
A_2A_4	2	1	0	-1	-1	0
A_3A_5	2	1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_3Z_4	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	R_3R_3	R_4R_4
dim		2	2	2	2	4	4
P_2P_4	2	1	1	0	0	-2	0
P_3P_5	2	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$U_2^+U_3^+$	$U_2^-U_3^-$	$U_4^+U_5^+$	$U_4^-U_5^-$	R_3R_3	R_4R_4	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		T_3T_3	T_4T_4	R_3R_3	R_4R_4
dim		4	4	4	4
E_5E_5	4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

T_3T_3	T_4T_4	R_3R_3	R_4R_4	
4	4	4	4	Bipartite graph

v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		S_3S_3	S_4S_4	R_3R_3	R_4R_4
	dim	4	4	4	4
Q_2Q_2	2	0	1	0	-1
Q_3Q_3	2	1	0	0	-1
Q_4Q_4	2	0	1	-1	0
Q_5Q_5	2	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_3	S_4S_4	R_3R_3	R_4R_4	
	4	4	4	4	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 61

Space group $Pbca$, No. 61, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	Y_3Y_4
	dim	2	2	4
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	X_3X_4
dim	2	2	4
Σ_5	2	1 1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1 1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C : $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Y_3Y_4	S_3S_3	S_4S_4
dim	4	4	4
C_2C_3	2	1	0 -2
C_4C_5	2	1	-2 0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Y_3Y_4	T_3T_3	T_4T_4
dim	4	4	4
H_5H_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_3Y_4	T_3T_3	T_4T_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	S_3S_3	S_4S_4
dim	4	4	4
D_5D_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	U_3U_3	U_4U_4
	dim	4	4	4
G_2G_3	2	1	-2	0
G_4G_5	2	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Z_3Z_4	T_3T_3	T_4T_4
	dim	4	4	4
B_2B_3	2	1	0	-2
B_4B_5	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_3	T_4T_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_3Z_4	U_3U_3	U_4U_4
	dim	4	4	4
A_5A_5	4	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_3Z_4	U_3U_3	U_4U_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	U_3U_3	U_4U_4	$R_2^+R_2^+$	$R_2^-R_2^-$	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_5^+$	$R_5^-R_5^-$
dim	4	4	2	2	2	2	2	2	2	2
P_2P_2	2	0	1	0	-1	0	0	-1	0	0
P_3P_3	2	1	0	-1	0	0	0	0	-1	0
P_4P_4	2	0	1	0	0	0	-1	0	0	-1
P_5P_5	2	1	0	0	0	-1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	U_3U_3	U_4U_4	$R_2^+R_2^+$	$R_2^-R_2^-$	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_5^+$	$R_5^-R_5^-$	Bipartite graph
	4	4	2	2	2	2	2	2	2	2	
v^1	0	1	0	0	0	0	1	0	1	0	$H_{3/2}^1$
v^2	0	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^4	0	1	0	1	0	1	0	0	0	0	$H_{3/2}^1$
v^5	1	0	0	0	0	0	0	1	0	1	$H_{3/2}^1$
v^6	1	0	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	1	0	0	0	0	0	$H_{3/2}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	T_3T_3	T_4T_4	$R_2^+R_2^+$	$R_2^-R_2^-$	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_5^+$	$R_5^-R_5^-$
dim	4	4	2	2	2	2	2	2	2	2
E_2E_2	2	1	0	0	0	0	0	0	-1	-1
E_3E_3	2	0	1	0	0	0	0	-1	0	0
E_4E_4	2	1	0	-1	0	0	-1	0	0	0
E_5E_5	2	0	1	0	-1	-1	0	0	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	T_3T_3	T_4T_4	$R_2^+R_2^+$	$R_2^-R_2^-$	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_5^+$	$R_5^-R_5^-$	Bipartite graph
	4	4	2	2	2	2	2	2	2	2	
v^1	0	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^2	0	1	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^3	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$

v^4	0	1	0	1	0	0	1	0	0	0	$H_{3/2}^1$
v^5	1	0	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^6	1	0	0	0	0	1	0	1	0	0	$H_{3/2}^1$
v^7	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	1	0	0	0	0	1	0	0	$H_{3/2}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q : $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_3S_3	S_4S_4	$R_2^+R_2^+$	$R_2^-R_2^-$	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_5^+$	$R_5^-R_5^-$
dim	4	4	2	2	2	2	2	2	2	2
Q_2Q_2	2	1	0	-1	0	0	0	0	0	-1
Q_3Q_3	2	0	1	0	-1	0	0	0	-1	0
Q_4Q_4	2	1	0	0	0	-1	-1	0	0	0
Q_5Q_5	2	0	1	0	0	-1	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	S_3S_3	S_4S_4	$R_2^+R_2^+$	$R_2^-R_2^-$	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_5^+$	$R_5^-R_5^-$	Bipartite graph
	4	4	2	2	2	2	2	2	2	2	
v^1	0	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^2	0	1	0	0	1	0	0	0	1	0	$H_{3/2}^1$
v^3	0	1	0	1	0	0	0	1	0	0	$H_{3/2}^1$
v^4	0	1	0	1	1	0	0	0	0	0	$H_{3/2}^1$
v^5	1	0	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	0	0	0	1	0	0	0	1	$H_{3/2}^1$
v^7	1	0	1	0	0	0	1	0	0	0	$H_{3/2}^1$
v^8	1	0	1	0	0	1	0	0	0	0	$H_{3/2}^1$

SG 62

Space group $Pnma$, No. 62, with SOC

Δ

High symmetry line $\Delta(-u, 0, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(-\frac{1}{2}, 0, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	Y_3Y_4
dim	2	2	4
Δ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	X_3X_4
dim	2	2	4
Σ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

C

High symmetry line $C(-\frac{1}{2}, u, 0)$, high symmetry points at both ends of C: $Y(-\frac{1}{2}, 0, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Y_3Y_4	S_3S_3	S_4S_4
dim		4	4	4
C_5C_5	4	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Y_3Y_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

H

High symmetry line $H(-\frac{1}{2}, 0, u)$, high symmetry points at both ends of H: $Y(-\frac{1}{2}, 0, 0), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Y_3Y_4	T_3T_4
dim		4	4
H_2H_5	2	1	-1
H_3H_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Y_3Y_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(-u, \frac{1}{2}, 0)$, high symmetry points at both ends of D: $X(0, \frac{1}{2}, 0), S(-\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_3X_4	S_3S_3	S_4S_4
dim		4	4	4
D_2D_5	2	1	0	-2
D_3D_4	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	S_3S_3	S_4S_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

G

High symmetry line $G(0, \frac{1}{2}, u)$, high symmetry points at both ends of G: $X(0, \frac{1}{2}, 0), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	$U_5^+U_5^+$	$U_5^-U_5^-$
dim	4	4	4
G_5G_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	$U_5^+U_5^+$	$U_5^-U_5^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

B

High symmetry line $B(-u, 0, \frac{1}{2})$, high symmetry points at both ends of B: $Z(0, 0, \frac{1}{2}), T(-\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Z_3Z_4	T_3T_4
dim	4	4
B_2B_3	2	1 -1
B_4B_5	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

A

High symmetry line $A(0, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), U(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	$U_5^+U_5^+$	$U_5^-U_5^-$
dim	4	4	4
A_5A_5	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_3Z_4	$U_5^+U_5^+$	$U_5^-U_5^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of P: $U(0, \frac{1}{2}, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$U_5^+U_5^+$	$U_5^-U_5^-$	R_3R_3	R_4R_4
dim		4	4	4	4
P_5P_5	4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$U_5^+U_5^+$	$U_5^-U_5^-$	R_3R_3	R_4R_4	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

E

High symmetry line $E(-\frac{1}{2}, u, \frac{1}{2})$, high symmetry points at both ends of E: $T(-\frac{1}{2}, 0, \frac{1}{2}), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		T_3T_4	R_3R_3	R_4R_4
dim		4	4	4
E_5E_5	4	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	T_3T_4	R_3R_3	R_4R_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(-\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of Q: $S(-\frac{1}{2}, \frac{1}{2}, 0), R(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_3S_3	S_4S_4	R_3R_3	R_4R_4
dim	4	4	4	4
Q_2Q_2	2	0	1	-1
Q_3Q_3	2	1	0	-1
Q_4Q_4	2	0	1	0
Q_5Q_5	2	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_3S_3	S_4S_4	R_3R_3	R_4R_4	
	4	4	4	4	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 63

Space group *Cmcm*, No. 63, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H: $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Y_5^+	Y_5^-	T_3T_4
dim	2	2	4
H_5	2	1	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	T_3T_4
dim	4	4
A_2A_3	2	1 -1
A_4A_5	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_2 R_2$
dim	2	2	4
$D_3 D_4$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_2 R_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 64

Space group *Cmca*, No. 64, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	$Z_3 Z_4$
dim	2	2	4
Λ_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	$Z_3 Z_4$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Y_5^+	Y_5^-	T_3T_4
dim	2	2	4
H_5	2	1	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	T_3T_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A : $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	T_3T_4
dim	4	4
A_2A_3	2	1 -1
A_4A_5	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_2S_2	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$
dim	4	2	2	2	2
D_3D_3	2	1	0	-1	-1
D_4D_4	2	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_2S_2	$R_3^+R_3^+$	$R_3^-R_3^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

SG 65

Space group $Cmmm$, No. 65, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
dim	2	2	2	2
Λ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim		2	2	2	2
Σ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Y_5^+	Y_5^-	T_5^+	T_5^-
dim		2	2	2	2
H_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_5^+	Y_5^-	T_5^+	T_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A : $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5^+	Z_5^-	T_5^+	T_5^-
dim		2	2	2	2
A_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5^+	Z_5^-	T_5^+	T_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$
dim	2	2	2	2
$D_3 D_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 66

Space group *Cccm*, No. 66, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$Z_3 Z_4$
dim	2	2	4
Λ_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Y_5^+	Y_5^-	T_3T_4
dim	2	2	4
H_5	2	1	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	T_3T_4
dim	4	4
A_2A_4	2	-1
A_3A_5	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$
dim	2	2	2	2
D_3D_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$S_3^+ S_4^+$	$S_3^- S_4^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 67

Space group *Cmma*, No. 67, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
dim		2	2	2	2
Λ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim		2	2	2	2
Σ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Y_5^+	Y_5^-	T_5^+	T_5^-
dim		2	2	2	2
H_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Y_5^+	Y_5^-	T_5^+	T_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5^+	Z_5^-	T_5^+	T_5^-
dim	2	2	2	2
A_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5^+	Z_5^-	T_5^+	T_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	S_2S_2	R_2R_2
dim	4	4
D_3D_3	2	1 -1
D_4D_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_2S_2	R_2R_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 68

Space group *Ccca*, No. 68, with SOC

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	4
Λ_5	2	1 1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Y(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2
Σ_5	2	1 1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

H

High symmetry line $H(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of H : $Y(\frac{1}{2}, \frac{1}{2}, 0), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Y_5^+	Y_5^-	T_3T_4
dim	2	2	4
H_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Y_5^+	Y_5^-	T_3T_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

A

High symmetry line $A(u, u, \frac{1}{2})$, high symmetry points at both ends of A: $Z(0, 0, \frac{1}{2}), T(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_3Z_4	T_3T_4
dim	4	4
A_2A_4	2	1
A_3A_5	2	1
		-1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	T_3T_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(0, \frac{1}{2}, u)$, high symmetry points at both ends of D: $S(0, \frac{1}{2}, 0), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	S_2S_2	R_2R_2
dim	4	4
D_3D_3	2	1
D_4D_4	2	1
		-1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_2S_2	R_2R_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 69

Space group $Fmmm$, No. 69, with SOC

Λ

High symmetry line $\Lambda(u, u, 0)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-
dim	2	2	2	2
Λ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Z_5^+	Z_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	X_5^+	X_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(0, -u, -u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(0, -\frac{1}{2}, -\frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	Y_5^+	Y_5^-
dim	2	2	2	2	2
Δ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	Y_5^+	Y_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 70

Space group $Fddd$, No. 70, with SOC

Λ

High symmetry line $\Lambda(u, u, 0)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_5^+	Γ_5^-	Z_3Z_4
dim	2	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Z_3Z_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_5^+	Γ_5^-	X_3X_4
dim	2	2	2	4
Σ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	X_3X_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, -u, -u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), Y(0, -\frac{1}{2}, -\frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	Y_3Y_4	
dim	2	2	4	
Δ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_5^+	Γ_5^-	Y_3Y_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 71

Space group *Immm*, No. 71, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	X_5^+	X_5^-	
dim	2	2	2	2	
Σ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	$R_3^+ R_4^+$	$R_3^- R_4^-$	W_5
dim	2	2	2
$Q_3 Q_4$	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$R_3^+ R_4^+$	$R_3^- R_4^-$	W_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	$S_3^+ S_4^+$	$S_3^- S_4^-$	W_5
dim	2	2	2
$D_3 D_4$	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$S_3^+ S_4^+$	$S_3^- S_4^-$	W_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	$T_3^+ T_4^+$	$T_3^- T_4^-$	W_5
dim	2	2	2
$P_3 P_4$	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$T_3^+ T_4^+$	$T_3^- T_4^-$	W_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 72

Space group *Ibam*, No. 72, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	X_5^+	X_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q : $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	$R_2 R_2$	$W_5 W_5$
dim	4	4
$Q_3 Q_3$	2	1
$Q_4 Q_4$	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$R_2 R_2$	$W_5 W_5$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		S_2S_2	W_5W_5
dim		4	4
D_3D_3	2	1	-1
D_4D_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	S_2S_2	W_5W_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		$T_3^+T_4^+$	$T_3^-T_4^-$	W_5W_5
dim		2	2	4
P_3P_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$T_3^+T_4^+$	$T_3^-T_4^-$	W_5W_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 73

Space group *Ibca*, No. 73, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_5^+	Γ_5^-	X_5^+	X_5^-
	dim	2	2	2	2
Σ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		R_2R_2	W_2W_2	W_3W_3	W_4W_4	W_5W_5
	dim	4	2	2	2	2
Q_3Q_3	2	1	0	-1	0	-1
Q_4Q_4	2	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_2R_2	W_2W_2	W_3W_3	W_4W_4	W_5W_5	Bipartite graph
	4	2	2	2	2	
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep		S_2S_2	W_2W_2	W_3W_3	W_4W_4	W_5W_5
	dim	4	2	2	2	2
D_3D_3	2	1	-1	-1	0	0
D_4D_4	2	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	S_2S_2	W_2W_2	W_3W_3	W_4W_4	W_5W_5	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	T_2T_2	W_2W_2	W_3W_3	W_4W_4	W_5W_5
dim	4	2	2	2	2
P_3P_3	2	1	-1	0	0
P_4P_4	2	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	T_2T_2	W_2W_2	W_3W_3	W_4W_4	W_5W_5	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

SG 74

Space group *Imma*, No. 74, with SOC

Σ

High symmetry line $\Sigma(u, -u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), X(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	X_5^+	X_5^-
dim	2	2	2	2
Σ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	X_5^+	X_5^-	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u + \frac{1}{2}, -u, -u)$, high symmetry points at both ends of Q: $R(\frac{1}{2}, 0, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	$R_3^+ R_4^+$	$R_3^- R_4^-$	$W_2 W_5$	$W_3 W_4$
dim	2	2	2	2
$Q_3 Q_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$R_3^+ R_4^+$	$R_3^- R_4^-$	$W_2 W_5$	$W_3 W_4$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(u + \frac{1}{2}, -u, u - \frac{1}{2})$, high symmetry points at both ends of D: $S(\frac{1}{2}, 0, -\frac{1}{2}), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$

A matrix

corep	$S_3^+ S_4^+$	$S_3^- S_4^-$	$W_2 W_5$	$W_3 W_4$
dim	2	2	2	2
$D_3 D_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$S_3^+ S_4^+$	$S_3^- S_4^-$	$W_2 W_5$	$W_3 W_4$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(u + \frac{1}{2}, u - \frac{1}{2}, -u)$, high symmetry points at both ends of P: $T(\frac{1}{2}, -\frac{1}{2}, 0), W(\frac{3}{4}, -\frac{1}{4}, -\frac{1}{4})$
A matrix

corep	T_2T_2	W_2W_5	W_3W_4
dim	4	2	2
P_3P_3	2	1	-1
P_4P_4	2	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	T_2T_2	W_2W_5	W_3W_4	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

SG 75

Space group $P4$, No. 75, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
dim	2	2	2
Δ_2	1	2	2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7
dim		2	2	2	2
Λ_5	1	1	0	-1	0
Λ_6	1	0	1	0	-1
Λ_7	1	0	1	0	-1
Λ_8	1	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7	
		2	2	2	2	Bipartite graph
v^1	0	1	0	1		$H_{2/2}^1$
v^2	1	0	1	0		$H_{2/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7
dim		2	2	2	2
Σ_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7	
		2	2	2	2	Bipartite graph
v^1	0	1	0	1		$H_{2/2}^1$
v^2	0	1	1	0		$H_{2/2}^1$
v^3	1	0	0	1		$H_{2/2}^1$
v^4	1	0	1	0		$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5M_8	M_6M_7	A_5A_8	A_6A_7
dim		2	2	2	2
V_5	1	1	0	-1	0
V_6	1	0	1	0	-1
V_7	1	0	1	0	-1
V_8	1	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5M_8	M_6M_7	A_5A_8	A_6A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_5M_8	M_6M_7	X_3X_4
dim	2	2	2
Y_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5M_8	M_6M_7	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_5Z_8	Z_6Z_7	R_3R_4
dim	2	2	2
U_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5Z_8	Z_6Z_7	R_3R_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_8	Z_6Z_7	A_5A_8	A_6A_7
	dim	2	2	2	2
S_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5Z_8	Z_6Z_7	A_5A_8	A_6A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_5A_8	A_6A_7	R_3R_4
	dim	2	2	2
T_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_5A_8	A_6A_7	R_3R_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_3R_4	X_3X_4
	dim	2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 76

Space group $P4_1$, No. 76, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
dim		2	2	2
Δ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_5	Z_6Z_8	Z_7Z_7
dim		2	2	2	2	2
Λ_5	1	1	0	-2	0	0
Λ_6	1	0	1	0	-1	0
Λ_7	1	0	1	0	0	-2
Λ_8	1	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_5	Z_6Z_8	Z_7Z_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	2	1	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7
dim		2	2	2	2
Σ_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5M_8	M_6M_7	A_5A_5	A_6A_8	A_7A_7
dim		2	2	2	2	2
V_5	1	1	0	-2	0	0
V_6	1	0	1	0	-1	0
V_7	1	0	1	0	0	-2
V_8	1	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5M_8	M_6M_7	A_5A_5	A_6A_8	A_7A_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	2	1	$H_{8/8}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y : $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_5M_8	M_6M_7	X_3X_4
dim		2	2	2
Y_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5M_8	M_6M_7	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	Z_6Z_8	Z_7Z_7	R_3R_3	R_4R_4
dim	2	2	2	2	2
U_2U_2	2	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_5Z_5	Z_6Z_8	Z_7Z_7	R_3R_3	R_4R_4	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	Z_6Z_8	Z_7Z_7	A_5A_5	A_6A_8	A_7A_7
dim	2	2	2	2	2	2
S_2S_2	2	1	1	1	-1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Z_5Z_5	Z_6Z_8	Z_7Z_7	A_5A_5	A_6A_8	A_7A_7	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$

v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5A_5	A_6A_8	A_7A_7	R_3R_3	R_4R_4
dim	2	2	2	2	2
T_2T_2	2	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_5A_5	A_6A_8	A_7A_7	R_3R_3	R_4R_4	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_3R_3	R_4R_4	X_3X_4
dim	2	2	2
W_3	1	2	0 -1
W_4	1	0	2 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_3	R_4R_4	X_3X_4	
	2	2	2	Bipartite graph
v^1	1	1	2	$H_{4/4}^2$

SG 77

Space group $P4_2$, No. 77, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
dim		2	2	2
Δ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7
dim		2	2	2	2
Λ_5	1	1	0	-1	0
Λ_6	1	0	1	-1	0
Λ_7	1	0	1	0	-1
Λ_8	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7
dim		2	2	2	2
Σ_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5M_8	M_6M_7	A_5A_8	A_6A_7
dim		2	2	2	2
V_5	1	1	0	-1	0
V_6	1	0	1	-1	0
V_7	1	0	1	0	-1
V_8	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5M_8	M_6M_7	A_5A_8	A_6A_7	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_5M_8	M_6M_7	X_3X_4
dim		2	2	2
Y_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5M_8	M_6M_7	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_8	Z_6Z_7	R_3R_4
dim		2	2	2
U_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5Z_8	Z_6Z_7	R_3R_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_8	Z_6Z_7	A_5A_8	A_6A_7
dim		2	2	2	2
S_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5Z_8	Z_6Z_7	A_5A_8	A_6A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_5A_8	A_6A_7	R_3R_4
dim		2	2	2
T_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_5A_8	A_6A_7	R_3R_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_3R_4	X_3X_4
dim		2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 78

Space group $P4_3$, No. 78, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
dim		2	2	2
Δ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_5	Z_6Z_8	Z_7Z_7
dim		2	2	2	2	2
Λ_5	1	1	0	0	-1	0
Λ_6	1	0	1	-2	0	0
Λ_7	1	0	1	0	-1	0
Λ_8	1	1	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_5	Z_6Z_8	Z_7Z_7	
		2	2	2	2	2	Bipartite graph
v^1		2	2	1	2	1	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7
dim		2	2	2	2
Σ_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7	
		2	2	2	2	Bipartite graph
v^1		0	1	0	1	$H_{2/2}^1$
v^2		0	1	1	0	$H_{2/2}^1$
v^3		1	0	0	1	$H_{2/2}^1$
v^4		1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5M_8	M_6M_7	A_5A_5	A_6A_8	A_7A_7
	dim	2	2	2	2	2
V_5	1	1	0	0	-1	0
V_6	1	0	1	-2	0	0
V_7	1	0	1	0	-1	0
V_8	1	1	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5M_8	M_6M_7	A_5A_5	A_6A_8	A_7A_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	2	1	$H_{8/8}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_5M_8	M_6M_7	X_3X_4
	dim	2	2	2
Y_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5M_8	M_6M_7	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_5Z_5	Z_6Z_8	Z_7Z_7	R_3R_3	R_4R_4
	dim	2	2	2	2	2
U_2U_2	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_5Z_5	Z_6Z_8	Z_7Z_7	R_3R_3	R_4R_4	Bipartite graph
	2	2	2	2	2	
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_5	Z_6Z_8	Z_7Z_7	A_5A_5	A_6A_8	A_7A_7
dim	2	2	2	2	2	2
S_2S_2	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Z_5Z_5	Z_6Z_8	Z_7Z_7	A_5A_5	A_6A_8	A_7A_7	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5A_5	A_6A_8	A_7A_7	R_3R_3	R_4R_4
dim	2	2	2	2	2
T_2T_2	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_5A_5	A_6A_8	A_7A_7	R_3R_3	R_4R_4	Bipartite graph
	2	2	2	2	2	
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_3R_3	R_4R_4	X_3X_4
	dim	2	2	2
W_3	1	2	0	-1
W_4	1	0	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_3	R_4R_4	X_3X_4	Bipartite graph
	2	2	2	
v^1	1	1	2	$H_{4/4}^2$

SG 79

Space group $I4$, No. 79, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7
	dim	2	2	2	2
Σ_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
	dim	2	2	2
Δ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_2N_2	P_3P_4
	dim	2	2
Q_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	N_2N_2	P_3P_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W : $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_3X_4	P_3P_4
	dim	2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	P_3P_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	Z_5Z_8	Z_6Z_7
	dim	2	2	2
Y_2	1	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	Z_5Z_8	Z_6Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 80

Space group $I4_1$, No. 80, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7
	dim	2	2	2	2
Σ_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
	dim	2	2	2
Δ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_2N_2	P_3	P_4P_4
	dim	2	1	2
Q_2	1	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	N_2N_2	P_3	P_4P_4	
	2	1	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_3X_4	P_3	P_4P_4
	dim	2	1	2
W_3	1	1	-1	0
W_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	P_3	P_4P_4	
	2	1	2	Bipartite graph
v^1	2	2	1	$H_{5/4}^3$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	Z_5Z_8	Z_6Z_7
	dim	2	2	2
Y_2	1	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	Z_5Z_8	Z_6Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 81

Space group $P\bar{4}$, No. 81, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
	dim	2	2	2
Δ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7
dim	2	2	2	2
$\Lambda_3\Lambda_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7
dim	2	2	2	2
Σ_2	1	2	2	-2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	M_5M_8	M_6M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_5M_8	M_6M_7	A_5A_8	A_6A_7
dim	2	2	2	2
V_3V_4	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_5M_8	M_6M_7	A_5A_8	A_6A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_5M_8	M_6M_7	X_3X_4
dim	2	2	2
Y_2	1	2	2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5M_8	M_6M_7	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5Z_8	Z_6Z_7	R_3R_4
dim	2	2	2
U_2	1	2	2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5Z_8	Z_6Z_7	R_3R_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_5Z_8	Z_6Z_7	A_5A_8	A_6A_7
dim		2	2	2	2
S_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5Z_8	Z_6Z_7	A_5A_8	A_6A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_5A_8	A_6A_7	R_3R_4
dim		2	2	2
T_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_5A_8	A_6A_7	R_3R_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_3R_4	X_3X_4
dim		2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 82

Space group $I\bar{4}$, No. 82, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7
dim		2	2	2	2
Σ_2	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	Z_5Z_8	Z_6Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4
dim		2	2	2
Δ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_5\Gamma_8$	$\Gamma_6\Gamma_7$	X_3X_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_2N_2	P_5	P_6	P_7	P_8
dim		2	1	1	1	1
Q_2	1	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	N_2N_2	P_5	P_6	P_7	P_8	
	2	1	1	1	1	Bipartite graph
v^1	1	0	0	0	2	$H_{3/2}^1$
v^2	1	0	0	1	1	$H_{3/2}^1$
v^3	1	0	0	2	0	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$
v^5	1	0	1	1	0	$H_{3/2}^1$
v^6	1	0	2	0	0	$H_{3/2}^1$
v^7	1	1	0	0	1	$H_{3/2}^1$
v^8	1	1	0	1	0	$H_{3/2}^1$
v^9	1	1	1	0	0	$H_{3/2}^1$
v^{10}	1	2	0	0	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_3X_4	P_5	P_6	P_7	P_8
	dim	2	1	1	1	1
W_3	1	1	-1	0	-1	0
W_4	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_3X_4	P_5	P_6	P_7	P_8	
	2	1	1	1	1	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	Z_5Z_8	Z_6Z_7
	dim	2	2	2
Y_2	1	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	Z_5Z_8	Z_6Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 83

Space group $P4/m$, No. 83, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$X_3^+X_4^+$	$X_3^-X_4^-$
	dim	2	2	2	2	2	2
$\Delta_3\Delta_4$	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$X_3^+X_4^+$	$X_3^-X_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$
dim		2	2	2	2	2	2	2	2
$\Lambda_5\Lambda_8$	2	1	1	0	0	-1	-1	0	0
$\Lambda_6\Lambda_7$	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$M_5^+M_6^+$	$M_5^-M_6^-$	$M_7^+M_8^+$	$M_7^-M_8^-$
dim		2	2	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$\Gamma_7^+ \Gamma_8^+$	$\Gamma_7^- \Gamma_8^-$	$M_5^+ M_6^+$	$M_5^- M_6^-$	$M_7^+ M_8^+$	$M_7^- M_8^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_5^+ M_6^+$	$M_5^- M_6^-$	$M_7^+ M_8^+$	$M_7^- M_8^-$	$A_5^+ A_6^+$	$A_5^- A_6^-$	$A_7^+ A_8^+$	$A_7^- A_8^-$	
dim	2	2	2	2	2	2	2	2	
$V_5 V_8$	2	1	1	0	0	-1	-1	0	0
$V_6 V_7$	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$M_5^+ M_6^+$	$M_5^- M_6^-$	$M_7^+ M_8^+$	$M_7^- M_8^-$	$A_5^+ A_6^+$	$A_5^- A_6^-$	$A_7^+ A_8^+$	$A_7^- A_8^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$M_5^+ M_6^+$	$M_5^- M_6^-$	$M_7^+ M_8^+$	$M_7^- M_8^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$
dim	2	2	2	2	2	2
$Y_3 Y_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$M_5^+ M_6^+$	$M_5^- M_6^-$	$M_7^+ M_8^+$	$M_7^- M_8^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$
dim	2	2	2	2	2	2
$U_3 U_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$	$A_5^+ A_6^+$	$A_5^- A_6^-$	$A_7^+ A_8^+$	$A_7^- A_8^-$
dim	2	2	2	2	2	2	2	2
$S_3 S_4$	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$	$A_5^+ A_6^+$	$A_5^- A_6^-$	$A_7^+ A_8^+$	$A_7^- A_8^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_5^+ A_6^+$	$A_5^- A_6^-$	$A_7^+ A_8^+$	$A_7^- A_8^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$
dim	2	2	2	2	2	2
$T_3 T_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

$A_5^+ A_6^+$	$A_5^- A_6^-$	$A_7^+ A_8^+$	$A_7^- A_8^-$	$R_3^+ R_4^+$	$R_3^- R_4^-$	Bipartite graph
2	2	2	2	2	2	

v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	$R_3^+ R_4^+$	$R_3^- R_4^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$
dim	2	2	2	2
$W_3 W_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$R_3^+ R_4^+$	$R_3^- R_4^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 84

Space group $P4_2/m$, No. 84, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$\Gamma_7^+ \Gamma_8^+$	$\Gamma_7^- \Gamma_8^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$
dim	2	2	2	2	2	2
$\Delta_3 \Delta_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$X_3^+X_4^+$	$X_3^-X_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	Z_3Z_4
	dim	2	2	2	2	4
$\Lambda_5\Lambda_8$	2	1	1	0	0	-1
$\Lambda_6\Lambda_7$	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	Z_3Z_4	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$M_5^+M_6^+$	$M_5^-M_6^-$	$M_7^+M_8^+$	$M_7^-M_8^-$
	dim	2	2	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$M_5^+M_6^+$	$M_5^-M_6^-$	$M_7^+M_8^+$	$M_7^-M_8^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$M_5^+M_6^+$	$M_5^-M_6^-$	$M_7^+M_8^+$	$M_7^-M_8^-$	A_3A_4
dim	2	2	2	2	4
V_5V_8 2	1	1	0	0	-1
V_6V_7 2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_5^+M_6^+$	$M_5^-M_6^-$	$M_7^+M_8^+$	$M_7^-M_8^-$	A_3A_4	Bipartite graph
	2	2	2	2	4	
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $X(0, \frac{1}{2}, 0)$
A matrix

corep	$M_5^+ M_6^+$	$M_5^- M_6^-$	$M_7^+ M_8^+$	$M_7^- M_8^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$
dim	2	2	2	2	2	2
$Y_3 Y_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$M_5^+ M_6^+$	$M_5^- M_6^-$	$M_7^+ M_8^+$	$M_7^- M_8^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2})$, $R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Z_3 Z_4$	$R_3^+ R_4^+$	$R_3^- R_4^-$
dim	4	2	2
$U_3 U_4$	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$Z_3 Z_4$	$R_3^+ R_4^+$	$R_3^- R_4^-$	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2})$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Z_3 Z_4$	$A_3 A_4$
dim	4	4
$S_3 S_4$	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	A_3A_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_3A_4	$R_3^+R_4^+$	$R_3^-R_4^-$
dim	4	2	2
T_3T_4	2	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_3A_4	$R_3^+R_4^+$	$R_3^-R_4^-$	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	$R_3^+R_4^+$	$R_3^-R_4^-$	$X_3^+X_4^+$	$X_3^-X_4^-$
dim	2	2	2	2
W_3W_4	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$R_3^+R_4^+$	$R_3^-R_4^-$	$X_3^+X_4^+$	$X_3^-X_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 85

Space group $P4/n$, No. 85, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	X_2X_2
dim	2	2	2	2	4
$\Delta_3\Delta_4$ 2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	X_2X_2	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$
dim	2	2	2	2	2	2	2	2
$\Lambda_5\Lambda_8$ 2	1	1	0	0	-1	-1	0	0
$\Lambda_6\Lambda_7$ 2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$\Gamma_7^+ \Gamma_8^+$	$\Gamma_7^- \Gamma_8^-$	$M_3 M_4$
dim	2	2	2	2	4
$\Sigma_3 \Sigma_4$	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$\Gamma_7^+ \Gamma_8^+$	$\Gamma_7^- \Gamma_8^-$	$M_3 M_4$	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_3 M_4$	$A_3 A_4$
dim	4	4
$V_5 V_6$	2	1
$V_7 V_8$	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$M_3 M_4$	$A_3 A_4$	Bipartite graph
	4	4	
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_3M_4	X_2X_2
dim	4	4
Y_3Y_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	X_2X_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$	R_2R_2
dim	2	2	2	2	4
U_3U_4	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$	R_2R_2	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$	$A_3 A_4$
dim	2	2	2	2	4
$S_3 S_4$	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$	$A_3 A_4$	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$A_3 A_4$	$R_2 R_2$
dim	4	4
$T_3 T_4$	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$A_3 A_4$	$R_2 R_2$	Bipartite graph
	4	4	
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	$R_2 R_2$	$X_2 X_2$
dim	4	4
$W_3 W_3$	2	-1
$W_4 W_4$	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_2R_2	X_2X_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 86

Space group $P4_2/n$, No. 86, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	X_2X_2
dim	2	2	2	2	4
$\Delta_3\Delta_4$	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	X_2X_2	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	Z_3Z_4
dim		2	2	2	2	4
$\Lambda_5\Lambda_8$	2	1	1	0	0	-1
$\Lambda_6\Lambda_7$	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	Z_3Z_4	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	M_3M_4
dim		2	2	2	2	4
$\Sigma_3\Sigma_4$	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	M_3M_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_3M_4	$A_5^+A_6^+$	$A_5^-A_6^-$	$A_7^+A_8^+$	$A_7^-A_8^-$
dim		4	2	2	2	2
V_5V_6	2	1	0	0	-1	-1
V_7V_8	2	1	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_3M_4	$A_5^+A_6^+$	$A_5^-A_6^-$	$A_7^+A_8^+$	$A_7^-A_8^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_3M_4	X_2X_2
dim		4	4
Y_3Y_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	X_2X_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_3Z_4	R_2R_2
dim		4	4
U_3U_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_3Z_4	R_2R_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_4	$A_5^+A_6^+$	$A_5^-A_6^-$	$A_7^+A_8^+$	$A_7^-A_8^-$
dim	4	2	2	2	2
S_3S_4	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_3Z_4	$A_5^+A_6^+$	$A_5^-A_6^-$	$A_7^+A_8^+$	$A_7^-A_8^-$	Bipartite graph
	4	2	2	2	2	
v^1	1	0	0	0	2	$H_{3/2}^1$
v^2	1	0	0	1	1	$H_{3/2}^1$
v^3	1	0	0	2	0	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$
v^5	1	0	1	1	0	$H_{3/2}^1$
v^6	1	0	2	0	0	$H_{3/2}^1$
v^7	1	1	0	0	1	$H_{3/2}^1$
v^8	1	1	0	1	0	$H_{3/2}^1$
v^9	1	1	1	0	0	$H_{3/2}^1$
v^{10}	1	2	0	0	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_5^+A_6^+$	$A_5^-A_6^-$	$A_7^+A_8^+$	$A_7^-A_8^-$	R_2R_2
dim	2	2	2	2	4
T_3T_4	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$A_5^+A_6^+$	$A_5^-A_6^-$	$A_7^+A_8^+$	$A_7^-A_8^-$	R_2R_2	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$

v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_2R_2	X_2X_2
	dim	4	4
W_3W_3	2	1	-1
W_4W_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_2R_2	X_2X_2	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 87

Space group $I4/m$, No. 87, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$
	dim	2	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$Z_5^+Z_6^+$	$Z_5^-Z_6^-$	$Z_7^+Z_8^+$	$Z_7^-Z_8^-$	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$

v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$\Gamma_7^+ \Gamma_8^+$	$\Gamma_7^- \Gamma_8^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$
dim	2	2	2	2	2	2
$\Delta_3 \Delta_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$\Gamma_7^+ \Gamma_8^+$	$\Gamma_7^- \Gamma_8^-$	$X_3^+ X_4^+$	$X_3^- X_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	$N_2^+ N_2^+$	$N_2^- N_2^-$	$P_5 P_8$	$P_6 P_7$
dim	2	2	2	2
$Q_2 Q_2$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$N_2^+ N_2^+$	$N_2^- N_2^-$	$P_5 P_8$	$P_6 P_7$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	$X_3^+ X_4^+$	$X_3^- X_4^-$	$P_5 P_8$	$P_6 P_7$
dim	2	2	2	2
$W_3 W_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$X_3^+ X_4^+$	$X_3^- X_4^-$	$P_5 P_8$	$P_6 P_7$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$X_3^+ X_4^+$	$X_3^- X_4^-$	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$
dim	2	2	2	2	2	2
$Y_3 Y_4$	2	1	1	-1	-1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$X_3^+ X_4^+$	$X_3^- X_4^-$	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$Z_7^+ Z_8^+$	$Z_7^- Z_8^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	$H_{2/1}^1$

v^6	1	0	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	1	0	0	0	$H_{2/1}^1$

SG 88

Space group $I4_1/a$, No. 88, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	Z_3Z_4
dim	2	2	2	2	4
$\Sigma_3\Sigma_4$	2	1	1	1	1
					-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	Z_3Z_4	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	X_2X_2
dim	2	2	2	2	4
$\Delta_3\Delta_4$	2	1	1	1	1
					-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	X_2X_2	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	$N_2^+N_2^+$	$N_2^-N_2^-$	P_5P_7	P_6P_6	P_8P_8
dim	2	2	2	2	2
Q_2Q_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$N_2^+N_2^+$	$N_2^-N_2^-$	P_5P_7	P_6P_6	P_8P_8	Bipartite graph
	2	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	X_2X_2	P_5P_7	P_6P_6	P_8P_8
dim	4	2	2	2
W_3W_3	2	1	-1	0
W_4W_4	2	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_2X_2	P_3P_7	P_6P_6	P_8P_8	
	4	2	2	2	Bipartite graph
v^1	1	1	0	1	$H_{3/2}^1$
v^2	1	1	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_2X_2	Z_3Z_4
dim	4	4
Y_3Y_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_2X_2	Z_3Z_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 89

Space group $P422$, No. 89, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_6	Γ_7	X_5
dim	2	2	2
Δ_3	1	1	-1
Δ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
dim		2	2	2	2
Λ_5	1	1	0	-1	0
Λ_6	1	0	1	0	-1
Λ_7	1	0	1	0	-1
Λ_8	1	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6	M_7
dim		2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	A_6	A_7
dim		2	2	2	2
V_5	1	1	0	-1	0
V_6	1	0	1	0	-1
V_7	1	0	1	0	-1
V_8	1	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

		M_6	M_7	A_6	A_7	
		2	2	2	2	Bipartite graph
v^1	0	1	0	1		$H_{2/2}^1$
v^2	1	0	1	0		$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
dim		2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

		M_6	M_7	X_5	
		2	2	2	Bipartite graph
v^1	0	1	1		$H_{2/2}^1$
v^2	1	0	1		$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	R_5
dim		2	2	2
U_3	1	1	1	-1
U_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6	Z_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	A_6	A_7
	dim	2	2	2	2
S_3	1	1	1	-1	-1
S_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_6	Z_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6	A_7	R_5
	dim	2	2	2
T_3	1	1	1	-1
T_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_6	A_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_5	X_5
dim		2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 90

Space group $P4_212$, No. 90, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
dim		2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Λ_5	1	1	0	-1	0
Λ_6	1	0	1	0	-1
Λ_7	1	0	1	0	-1
Λ_8	1	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6M_7	A_6A_7
	dim	4	4
V_5V_7	2	1	-1
V_6V_8	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6M_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_6M_7	X_2X_3	X_4X_5
dim	4	2	2
Y_3Y_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6	Z_7	R_2R_3	R_4R_5
dim	2	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	A_6A_7
	dim	2	2	4
S_3	1	1	1	-2
S_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	A_6A_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6A_7	R_2R_3	R_4R_5
	dim	4	2	2
T_3T_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_6A_7	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_2R_3	R_4R_5	X_2X_3	X_4X_5
	dim	2	2	2	2
W_3W_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 91

Space group $P4_122$, No. 91, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_6	Γ_7	X_5
dim	2	2	2
Δ_3	1	1	-1
Δ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7
dim	2	2	2	2	2
Λ_5	1	1	0	-2	0
Λ_6	1	0	1	0	-1
Λ_7	1	0	1	0	-2
Λ_8	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	1	2	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	A_3A_5	A_4A_6	A_7
	dim	2	2	2	2	2
V_5	1	1	0	-2	0	0
V_6	1	0	1	0	0	-1
V_7	1	0	1	0	-2	0
V_8	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6	M_7	A_3A_5	A_4A_6	A_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	1	2	$H_{8/8}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_3Z_5	Z_4Z_6	Z_7	R_2R_3	R_4R_5
	dim	2	2	2	2	2
U_3U_4	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_3Z_5	Z_4Z_6	Z_7	R_2R_3	R_4R_5	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_5	Z_4Z_6	Z_7	A_3A_5	A_4A_6	A_7
dim	2	2	2	2	2	2
S_3S_4	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Z_3Z_5	Z_4Z_6	Z_7	A_3A_5	A_4A_6	A_7	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_3A_5	A_4A_6	A_7	R_2R_3	R_4R_5
dim	2	2	2	2	2
T_3T_4	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_3A_5	A_4A_6	A_7	R_2R_3	R_4R_5	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_2R_3	R_4R_5	X_5
	dim	2	2	2
W_3	1	0	2	-1
W_4	1	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_2R_3	R_4R_5	X_5	
	2	2	2	Bipartite graph
v^1	1	1	2	$H_{4/4}^2$

SG 92

Space group $P4_12_12$, No. 92, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7
	dim	2	2	2	2	2
Λ_5	1	1	0	-2	0	0
Λ_6	1	0	1	0	0	-1
Λ_7	1	0	1	0	-2	0
Λ_8	1	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	1	2	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_6M_7	A_3A_6	A_4A_5	A_7A_7
	dim	4	2	2	4
V_5V_7	2	1	0	0	-2
V_6V_8	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	A_3A_6	A_4A_5	A_7A_7	
	4	2	2	4	Bipartite graph
v^1	2	0	2	1	$H_{5/4}^3$
v^2	2	1	1	1	$H_{5/4}^3$
v^3	2	2	0	1	$H_{5/4}^3$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_6M_7	X_2X_3	X_4X_5
dim	4	2	2
Y_3Y_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_5	Z_4Z_6	Z_7	R_5R_5
dim	2	2	2	4
U_3U_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_3Z_5	Z_4Z_6	Z_7	R_5R_5	
	2	2	2	4	Bipartite graph
v^1	0	0	2	1	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	2	0	1	$H_{3/2}^1$
v^4	1	0	1	1	$H_{3/2}^1$
v^5	1	1	0	1	$H_{3/2}^1$
v^6	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_5	Z_4Z_6	Z_7	A_3A_6	A_4A_5	A_7A_7	
dim	2	2	2	2	2	4	
S_3S_4	2	1	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_3Z_5	Z_4Z_6	Z_7	A_3A_6	A_4A_5	A_7A_7	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	1	1	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	0	1	$H_{3/2}^1$
v^{11}	1	1	0	0	0	1	$H_{3/2}^1$
v^{12}	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_3A_6	A_4A_5	A_7A_7	R_5R_5	
dim	2	2	4	4	
T_3T_3	2	1	0	1	-1
T_4T_4	2	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_3A_6	A_4A_5	A_7A_7	R_5R_5	
	2	2	4	4	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	1	1	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	R_5R_5	X_2X_3	X_4X_5
dim	4	2	2
W_3W_4	2	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	R_5R_5	X_2X_3	X_4X_5	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

SG 93

Space group $P4_222$, No. 93, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_6	Γ_7	X_5
dim	2	2	2
Δ_3	1	1	1 -1
Δ_4	1	1	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_6	Γ_7	Z_6	Z_7
dim	2	2	2	2
Λ_5	1	1	0	-1 0
Λ_6	1	0	1	-1 0
Λ_7	1	0	1	0 -1
Λ_8	1	1	0	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_6	M_7	A_6	A_7
	dim	2	2	2	2
V_5	1	1	0	-1	0
V_6	1	0	1	-1	0
V_7	1	0	1	0	-1
V_8	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6	M_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	R_5
	dim	2	2	2
U_3	1	1	1	-1
U_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6	Z_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	A_6	A_7
	dim	2	2	2	2
S_3	1	1	1	-1	-1
S_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_6	Z_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_6	A_7	R_5
	dim	2	2	2
T_3	1	1	1	-1
T_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_6	A_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_5	X_5
	dim	2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 94

Space group $P4_22_12$, No. 94, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
dim		2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
dim		2	2	2	2
Λ_5	1	1	0	-1	0
Λ_6	1	0	1	-1	0
Λ_7	1	0	1	0	-1
Λ_8	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6M_7	A_6A_7
	dim	4	4
V_5V_7	2	1	-1
V_6V_8	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6M_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6M_7	X_2X_3	X_4X_5
	dim	4	2	2
Y_3Y_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	R_2R_3	R_4R_5
	dim	2	2	2	2
U_3	1	1	1	0	-2
U_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	A_6A_7
	dim	2	2	4
S_3	1	1	1	-2
S_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	A_6A_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_6A_7	R_2R_3	R_4R_5
dim	4	2	2
T_3T_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_6A_7	R_2R_3	R_4R_5	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_2R_3	R_4R_5	X_2X_3	X_4X_5
dim	2	2	2	2
W_3W_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 95

Space group $P4_322$, No. 95, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7
	dim	2	2	2	2	2
Λ_5	1	1	0	0	0	-1
Λ_6	1	0	1	-2	0	0
Λ_7	1	0	1	0	0	-1
Λ_8	1	1	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	1	2	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_6	M_7	A_3A_5	A_4A_6	A_7
	dim	2	2	2	2	2
V_5	1	1	0	0	0	-1
V_6	1	0	1	-2	0	0
V_7	1	0	1	0	0	-1
V_8	1	1	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6	M_7	A_3A_5	A_4A_6	A_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	1	2	$H_{8/8}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_5	Z_4Z_6	Z_7	R_2R_3	R_4R_5	
dim	2	2	2	2	2	
U_3U_4	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_3Z_5	Z_4Z_6	Z_7	R_2R_3	R_4R_5	
	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_5	Z_4Z_6	Z_7	A_3A_5	A_4A_6	A_7	
dim	2	2	2	2	2	2	
S_3S_4	2	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Z_3Z_5	Z_4Z_6	Z_7	A_3A_5	A_4A_6	A_7	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_3A_5	A_4A_6	A_7	R_2R_3	R_4R_5
dim	2	2	2	2	2
T_3T_4	2	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_3A_5	A_4A_6	A_7	R_2R_3	R_4R_5	Bipartite graph
	2	2	2	2	2	
v^1	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	0	1	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_2R_3	R_4R_5	X_5
dim	2	2	2
W_3	1	0	2 -1
W_4	1	2	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_2R_3	R_4R_5	X_5	Bipartite graph
	2	2	2	
v^1	1	1	2	$H_{4/4}^2$

SG 96

Space group $P4_32_12$, No. 96, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7
	dim	2	2	2	2	2
Λ_5	1	1	0	0	0	-1
Λ_6	1	0	1	-2	0	0
Λ_7	1	0	1	0	0	-1
Λ_8	1	1	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	Z_3Z_5	Z_4Z_6	Z_7	
	2	2	2	2	2	Bipartite graph
v^1	2	2	1	1	2	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_6M_7	A_3A_6	A_4A_5	A_7A_7
dim	4	2	2	4
V_5V_7	2	1	-1	-1
V_6V_8	2	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	A_3A_6	A_4A_5	A_7A_7	
	4	2	2	4	Bipartite graph
v^1	2	0	2	1	$H_{5/4}^3$
v^2	2	1	1	1	$H_{5/4}^3$
v^3	2	2	0	1	$H_{5/4}^3$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_6M_7	X_2X_3	X_4X_5
dim	4	2	2
Y_3Y_4	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_5	Z_4Z_6	Z_7	R_5R_5
dim	2	2	2	4
U_3U_4	2	1	1	1
				-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_3Z_5	Z_4Z_6	Z_7	R_5R_5	Bipartite graph
	2	2	2	4	
v^1	0	0	2	1	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	2	0	1	$H_{3/2}^1$
v^4	1	0	1	1	$H_{3/2}^1$
v^5	1	1	0	1	$H_{3/2}^1$
v^6	2	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_3Z_5	Z_4Z_6	Z_7	A_3A_6	A_4A_5	A_7A_7
dim	2	2	2	2	2	4
S_3S_4	2	1	1	1	-1	-1
						-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Z_3Z_5	Z_4Z_6	Z_7	A_3A_6	A_4A_5	A_7A_7	Bipartite graph
	2	2	2	2	2	4	
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	1	1	0	0	1	$H_{3/2}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	1	0	0	1	$H_{3/2}^1$
v^{11}	1	1	0	0	0	1	$H_{3/2}^1$
v^{12}	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_3A_6	A_4A_5	A_7A_7	R_5R_5
dim	2	2	4	4
T_3T_3	2	1	0	1
T_4T_4	2	0	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_3A_6	A_4A_5	A_7A_7	R_5R_5	Bipartite graph
	2	2	4	4	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	1	1	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_5R_5	X_2X_3	X_4X_5
dim	4	2	2
W_3W_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	R_5R_5	X_2X_3	X_4X_5	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

SG 97

Space group *I*422, No. 97, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix



corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_3N_4	P_5
	dim	2	2
Q_3	1	1	-1
Q_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	N_3N_4	P_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_5	P_5
	dim	2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	P_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	Z_6	Z_7
	dim	2	2	2
Y_3	1	1	-1	-1
Y_4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	Z_6	Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 98

Space group $I4_122$, No. 98, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
dim		2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	X_5
dim		2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_3N_4	P_2P_5	P_3	P_4
	dim	2	2	1	1
Q_3	1	1	-1	-1	0
Q_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	N_3N_4	P_2P_5	P_3	P_4	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_5	P_2P_5	P_3	P_4
	dim	2	2	1	1
W_3	1	1	0	-1	-1
W_4	1	1	-2	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_5	P_2P_5	P_3	P_4	
	2	2	1	1	Bipartite graph
v^1	2	1	0	2	$H_{5/4}^3$
v^2	2	1	1	1	$H_{5/4}^3$
v^3	2	1	2	0	$H_{5/4}^3$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_5	Z_6	Z_7
	dim	2	2	2
Y_3	1	1	-1	-1
Y_4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	Z_6	Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 99

Space group $P4mm$, No. 99, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	X_5
dim		2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
dim		2	2	2	2
Λ_6	2	1	0	-1	0
Λ_7	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	A_6	A_7
	dim	2	2	2	2
V_6	2	1	0	-1	0
V_7	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y : $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_6	Z_7	R_5
	dim	2	2	2
U_3	1	1	1	-1
U_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6	Z_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_6	Z_7	A_6	A_7
	dim	2	2	2	2
S_3	1	1	1	-1	-1
S_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_6	Z_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_6	A_7	R_5
	dim	2	2	2
T_3	1	1	1	-1
T_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_6	A_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_5	X_5
	dim	2	2
W_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 100

Space group $P4bm$, No. 100, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Λ_6	2	1	0	-1	0
Λ_7	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_6M_7	A_6A_7
	dim	4	4
V_6	2	1	-1
V_7	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6M_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_6M_7	X_2X_3	X_4X_5
	dim	4	2	2
Y_3Y_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	R_2R_3	R_4R_5
	dim	2	2	2	2
U_3	1	1	1	0	-2
U_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	A_6A_7
	dim	2	2	4
S_3	1	1	1	-2
S_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	A_6A_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6A_7	R_2R_3	R_4R_5
	dim	4	2	2
T_3T_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_6A_7	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_2R_3	R_4R_5	X_2X_3	X_4X_5
	dim	2	2	2	2
W_2	1	1	0	-1	0
W_3	1	0	1	0	-1
W_4	1	1	0	-1	0
W_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 101

Space group $P4_2cm$, No. 101, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	$Z_6 Z_7$
	dim	2	2	4
Λ_6	2	1	0	-1
Λ_7	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	A_6A_7
	dim	2	2	4
V_6	2	1	0	-1
V_7	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6	M_7	A_6A_7	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	R_5R_5
	dim	4	4
U_3U_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	A_6A_7
	dim	4	4
S_3	1	2	-2
S_4	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6A_7	R_5R_5
	dim	4	4
T_3T_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_6A_7	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix



corep		R_5R_5	X_5
	dim	4	2
W_5	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5R_5	X_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/2}^1$

SG 102

Space group $P4_2nm$, No. 102, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6Z_7
	dim	2	2	4
Λ_6	2	1	0	-1
Λ_7	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	$M_6 M_7$
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$M_6 M_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$M_6 M_7$	A_6	A_7
	dim	4	2	2
V_6	2	1	-1	0
V_7	2	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$M_6 M_7$	A_6	A_7	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y : $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_6M_7	X_2X_3	X_4X_5
dim	4	2	2
Y_3Y_4	2	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6Z_7	R_2R_3	R_4R_5
dim	4	2	2
U_3U_4	2	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6Z_7	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6Z_7	A_6	A_7
dim	4	2	2
S_3	1	2	-1 -1
S_4	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$Z_6 Z_7$	A_6	A_7	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	1	2	0	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_6	A_7	$R_2 R_3$	$R_4 R_5$
dim	2	2	2	2
T_3	1	1	1	-2 0
T_4	1	1	1	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_6	A_7	$R_2 R_3$	$R_4 R_5$	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	$R_2 R_3$	$R_4 R_5$	$X_2 X_3$	$X_4 X_5$
dim	2	2	2	2
W_2	1	0	1	-1 0
W_3	1	0	1	0 -1
W_4	1	1	0	-1 0
W_5	1	1	0	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$R_2 R_3$	$R_4 R_5$	$X_2 X_3$	$X_4 X_5$	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 103

Space group $P4cc$, No. 103, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6Z_6	Z_7Z_7
	dim	2	2	4	4
Λ_6	2	1	0	-2	0
Λ_7	2	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Z_6Z_6	Z_7Z_7	
	2	2	4	4	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	2	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	A_6A_6	A_7A_7
	dim	2	2	4	4
V_6	2	1	0	-2	0
V_7	2	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	A_6A_6	A_7A_7	
	2	2	4	4	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	2	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6Z_6	Z_7Z_7	R_5R_5
dim	4	4	4
U_3U_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6Z_6	Z_7Z_7	R_5R_5	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6Z_6	Z_7Z_7	A_6A_6	A_7A_7
dim	4	4	4	4
S_3S_4	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_6Z_6	Z_7Z_7	A_6A_6	A_7A_7	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T : $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_6A_6	A_7A_7	R_5R_5
dim	4	4	4
T_3T_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_6A_6	A_7A_7	R_5R_5	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_5R_5	X_5
dim	4	2
W_5	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5R_5	X_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/2}^1$

SG 104

Space group $P4nc$, No. 104, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_6	Γ_7	X_2X_3	X_4X_5
dim	2	2	2	2
Δ_3	1	1	0	-2
Δ_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6Z_6	Z_7Z_7
	dim	2	2	4	4
Λ_6	2	1	0	-2	0
Λ_7	2	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Z_6Z_6	Z_7Z_7	
	2	2	4	4	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	2	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6M_7	A_6A_7
dim		4	4
V_6	2	1	-1
V_7	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6M_7	A_6A_7	Bipartite graph
	4	4	
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6M_7	X_2X_3	X_4X_5
dim		4	2	2
Y_3Y_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_6	Z_7Z_7	R_2R_3	R_4R_5
dim		4	4	2	2
U_3U_4	2	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Z_6Z_6	Z_7Z_7	R_2R_3	R_4R_5	
	4	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	2	$H_{3/2}^1$
v^5	1	0	1	1	$H_{3/2}^1$
v^6	1	0	2	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6Z_6	Z_7Z_7	A_6A_7
dim	4	4	4
S_3S_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6Z_6	Z_7Z_7	A_6A_7	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_6A_7	R_2R_3	R_4R_5
dim	4	2	2
T_3	1	2	-2
T_4	1	2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_6A_7	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/4}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_2R_3	R_4R_5	X_2X_3	X_4X_5
dim		2	2	2	2
W_2	1	0	1	-1	0
W_3	1	0	1	0	-1
W_4	1	1	0	-1	0
W_5	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 105

Space group $P4_2mc$, No. 105, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_5
dim		2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	$Z_6 Z_7$
	dim	2	2	4
Λ_6	2	1	0	-1
Λ_7	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	$A_6 A_7$
	dim	2	2	4
V_6	2	1	0	-1
V_7	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6	M_7	$A_6 A_7$	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$Z_6 Z_7$	R_5
	dim	4	2
U_3	1	2	-1
U_4	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$Z_6 Z_7$	R_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/4}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6Z_7	A_6A_7
dim	4	4
S_3S_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_6A_7	R_5
dim		4	2
T_3	1	2	-1
T_4	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_6A_7	R_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/4}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_5	X_5
dim		2	2
W_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 106

Space group $P4_2bc$, No. 106, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6Z_7
	dim	2	2	4
Λ_6	2	1	0	-1
Λ_7	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_6	Γ_7	Z_6Z_7	
	2	2	4	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_6M_7	A_6A_6	A_7A_7
	dim	4	4	4
V_6	2	1	-2	0
V_7	2	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6M_7	A_6A_6	A_7A_7	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_6M_7	X_2X_3	X_4X_5
	dim	4	2	2
Y_3Y_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	R_2R_3	R_4R_5
	dim	4	2	2
U_3	1	2	0	-2
U_4	1	2	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	R_2R_3	R_4R_5	Bipartite graph
	4	2	2	
v^1	1	1	1	$H_{3/4}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	A_6A_6	A_7A_7
	dim	4	4	4
S_3S_4	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6Z_7	A_6A_6	A_7A_7	Bipartite graph
	4	4	4	
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6A_6	A_7A_7	R_2R_3	R_4R_5
	dim	4	4	2	2
T_3T_4	2	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_6A_6	A_7A_7	R_2R_3	R_4R_5	
	4	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	2	$H_{3/2}^1$
v^5	1	0	1	1	$H_{3/2}^1$
v^6	1	0	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_2R_3	R_4R_5	X_2X_3	X_4X_5
	dim	2	2	2	2
W_2	1	1	0	-1	0
W_3	1	0	1	0	-1
W_4	1	1	0	-1	0
W_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 107

Space group $I4mm$, No. 107, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_3N_4	P_5
	dim	2	2
Q_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	N_3N_4	P_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_5	P_5
	dim	2	2
W_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	P_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	Z_6	Z_7
	dim	2	2	2
Y_3	1	1	-1	-1
Y_4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	Z_6	Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 108

Space group $I4cm$, No. 108, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		N_3N_3	N_4N_4	P_5P_5
	dim	2	2	4
Q_2Q_2	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	N_3N_3	N_4N_4	P_5P_5	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	X_5	P_5P_5
dim	2	4
W_5	2	1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	P_5P_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_5	Z_6	Z_7
dim	2	2	2
Y_3	1	1	-1 -1
Y_4	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	Z_6	Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 109

Space group $I4_1md$, No. 109, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	$Z_6 Z_7$
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	$X_2 X_3$	$X_4 X_5$
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$X_2 X_3$	$X_4 X_5$	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		$N_3 N_4$	P_2	P_3	$P_4 P_5$
	dim	2	1	1	2
Q_2	1	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	N_3N_4	P_2	P_3	P_4P_5	
	2	1	1	2	Bipartite graph
v^1	1	0	0	1	$H_{2/2}^1$
v^2	1	0	2	0	$H_{3/2}^1$
v^3	1	1	1	0	$H_{3/2}^1$
v^4	1	2	0	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_2X_3	X_4X_5	P_2	P_3	P_4P_5
	dim	2	2	1	1	2
W_2	1	1	0	0	-1	0
W_3	1	0	1	0	0	-1
W_4	1	1	0	0	0	-1
W_5	1	0	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_2X_3	X_4X_5	P_2	P_3	P_4P_5	
	2	2	1	1	2	Bipartite graph
v^1	1	1	1	1	1	$H_{5/4}^3$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_2X_3	X_4X_5	Z_6Z_7
	dim	2	2	4
Y_3Y_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	Z_6Z_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 110

Space group $I4_1cd$, No. 110, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6Z_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Z_6Z_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_3N_3	N_4N_4	P_2P_3	P_4P_4	P_5P_5
dim		2	2	2	2	2
Q_2Q_2	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	N_3N_3	N_4N_4	P_2P_3	P_4P_4	P_5P_5	Bipartite graph
	2	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_2X_3	X_4X_5	P_2P_3	P_4P_4	P_5P_5
dim		2	2	2	2	2
W_2	1	1	0	-1	0	0
W_3	1	0	1	0	-2	0
W_4	1	1	0	0	0	-2
W_5	1	0	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_2X_3	X_4X_5	P_2P_3	P_4P_4	P_5P_5	Bipartite graph
	2	2	2	2	2	
v^1	2	2	2	1	1	$H_{8/8}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_2X_3	X_4X_5	Z_6Z_7
dim		2	2	4
Y_3Y_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	Z_6Z_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 111

Space group $P\bar{4}2m$, No. 111, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_6	Γ_7	X_5
dim	2	2	2
Δ_3	1	1	1
Δ_4	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_6	Γ_7	Z_6	Z_7
dim	2	2	2	2
Λ_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6	M_7
dim		2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	A_6	A_7
dim		2	2	2	2
V_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_6	M_7	A_6	A_7	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y : $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix



corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_6	Z_7	R_5
	dim	2	2	2
U_3	1	1	1	-1
U_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6	Z_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_6	Z_7	A_6	A_7
	dim	2	2	2	2
S_3	1	1	1	-1	-1
S_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_6	Z_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_6	A_7	R_5
	dim	2	2	2
T_3	1	1	1	-1
T_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_6	A_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_5	X_5
	dim	2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 112

Space group $P\bar{4}2c$, No. 112, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	$Z_6 Z_7$
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_6	M_7	A_6A_7
dim	2	2	4
V_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6	M_7	A_6A_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_6	M_7	X_5
dim	2	2	2
Y_3	1	1	1
Y_4	1	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	R_5
dim		4	2
U_3	1	2	-1
U_4	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	R_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/4}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	A_6A_7
dim		4	4
S_3S_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6A_7	R_5
dim		4	2
T_3	1	2	-1
T_4	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_6A_7	R_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/4}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_5	X_5
dim		2	2
W_3	1	1	-1
W_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 113

Space group $P\bar{4}2_1m$, No. 113, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
dim		2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Λ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6M_7	A_6A_7
	dim	4	4
V_5V_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6M_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_6M_7	X_2X_3	X_4X_5
dim	4	2	2
Y_3Y_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6	Z_7	R_2R_3	R_4R_5
dim	2	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	A_6A_7
	dim	2	2	4
S_3	1	1	1	-2
S_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	A_6A_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6A_7	R_2R_3	R_4R_5
	dim	4	2	2
T_3T_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_6A_7	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_2R_3	R_4R_5	X_2X_3	X_4X_5
	dim	2	2	2	2
W_3W_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 114

Space group $P\bar{4}2_1c$, No. 114, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6Z_7
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	$M_6 M_7$
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$M_6 M_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$M_6 M_7$	$A_6 A_6$	$A_7 A_7$
	dim	4	4	4
$V_5 V_5$	4	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$M_6 M_7$	$A_6 A_6$	$A_7 A_7$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6M_7	X_2X_3	X_4X_5
dim		4	2	2
Y_3Y_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	R_2R_3	R_4R_5
dim		4	2	2
U_3	1	2	-2	0
U_4	1	2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	R_2R_3	R_4R_5	
	4	2	2	Bipartite graph
v^1	1	1	1	$H_{3/4}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6Z_7	A_6A_6	A_7A_7
dim		4	4	4
S_3S_4	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6Z_7	A_6A_6	A_7A_7	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_6A_6	A_7A_7	R_2R_3	R_4R_5
dim	4	4	2	2
T_3T_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_6A_6	A_7A_7	R_2R_3	R_4R_5	
	4	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	2	$H_{3/2}^1$
v^5	1	0	1	1	$H_{3/2}^1$
v^6	1	0	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	R_2R_3	R_4R_5	X_2X_3	X_4X_5
dim	2	2	2	2
W_3W_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 115

Space group $P\bar{4}m2$, No. 115, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Λ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	A_6	A_7
	dim	2	2	2	2
V_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_6	M_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6	M_7	X_5
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	R_5
dim		2	2	2
U_3	1	1	1	-1
U_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_6	Z_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6	Z_7	A_6	A_7
dim		2	2	2	2
S_3	1	1	1	-1	-1
S_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_6	Z_7	A_6	A_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6	A_7	R_5
dim		2	2	2
T_3	1	1	1	-1
T_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_6	A_7	R_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_5	X_5
dim		2	2
W_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5	X_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 116

Space group $P\bar{4}c2$, No. 116, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	$Z_6 Z_7$
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	M_6	M_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_6	M_7	A_6A_7
dim	2	2	4
V_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6	M_7	A_6A_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_6	M_7	X_5
dim	2	2	2
Y_3	1	1	1
Y_4	1	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6Z_7	R_5R_5
dim	4	4
U_3U_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6Z_7	A_6A_7
dim	4	4
S_3	1	2
S_4	1	2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_6Z_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_6A_7	R_5R_5
dim	4	4
T_3T_4	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_6A_7	R_5R_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_5R_5	X_5
dim		4	2
W_5	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_5R_5	X_5	
	4	2	Bipartite graph
v^1	1	2	$H_{3/2}^1$

SG 117

Space group $P\bar{4}b2$, No. 117, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
dim		2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Λ_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	M_6M_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	M_6M_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6M_7	A_6A_7
	dim	4	4
V_5	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_6M_7	A_6A_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_6M_7	X_2X_3	X_4X_5
dim	4	2	2
Y_3Y_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6	Z_7	R_2R_3	R_4R_5
dim	2	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	R_2R_3	R_4R_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6	Z_7	A_6A_7
dim	2	2	4
S_3	1	1	1
S_4	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6	Z_7	A_6A_7	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_6A_7	R_2R_3	R_4R_5
dim	4	2	2
T_3T_4	2	2	2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_6A_7	R_2R_3	R_4R_5	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_2R_3	R_4R_5	X_2X_3	X_4X_5
dim	2	2	2	2

W_2	1	1	0	-1	0
W_3	1	0	1	0	-1
W_4	1	1	0	-1	0
W_5	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_2R_3	R_4R_5	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 118

Space group $P\bar{4}n2$, No. 118, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
	dim	2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6Z_7
	dim	2	2	4
Λ_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$Z_6 Z_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	$M_6 M_7$
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	$M_6 M_7$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		$M_6 M_7$	A_6	A_7
	dim	4	2	2
V_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$M_6 M_7$	A_6	A_7	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_6M_7	X_2X_3	X_4X_5
dim	4	2	2
Y_3Y_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6M_7	X_2X_3	X_4X_5	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6Z_7	R_2R_3	R_4R_5
dim	4	2	2
U_3U_4	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_6Z_7	R_2R_3	R_4R_5	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6Z_7	A_6	A_7
dim	4	2	2
S_3	1	2	-1 -1
S_4	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$Z_6 Z_7$	A_6	A_7	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	1	2	0	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_6	A_7	$R_2 R_3$	$R_4 R_5$
dim	2	2	2	2
T_3	1	1	1	-2 0
T_4	1	1	1	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_6	A_7	$R_2 R_3$	$R_4 R_5$	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	$R_2 R_3$	$R_4 R_5$	$X_2 X_3$	$X_4 X_5$
dim	2	2	2	2
W_2	1	0	1	-1 0
W_3	1	0	1	0 -1
W_4	1	1	0	-1 0
W_5	1	1	0	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$R_2 R_3$	$R_4 R_5$	$X_2 X_3$	$X_4 X_5$	
	2	2	2	2	Bipartite graph
v^1	1	1	1	1	$H_{4/4}^2$

SG 119

Space group $I\bar{4}m2$, No. 119, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		$N_3 N_4$	P_5	P_6	P_7	P_8
	dim	2	1	1	1	1
Q_2	1	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	$N_3 N_4$	P_5	P_6	P_7	P_8	Bipartite graph
	2	1	1	1	1	
v^1	1	0	0	0	2	$H_{3/2}^1$
v^2	1	0	0	1	1	$H_{3/2}^1$
v^3	1	0	0	2	0	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$
v^5	1	0	1	1	0	$H_{3/2}^1$
v^6	1	0	2	0	0	$H_{3/2}^1$
v^7	1	1	0	0	1	$H_{3/2}^1$
v^8	1	1	0	1	0	$H_{3/2}^1$
v^9	1	1	1	0	0	$H_{3/2}^1$
v^{10}	1	2	0	0	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		X_5	P_5	P_6	P_7	P_8
	dim	2	1	1	1	1
W_3	1	1	-1	0	-1	0
W_4	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_5	P_5	P_6	P_7	P_8	Bipartite graph
	2	1	1	1	1	
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	Z_6	Z_7
	dim	2	2	2
Y_3	1	1	-1	-1
Y_4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	Z_6	Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 120

Space group $I\bar{4}c2$, No. 120, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
	dim	2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_3N_3	N_4N_4	P_5P_7	P_6P_8
	dim	2	2	2	2
Q_2Q_2	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	N_3N_3	N_4N_4	P_5P_7	P_6P_8	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_5	P_5P_7	P_6P_8
	dim	2	2	2
W_3	1	1	-2	0
W_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	P_5P_7	P_6P_8	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	Z_6	Z_7
dim		2	2	2
Y_3	1	1	-1	-1
Y_4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	Z_6	Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 121

Space group $I\bar{4}2m$, No. 121, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6	Z_7
dim		2	2	2	2
Σ_3	1	1	1	-1	-1
Σ_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Z_6	Z_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	X_5
	dim	2	2	2
Δ_3	1	1	1	-1
Δ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	X_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_3N_4	P_6	P_7
	dim	2	2	2
Q_3	1	1	-1	-1
Q_4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	N_3N_4	P_6	P_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W : $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_5	P_6	P_7
	dim	2	2	2
W_5	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	P_6	P_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	Z_6	Z_7
	dim	2	2	2
Y_3	1	1	-1	-1
Y_4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	Z_6	Z_7	
	2	2	2	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

SG 122

Space group $I\bar{4}2d$, No. 122, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Z_6Z_7
	dim	2	2	4
Σ_3	1	1	1	-2
Σ_4	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Z_6Z_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	X_2X_3	X_4X_5
dim		2	2	2	2
Δ_3	1	1	1	0	-2
Δ_4	1	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	X_2X_3	X_4X_5	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		N_3N_4	P_3	P_4	P_5	P_6	P_7
dim		2	1	1	1	1	2
Q_3	1	1	0	-1	-1	0	-1
Q_4	1	1	-1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	N_3N_4	P_3	P_4	P_5	P_6	P_7	
	2	1	1	1	1	2	Bipartite graph
v^1	1	0	0	0	0	1	$H_{2/2}^1$
v^2	1	0	0	1	1	0	$H_{3/2}^1$
v^3	1	0	1	0	1	0	$H_{3/2}^1$
v^4	1	1	0	1	0	0	$H_{3/2}^1$
v^5	1	1	1	0	0	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W : $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_2X_3	X_4X_5	P_3	P_4	P_5	P_6	P_7
	dim	2	2	1	1	1	1	2
W_2	1	1	0	-1	0	-1	0	0
W_3	1	0	1	0	0	0	0	-1
W_4	1	1	0	0	0	0	0	-1
W_5	1	0	1	0	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_2X_3	X_4X_5	P_3	P_4	P_5	P_6	P_7	
	2	2	1	1	1	1	2	Bipartite graph
v^1	1	1	0	0	1	1	1	$H_{5/4}^3$
v^2	1	1	0	1	1	0	1	$H_{5/4}^3$
v^3	1	1	1	0	0	1	1	$H_{5/4}^3$
v^4	1	1	1	1	0	0	1	$H_{5/4}^3$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_2X_3	X_4X_5	Z_6Z_7
dim	2	2	4
Y_3Y_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	Z_6Z_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

SG 123

Space group $P4/mmm$, No. 123, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-
	dim	2	2	2	2	2	2
Δ_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-
	dim	2	2	2	2	2	2	2	2
Λ_6	2	1	1	0	0	-1	-1	0	0
Λ_7	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-
dim	2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_6^+	M_6^-	M_7^+	M_7^-	A_6^+	A_6^-	A_7^+	A_7^-
dim	2	2	2	2	2	2	2	2
V_6	2	1	1	0	0	-1	-1	0
V_7	2	0	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_6^+	M_6^-	M_7^+	M_7^-	A_6^+	A_6^-	A_7^+	A_7^-	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-
	dim	2	2	2	2	2	2
Y_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_6^+	Z_6^-	Z_7^+	Z_7^-	R_5^+	R_5^-
	dim	2	2	2	2	2	2
U_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	R_5^+	R_5^-	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6^+	Z_6^-	Z_7^+	Z_7^-	A_6^+	A_6^-	A_7^+	A_7^-
dim	2	2	2	2	2	2	2	2
S_5	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	A_6^+	A_6^-	A_7^+	A_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_6^+	A_6^-	A_7^+	A_7^-	R_5^+	R_5^-
dim		2	2	2	2	2	2
T_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_6^+	A_6^-	A_7^+	A_7^-	R_5^+	R_5^-	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_5^+	R_5^-	X_5^+	X_5^-
dim		2	2	2	2
W_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_5^+	R_5^-	X_5^+	X_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 124

Space group $P4/mcc$, No. 124, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-
	dim	2	2	2	2	2	2
Δ_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8
	dim	2	2	2	2	4	4
Λ_6	2	1	1	0	0	-2	0
Λ_7	2	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-
dim	2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_6^+	M_6^-	M_7^+	M_7^-	$A_5 A_6$	$A_7 A_8$
dim	2	2	2	2	4	4
V_6	2	1	1	0	0	-2
V_7	2	0	0	1	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_6^+	M_6^-	M_7^+	M_7^-	A_5A_6	A_7A_8	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-
dim		2	2	2	2	2	2
Y_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_6	Z_7Z_8	R_3R_4
dim		4	4	4
U_2U_5	2	1	1	-1
U_3U_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5Z_6	Z_7Z_8	R_3R_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_6	Z_7Z_8	A_5A_6	A_7A_8
	dim	4	4	4	4
S_2S_5	2	1	1	-1	-1
S_3S_4	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5Z_6	Z_7Z_8	A_5A_6	A_7A_8	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_5A_6	A_7A_8	R_3R_4
	dim	4	4	4
T_2T_5	2	1	1	-1
T_3T_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_5A_6	A_7A_8	R_3R_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_3R_4	X_5^+	X_5^-
dim		4	2	2
W_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	R_3R_4	X_5^+	X_5^-	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

SG 125

Space group $P4/nbm$, No. 125, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim		2	2	2	2	4
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-
	dim	2	2	2	2	2	2	2	2
Λ_6	2	1	1	0	0	-1	-1	0	0
Λ_7	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	A_5
dim		4	4
V_6	2	1	-1
V_7	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		M_5	A_5	
		4	4	Bipartite graph
v^1	1	1		$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_5	$X_3 X_4$
dim		4	4
$Y_2 Y_5$	2	1	-1
$Y_3 Y_4$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		M_5	$X_3 X_4$	
		4	4	Bipartite graph
v^1	1	1		$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_6^+	Z_6^-	Z_7^+	Z_7^-	$R_3 R_4$
dim		2	2	2	2	4
U_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	R_3R_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_6^+	Z_6^-	Z_7^+	Z_7^-	A_5
	dim	2	2	2	2	4
S_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	A_5	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix



corep		A_5	R_3R_4
	dim	4	4
T_2T_5	2	1	-1
T_3T_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_5	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W : $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_3R_4	X_3X_4
	dim	4	4
W_2W_5	2	1	-1
W_3W_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 126

Space group $P4/nnc$, No. 126, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
	dim	2	2	2	2	4
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8
	dim	2	2	2	2	4	4
Λ_6	2	1	1	0	0	-2	0
Λ_7	2	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	A_5
	dim	4	4
V_6	2	1	-1
V_7	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	A_5	Bipartite graph
	4	4	
v^1	1	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_5	X_3X_4
	dim	4	4
Y_2Y_5	2	1	-1
Y_3Y_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_6	Z_7Z_8	R_3R_4
dim		4	4	4
U_2U_5	2	1	1	-1
U_3U_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5Z_6	Z_7Z_8	R_3R_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_6	Z_7Z_8	A_5
dim		4	4	4
S_2S_5	2	1	1	-1
S_3S_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5Z_6	Z_7Z_8	A_5	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5	R_3R_4
dim	4	4
T_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_5	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_3R_4	X_3X_4
dim	4	4
W_2W_5	2	-1
W_3W_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 127

Space group $P4/mbm$, No. 127, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim	2	2	2	2	4
Δ_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-
	dim	2	2	2	2	2	2	2	2
Λ_6	2	1	1	0	0	-1	-1	0	0
Λ_7	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+ M_7^+$	$M_6^- M_7^-$
dim	2	2	2	2		4	4
Σ_5	2	1	1	1	1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+ M_7^+$	$M_6^- M_7^-$	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	0	2	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	1	$H_{3/2}^1$
v^4	0	0	1	1	1	0	$H_{3/2}^1$
v^5	0	0	2	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	0	$H_{3/2}^1$
v^9	0	1	1	0	0	1	$H_{3/2}^1$
v^{10}	0	1	1	0	1	0	$H_{3/2}^1$
v^{11}	0	2	0	0	0	1	$H_{3/2}^1$
v^{12}	0	2	0	0	1	0	$H_{3/2}^1$
v^{13}	1	0	0	1	0	1	$H_{3/2}^1$
v^{14}	1	0	0	1	1	0	$H_{3/2}^1$
v^{15}	1	0	1	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	1	0	$H_{3/2}^1$
v^{17}	1	1	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	1	0	$H_{3/2}^1$
v^{19}	2	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		$M_6^+ M_7^+$	$M_6^- M_7^-$	$A_6^+ A_7^+$	$A_6^- A_7^-$
dim	4	4		4	4
$V_6 V_7$	4	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

$M_6^+ M_7^+$	$M_6^- M_7^-$	$A_6^+ A_7^+$	$A_6^- A_7^-$	
4	4	4	4	Bipartite graph

v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$
dim	4	4	4
$Y_5 Y_5$	4	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_6^+	Z_6^-	Z_7^+	Z_7^-	$R_3 R_4$
dim	2	2	2	2	4
U_5	2	1	1	1	1
					-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	$R_3 R_4$	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6^+	Z_6^-	Z_7^+	Z_7^-	$A_6^+ A_7^+$	$A_6^- A_7^-$
dim	2	2	2	2	4	4
S_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	$A_6^+ A_7^+$	$A_6^- A_7^-$	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	0	2	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	1	$H_{3/2}^1$
v^4	0	0	1	1	1	0	$H_{3/2}^1$
v^5	0	0	2	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	0	$H_{3/2}^1$
v^9	0	1	1	0	0	1	$H_{3/2}^1$
v^{10}	0	1	1	0	1	0	$H_{3/2}^1$
v^{11}	0	2	0	0	0	1	$H_{3/2}^1$
v^{12}	0	2	0	0	1	0	$H_{3/2}^1$
v^{13}	1	0	0	1	0	1	$H_{3/2}^1$
v^{14}	1	0	0	1	1	0	$H_{3/2}^1$
v^{15}	1	0	1	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	1	0	$H_{3/2}^1$
v^{17}	1	1	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	1	0	$H_{3/2}^1$
v^{19}	2	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_6^+ A_7^+$	$A_6^- A_7^-$	$R_3 R_4$
dim	4	4	4
$T_5 T_5$	4	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$A_6^+ A_7^+$	$A_6^- A_7^-$	$R_3 R_4$	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		$R_3 R_4$	$X_3 X_4$
	dim	4	4
$W_2 W_3$	2	1	-1
$W_4 W_5$	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$R_3 R_4$	$X_3 X_4$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 128

Space group $P4/mnc$, No. 128, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$X_3 X_4$
	dim	2	2	2	2	4
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$X_3 X_4$	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$

v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8
	dim	2	2	2	2	4	4
Λ_6	2	1	1	0	0	-2	0
Λ_7	2	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+M_7^+$	$M_6^-M_7^-$
	dim	2	2	2	2	4	4
Σ_5	2	1	1	1	1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+M_7^+$	$M_6^-M_7^-$	
	2	2	2	2	4	4	Bipartite graph

v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	0	2	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	1	$H_{3/2}^1$
v^4	0	0	1	1	1	0	$H_{3/2}^1$
v^5	0	0	2	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	0	$H_{3/2}^1$
v^9	0	1	1	0	0	1	$H_{3/2}^1$
v^{10}	0	1	1	0	1	0	$H_{3/2}^1$
v^{11}	0	2	0	0	0	1	$H_{3/2}^1$
v^{12}	0	2	0	0	1	0	$H_{3/2}^1$
v^{13}	1	0	0	1	0	1	$H_{3/2}^1$
v^{14}	1	0	0	1	1	0	$H_{3/2}^1$
v^{15}	1	0	1	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	1	0	$H_{3/2}^1$
v^{17}	1	1	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	1	0	$H_{3/2}^1$
v^{19}	2	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_6^+ M_7^+$	$M_6^- M_7^-$	$A_5 A_6$	$A_7 A_8$
dim	4	4	4	4
$V_6 V_7$	4	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_6^+ M_7^+$	$M_6^- M_7^-$	$A_5 A_6$	$A_7 A_8$	Bipartite graph
	4	4	4	4	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$
dim	4	4	4
$Y_5 Y_5$	4	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$Z_5 Z_6$	$Z_7 Z_8$	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$
dim	4	4	2	2	2	2
$U_2 U_5$	2	1	1	0	0	-1
$U_3 U_4$	2	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$Z_5 Z_6$	$Z_7 Z_8$	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	
	4	4	2	2	2	2	Bipartite graph
v^1	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	1	0	$H_{3/2}^1$
v^5	1	0	0	1	0	1	$H_{3/2}^1$
v^6	1	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$Z_5 Z_6$	$Z_7 Z_8$	$A_5 A_6$	$A_7 A_8$
dim	4	4	4	4
$S_2 S_5$	2	1	1	-1
$S_3 S_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$Z_5 Z_6$	$Z_7 Z_8$	$A_5 A_6$	$A_7 A_8$	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$A_5 A_6$	$A_7 A_8$	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	
dim	4	4	2	2	2	2	
$T_2 T_4$	2	1	1	0	-1	-1	0
$T_3 T_5$	2	1	1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$A_5 A_6$	$A_7 A_8$	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	
	4	4	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	1	0	1	$H_{3/2}^1$
v^3	0	1	1	0	1	0	$H_{3/2}^1$
v^4	0	1	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	1	$H_{3/2}^1$
v^7	1	0	1	0	1	0	$H_{3/2}^1$
v^8	1	0	1	1	0	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	$X_3 X_4$	
dim	2	2	2	2	4	
$W_2 W_3$	2	1	0	0	1	-1
$W_4 W_5$	2	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	$X_3 X_4$	
	2	2	2	2	4	Bipartite graph
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

SG 129

Space group $P4/nmm$, No. 129, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$X_3 X_4$
dim	2	2	2	2	4
Δ_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$X_3 X_4$	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-
	dim	2	2	2	2	2	2	2	2
Λ_6	2	1	1	0	0	-1	-1	0	0
Λ_7	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_5	A_5
dim	4	4
V_6V_7	4	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

M_5	A_5	Bipartite graph
4	4	
v^1	1	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_5	X_3X_4
dim	4	4
Y_2Y_5	2	-1
Y_3Y_4	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

M_5	X_3X_4	Bipartite graph
4	4	
v^1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_6^+	Z_6^-	Z_7^+	Z_7^-	R_3R_4
dim	2	2	2	2	4
U_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	R_3R_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Z_6^+	Z_6^-	Z_7^+	Z_7^-	A_5
	dim	2	2	2	2	4
S_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_6^+	Z_6^-	Z_7^+	Z_7^-	A_5	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix



corep	A_5	R_3R_4
dim	4	4
T_2T_5	2	1 -1
T_3T_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

A_5	R_3R_4	
4	4	Bipartite graph
v^1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_3R_4	X_3X_4
dim	4	4
W_5W_5	4	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

R_3R_4	X_3X_4	
4	4	Bipartite graph
v^1	1	$H_{2/1}^1$

SG 130

Space group $P4/ncc$, No. 130, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim	2	2	2	2	4
Δ_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8
	dim	2	2	2	2	4	4
Λ_6	2	1	1	0	0	-2	0
Λ_7	2	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5Z_6	Z_7Z_8	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	$H_{3/2}^1$
v^6	2	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_5	$A_5 A_5$
dim	4	8
$V_6 V_7$	4	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	$A_5 A_5$	
	4	8	Bipartite graph
v^1	2	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_5	$X_3 X_4$
dim	4	4
$Y_2 Y_5$	2	-1
$Y_3 Y_4$	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_6	Z_7Z_8	R_3R_3	R_4R_4
	dim	4	4	4	4
U_2U_5	2	1	1	-2	0
U_3U_4	2	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5Z_6	Z_7Z_8	R_3R_3	R_4R_4	
	4	4	4	4	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5Z_6	Z_7Z_8	A_5A_5
	dim	4	4	8
S_2S_5	2	1	1	-2
S_3S_4	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5Z_6	Z_7Z_8	A_5A_5	
	4	4	8	Bipartite graph
v^1	0	2	1	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	2	0	1	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5A_5	R_3R_3	R_4R_4
dim	8	4	4
T_5T_5	4	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_5A_5	R_3R_3	R_4R_4	
	8	4	4	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_3R_3	R_4R_4	X_3X_4
dim	4	4	4
W_5W_5	4	1	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	R_3R_3	R_4R_4	X_3X_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

SG 131

Space group $P4_2/mmc$, No. 131, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-
dim		2	2	2	2	2	2
Δ_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
dim		2	2	2	2	4
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-
dim	2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0)$, $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_6^+	M_6^-	M_7^+	M_7^-	A_5
dim	2	2	2	2	4
V_6	2	1	1	0	0
V_7	2	0	0	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_6^+	M_6^-	M_7^+	M_7^-	A_5	Bipartite graph
	2	2	2	2	4	
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-
	dim	2	2	2	2	2	2
Y_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	R_5^+	R_5^-
	dim	4	2	2
U_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Z_5	R_5^+	R_5^-	Bipartite graph
	4	2	2	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	A_5
	dim	4	4
S_2S_3	2	1	-1
S_4S_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	A_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_5	R_5^+	R_5^-
	dim	4	2	2
T_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_5	R_5^+	R_5^-	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_5^+	R_5^-	X_5^+	X_5^-
	dim	2	2	2	2
W_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	R_5^+	R_5^-	X_5^+	X_5^-	
	2	2	2	2	Bipartite graph

v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

SG 132

Space group $P4_2/mcm$, No. 132, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-
	dim	2	2	2	2	2	2
Δ_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
	dim	2	2	2	2	4
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	Bipartite graph
	2	2	2	2	4	
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-
dim	2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_6^+	M_6^-	M_7^+	M_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_6^+	M_6^-	M_7^+	M_7^-	A_5
	dim	2	2	2	2	4
V_6	2	1	1	0	0	-1
V_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_6^+	M_6^-	M_7^+	M_7^-	A_5	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-
	dim	2	2	2	2	2	2
Y_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_6^+	M_6^-	M_7^+	M_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	R_3R_4
dim	4	4
U_2U_5	2	1 -1
U_3U_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	A_5
dim	4	4
S_5	2	2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	A_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5	R_3R_4
dim	4	4
T_2T_5	2	1 -1
T_3T_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_5	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep		R_3R_4	X_5^+	X_5^-
dim		4	2	2
W_5	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	R_3R_4	X_5^+	X_5^-	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

SG 133

Space group $P4_2/nbc$, No. 133, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim		2	2	2	2	4
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
	dim	2	2	2	2	4
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	A_5A_6	A_7A_8
	dim	4	4	4
V_6	2	1	0	-2
V_7	2	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	A_5A_6	A_7A_8	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		M_5	X_3X_4
	dim	4	4
Y_2Y_5	2	1	-1
Y_3Y_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Z_5	R_3R_4
	dim	4	4
U_5	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	A_5A_6	A_7A_8
dim	4	4	4
S_2S_3	2	1	-1
S_4S_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5	A_5A_6	A_7A_8	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5A_6	A_7A_8	R_3R_4
dim	4	4	4
T_2T_5	2	1	1
T_3T_4	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_5A_6	A_7A_8	R_3R_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_3R_4	X_3X_4
dim	4	4
W_2W_5	2	1
W_3W_4	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 134

Space group $P4_2/nmm$, No. 134, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim		2	2	2	2	4
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
dim		2	2	2	2	4
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	Bipartite graph
	2	2	2	2	4	
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
dim	2	2	2	2	4
Σ_5	2	1	1	1	1
					-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_5	A_6^+	A_6^-	A_7^+	A_7^-
dim	4	2	2	2	2
V_6	2	1	0	0	-1
V_7	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_5	A_6^+	A_6^-	A_7^+	A_7^-	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	M_5	X_3X_4
dim	4	4
Y_2Y_5	2	1 -1
Y_3Y_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_5	R_3R_4
dim	4	4
U_2U_5	2	1 -1
U_3U_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	R_3R_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	A_6^+	A_6^-	A_7^+	A_7^-
dim	4	2	2	2	2
S_5	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Z_5	A_6^+	A_6^-	A_7^+	A_7^-	Bipartite graph
	4	2	2	2	2	
v^1	1	0	0	0	2	$H_{3/2}^1$
v^2	1	0	0	1	1	$H_{3/2}^1$
v^3	1	0	0	2	0	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$
v^5	1	0	1	1	0	$H_{3/2}^1$
v^6	1	0	2	0	0	$H_{3/2}^1$
v^7	1	1	0	0	1	$H_{3/2}^1$
v^8	1	1	0	1	0	$H_{3/2}^1$
v^9	1	1	1	0	0	$H_{3/2}^1$
v^{10}	1	2	0	0	0	$H_{3/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_6^+	A_6^-	A_7^+	A_7^-	R_3R_4
dim	2	2	2	2	4
T_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	A_6^+	A_6^-	A_7^+	A_7^-	R_3R_4	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$

v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$

A matrix

corep		R_3R_4	X_3X_4
	dim	4	4
W_2W_5	2	1	-1
W_3W_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 135

Space group $P4_2/mbc$, No. 135, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
	dim	2	2	2	2	4
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$

v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
	dim	2	2	2	2	4
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+ M_7^+$	$M_6^- M_7^-$
	dim	2	2	2	2	4	4
Σ_5	2	1	1	1	1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+ M_7^+$	$M_6^- M_7^-$	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	0	2	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	1	$H_{3/2}^1$
v^4	0	0	1	1	1	0	$H_{3/2}^1$
v^5	0	0	2	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	$H_{3/2}^1$

v^8	0	1	0	1	1	0	$H_{3/2}^1$
v^9	0	1	1	0	0	1	$H_{3/2}^1$
v^{10}	0	1	1	0	1	0	$H_{3/2}^1$
v^{11}	0	2	0	0	0	1	$H_{3/2}^1$
v^{12}	0	2	0	0	1	0	$H_{3/2}^1$
v^{13}	1	0	0	1	0	1	$H_{3/2}^1$
v^{14}	1	0	0	1	1	0	$H_{3/2}^1$
v^{15}	1	0	1	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	1	0	$H_{3/2}^1$
v^{17}	1	1	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	1	0	$H_{3/2}^1$
v^{19}	2	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$M_6^+ M_7^+$	$M_6^- M_7^-$	$A_5 A_5$
dim	4	4	8
$V_6 V_7$	4	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$M_6^+ M_7^+$	$M_6^- M_7^-$	$A_5 A_5$	Bipartite graph
	4	4	8	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$
dim	4	4	4
$Y_5 Y_5$	4	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	$R_3 R_4$	
dim	4	4	
U_5	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	$R_3 R_4$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	$A_5 A_5$	
dim	4	8	
$S_2 S_3$	2	1	-2
$S_4 S_5$	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	$A_5 A_5$	
	4	8	Bipartite graph
v^1	2	1	$H_{3/4}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5A_5	R_3R_4
dim	8	4
T_5T_5	4	2
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	A_5A_5	R_3R_4	
	8	4	Bipartite graph
v^1	1	2	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_3R_4	X_3X_4
dim	4	4
W_2W_3	2	1
		-1
W_4W_5	2	1
		-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 136

Space group $P4_2/mnm$, No. 136, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim	2	2	2	2	4
Δ_5	2	1	1	1	1
					-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
	dim	2	2	2	2	4
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+M_7^+$	$M_6^-M_7^-$
	dim	2	2	2	2	4	4
Σ_5	2	1	1	1	1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	$M_6^+ M_7^+$	$M_6^- M_7^-$	Bipartite graph
	2	2	2	2	4	4	
v^1	0	0	0	2	0	1	$H_{3/2}^1$
v^2	0	0	0	2	1	0	$H_{3/2}^1$
v^3	0	0	1	1	0	1	$H_{3/2}^1$
v^4	0	0	1	1	1	0	$H_{3/2}^1$
v^5	0	0	2	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	$H_{3/2}^1$
v^8	0	1	0	1	1	0	$H_{3/2}^1$
v^9	0	1	1	0	0	1	$H_{3/2}^1$
v^{10}	0	1	1	0	1	0	$H_{3/2}^1$
v^{11}	0	2	0	0	0	1	$H_{3/2}^1$
v^{12}	0	2	0	0	1	0	$H_{3/2}^1$
v^{13}	1	0	0	1	0	1	$H_{3/2}^1$
v^{14}	1	0	0	1	1	0	$H_{3/2}^1$
v^{15}	1	0	1	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	1	0	$H_{3/2}^1$
v^{17}	1	1	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	1	0	$H_{3/2}^1$
v^{19}	2	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	1	0	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$M_6^+ M_7^+$	$M_6^- M_7^-$	A_5
dim	4	4	4
$V_6 V_7$	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$M_6^+ M_7^+$	$M_6^- M_7^-$	A_5	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$
dim	4	4	4
$Y_5 Y_5$	4	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$M_6^+ M_7^+$	$M_6^- M_7^-$	$X_3 X_4$	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_5	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$
dim	4	2	2	2	2
$U_2 U_5$	2	1	0	0	-1
$U_3 U_4$	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Z_5	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	1	0	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Z_5	A_5
dim	4	4
S_5	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	A_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	
dim	4	2	2	2	2	
$T_2 T_5$	2	1	0	-1	-1	0
$T_3 T_4$	2	1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_5	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	Bipartite graph
	4	2	2	2	2	
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	0	1	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	$X_3 X_4$	
dim	2	2	2	2	4	
$W_2 W_3$	2	1	0	0	1	-1
$W_4 W_5$	2	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$R_2^+ R_3^+$	$R_2^- R_3^-$	$R_4^+ R_5^+$	$R_4^- R_5^-$	$X_3 X_4$	Bipartite graph
	2	2	2	2	4	
v^1	0	0	1	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	1	0	1	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$

SG 137

Space group $P4_2/nmc$, No. 137, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
dim	2	2	2	2	4	
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
dim	2	2	2	2	4	
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
dim	2	2	2	2	4
Σ_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V : $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_5	A_5A_6	A_7A_8
dim	4	4	4
V_6V_7	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	A_5A_6	A_7A_8	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y : $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix



corep	M_5	$X_3 X_4$
dim	4	4
$Y_2 Y_4$	2	1 -1
$Y_3 Y_5$	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	$X_3 X_4$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	$R_3 R_4$
dim	4	4
U_5	2	2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	$R_3 R_4$	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	$A_5 A_6$	$A_7 A_8$
dim	4	4	4
$S_2 S_3$	2	1 -1	-1
$S_4 S_5$	2	1 -1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5	$A_5 A_6$	$A_7 A_8$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_5A_6	A_7A_8	R_3R_4
dim	4	4	4
T_2T_4	2	1	1
T_3T_5	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_5A_6	A_7A_8	R_3R_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

W

High symmetry line $W(0, \frac{1}{2}, u)$, high symmetry points at both ends of W: $R(0, \frac{1}{2}, \frac{1}{2}), X(0, \frac{1}{2}, 0)$
A matrix

corep	R_3R_4	X_3X_4
dim	4	4
W_5W_5	4	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	R_3R_4	X_3X_4	
	4	4	Bipartite graph
v^1	1	1	$H_{2/1}^1$

SG 138

Space group $P4_2/nm$, No. 138, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim	2	2	2	2	4
Δ_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(0, 0, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), Z(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
	dim	2	2	2	2	4
Λ_6	2	1	1	0	0	-1
Λ_7	2	0	0	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	M_5	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

V

High symmetry line $V(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of V: $M(\frac{1}{2}, \frac{1}{2}, 0), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_5	$A_6^+ A_7^+$	$A_6^- A_7^-$
dim	4	4	4
$V_6 V_7$	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	$A_6^+ A_7^+$	$A_6^- A_7^-$	Bipartite graph
	4	4	4	
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Y: $M(\frac{1}{2}, \frac{1}{2}, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep	M_5	$X_3 X_4$
dim	4	4
$Y_2 Y_4$	2	1
$Y_3 Y_5$	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	$X_3 X_4$	Bipartite graph
	4	4	
v^1	1	1	$H_{2/2}^1$

U

High symmetry line $U(0, u, \frac{1}{2})$, high symmetry points at both ends of U: $Z(0, 0, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	R_3R_3	R_4R_4
dim	4	4	4
U_2U_5	2	1	-2 0
U_3U_4	2	1	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Z_5	R_3R_3	R_4R_4	
	4	4	4	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

S

High symmetry line $S(u, u, \frac{1}{2})$, high symmetry points at both ends of S: $Z(0, 0, \frac{1}{2}), A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Z_5	$A_6^+A_7^+$	$A_6^-A_7^-$
dim	4	4	4
S_5	2	2	-2 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_5	$A_6^+A_7^+$	$A_6^-A_7^-$	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	1	0	$H_{2/2}^1$

T

High symmetry line $T(u, \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of T: $A(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), R(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_6^+A_7^+$	$A_6^-A_7^-$	R_3R_3	R_4R_4
dim	4	4	4	4
T_5T_5	4	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-
	dim	2	2	2	2	2	2
Δ_5	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	$N_3^+ N_4^+$	$N_3^- N_4^-$	P_6	P_7
dim	2	2	2	2
$Q_3 Q_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$N_3^+ N_4^+$	$N_3^- N_4^-$	P_6	P_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	X_5^+	X_5^-	P_6	P_7
dim	2	2	2	2
W_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_5^+	X_5^-	P_6	P_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_5^+	X_5^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-
dim	2	2	2	2	2	2
Y_5	2	1	1	-1	-1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_5^+	X_5^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	Bipartite graph
	2	2	2	2	2	2	
v^1	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	1	0	0	0	$H_{2/1}^1$

SG 140

Space group $I4/mcm$, No. 140, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-
dim	2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$

$$\underline{v^{16} \mid 1 \ 0 \ 0 \ 0 \mid 1 \ 0 \ 0 \ 0 \mid H_{2/1}^1}$$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-
dim	2	2	2	2	2	2
Δ_5	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_5^+	X_5^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	N_2N_2	P_6P_7
dim	4	4
Q_3Q_3	2	1 -1
Q_4Q_4	2	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	N_2N_2	P_6P_7	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	X_5^+	X_5^-	P_6P_7	
dim	2	2	4	
W_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_5^+	X_5^-	P_6P_7	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_5^+	X_5^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	
dim	2	2	2	2	2	2	
Y_5	2	1	1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_5^+	X_5^-	Z_6^+	Z_6^-	Z_7^+	Z_7^-	
	2	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	1	1	0	0	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	$H_{2/1}^1$
v^6	1	0	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	1	0	0	0	$H_{2/1}^1$

SG 141

Space group $I4_1/amd$, No. 141, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
	dim	2	2	2	2	4
Δ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$

v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q: $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	$N_3^+ N_4^+$	$N_3^- N_4^-$	$P_3 P_4$	$P_5 P_6$	P_7	
dim	2	2	2	2	2	
$Q_3 Q_4$	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$N_3^+ N_4^+$	$N_3^- N_4^-$	$P_3 P_4$	$P_5 P_6$	P_7	Bipartite graph
	2	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W: $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep	$X_3 X_4$	$P_3 P_4$	$P_5 P_6$	P_7	
dim	4	2	2	2	
$W_2 W_3$	2	1	0	0	-1
$W_4 W_5$	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$X_3 X_4$	$P_3 P_4$	$P_5 P_6$	P_7	Bipartite graph
	4	2	2	2	
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	1	$H_{3/2}^1$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	Z_5
	dim	4	4
Y_2Y_3	2	1	-1
Y_4Y_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	Z_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 142

Space group $I4_1/acd$, No. 142, with SOC

Σ

High symmetry line $\Sigma(-u, u, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5
	dim	2	2	2	2	4
Σ_5	2	1	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Z_5	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, 0, \frac{1}{2})$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4
dim	2	2	2	2	4
Δ_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	X_3X_4	
	2	2	2	2	4	Bipartite graph
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(u, \frac{1}{2} - u, u)$, high symmetry points at both ends of Q : $N(0, \frac{1}{2}, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	N_2N_2	P_3P_6	P_4P_5	P_7P_7
dim	4	2	2	4
Q_3Q_3	2	1	0	-1
Q_4Q_4	2	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	N_2N_2	P_3P_6	P_4P_5	P_7P_7	
	4	2	2	4	Bipartite graph
v^1	1	0	0	1	$H_{2/2}^1$
v^2	1	1	1	0	$H_{3/2}^1$

W

High symmetry line $W(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of W : $X(0, 0, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		X_3X_4	P_3P_6	P_4P_5	P_7P_7
	dim	4	2	2	4
W_2W_3	2	1	0	0	-2
W_4W_5	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_3X_4	P_3P_6	P_4P_5	P_7P_7	
	4	2	2	4	Bipartite graph
v^1	2	0	2	1	$H_{5/4}^3$
v^2	2	1	1	1	$H_{5/4}^3$
v^3	2	2	0	1	$H_{5/4}^3$

Y

High symmetry line $Y(-u, u, \frac{1}{2})$, high symmetry points at both ends of Y: $X(0, 0, \frac{1}{2}), Z(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_3X_4	Z_5
	dim	4	4
Y_2Y_3	2	1	-1
Y_4Y_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	Z_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

SG 143

Space group $P3$, No. 143, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	A_4A_5	A_6A_6
	dim	2	2	2	2
Δ_4	1	1	0	-1	0
Δ_5	1	1	0	-1	0
Δ_6	1	0	2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	A_4A_5	A_6A_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	K_4	K_5	K_6
	dim	2	2	1	1	1
T_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	K_4	K_5	K_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	M_2M_2
	dim	2	2	2
Σ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	M_2M_2	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_2M_2	L_2L_2
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_2M_2	L_2L_2	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_2M_2	K_4	K_5	K_6
dim		2	1	1	1
T'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_2M_2	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4A_5	A_6A_6	H_4	H_5	H_6
	dim	2	2	1	1	1
S_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4A_5	A_6A_6	H_4	H_5	H_6	Bipartite graph
	2	2	1	1	1	
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_4A_5	A_6A_6	L_2L_2
	dim	2	2	2
R_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_5	A_6A_6	L_2L_2	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_2L_2	H_4	H_5	H_6
	dim	2	1	1	1
S'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_2L_2	H_4	H_5	H_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

SG 144

Space group $P3_1$, No. 144, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	A_4A_5	A_6A_6
	dim	2	2	2	2
Δ_4	1	1	0	0	-2
Δ_5	1	1	0	-1	0
Δ_6	1	0	2	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	A_4A_5	A_6A_6	
	2	2	2	2	Bipartite graph
v^1	2	1	2	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	K_4	K_5	K_6
dim		2	2	1	1	1
T_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	K_4	K_5	K_6	Bipartite graph
	2	2	1	1	1	
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	M_2M_2
dim		2	2	2
Σ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	M_2M_2	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_2M_2	L_2L_2
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_2M_2	L_2L_2	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_2M_2	K_4	K_5	K_6
dim		2	1	1	1
T'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_2M_2	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4A_5	A_6A_6	H_4	H_5	H_6
dim		2	2	1	1	1
S_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4A_5	A_6A_6	H_4	H_5	H_6	Bipartite graph
	2	2	1	1	1	
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_4A_5	A_6A_6	L_2L_2
dim	2	2	2
R_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_5	A_6A_6	L_2L_2	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_2L_2	H_4	H_5	H_6
dim	2	1	1	1
S'_2	1	2	-1	-1
			-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_2L_2	H_4	H_5	H_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

SG 145

Space group $P3_2$, No. 145, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	A_4A_5	A_6A_6
	dim	2	2	2	2
Δ_4	1	1	0	-1	0
Δ_5	1	1	0	0	-2
Δ_6	1	0	2	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	A_4A_5	A_6A_6	
	2	2	2	2	Bipartite graph
v^1	2	1	2	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	K_4	K_5	K_6
	dim	2	2	1	1	1
T_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	K_4	K_5	K_6	Bipartite graph
	2	2	1	1	1	
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	M_2M_2
dim	2	2	2
Σ_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	M_2M_2	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_2M_2	L_2L_2
dim	2	2
U_2	1	2
		-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_2M_2	L_2L_2	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_2M_2	K_4	K_5	K_6
dim		2	1	1	1
T'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_2M_2	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4A_5	A_6A_6	H_4	H_5	H_6
dim		2	2	1	1	1
S_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4A_5	A_6A_6	H_4	H_5	H_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$

v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_4A_5	A_6A_6	L_2L_2
dim		2	2	2
R_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_5	A_6A_6	L_2L_2	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_2L_2	H_4	H_5	H_6
dim		2	1	1	1
S'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_2L_2	H_4	H_5	H_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

SG 146

Space group $R3$, No. 146, with SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$

A matrix

corep		$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	F_2F_2
	dim	2	2	2
Σ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_4\Gamma_5$	$\Gamma_6\Gamma_6$	F_2F_2	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y : $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep		Z_4Z_5	Z_6Z_6	L_2L_2
	dim	2	2	2
Y_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_4Z_5	Z_6Z_6	L_2L_2	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

SG 147

Space group $P\bar{3}$, No. 147, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep	$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	$A_4^+A_5^+$	$A_4^-A_5^-$	$A_6^+A_6^+$	$A_6^-A_6^-$	
dim	2	2	2	2	2	2	2	2	
$\Delta_4\Delta_5$	2	1	1	0	0	-1	-1	0	0
$\Delta_6\Delta_6$	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	$A_4^+A_5^+$	$A_4^-A_5^-$	$A_6^+A_6^+$	$A_6^-A_6^-$	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	K_4K_5	K_6K_6	
dim	2	2	2	2	2	2	
T_2T_2	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	K_4K_5	K_6K_6	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	$M_2^+M_2^+$	$M_2^-M_2^-$
dim	2	2	2	2	2	2
$\Sigma_2\Sigma_2$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	$M_2^+M_2^+$	$M_2^-M_2^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_2^+M_2^+$	$M_2^-M_2^-$	$L_2^+L_2^+$	$L_2^-L_2^-$
dim	2	2	2	2
U_2U_2	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_2^+M_2^+$	$M_2^-M_2^-$	$L_2^+L_2^+$	$L_2^-L_2^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	$M_2^+ M_2^+$	$M_2^- M_2^-$	$K_4 K_5$	$K_6 K_6$
dim	2	2	2	2
$T_2' T_2'$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_2^+ M_2^+$	$M_2^- M_2^-$	$K_4 K_5$	$K_6 K_6$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	$A_4^+ A_5^+$	$A_4^- A_5^-$	$A_6^+ A_6^+$	$A_6^- A_6^-$	$H_4 H_5$	$H_6 H_6$
dim	2	2	2	2	2	2
$S_2 S_2$	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$A_4^+ A_5^+$	$A_4^- A_5^-$	$A_6^+ A_6^+$	$A_6^- A_6^-$	$H_4 H_5$	$H_6 H_6$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$A_4^+ A_5^+$	$A_4^- A_5^-$	$A_6^+ A_6^+$	$A_6^- A_6^-$	$L_2^+ L_2^+$	$L_2^- L_2^-$
dim	2	2	2	2	2	2
$R_2 R_2$	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$A_4^+ A_5^+$	$A_4^- A_5^-$	$A_6^+ A_6^+$	$A_6^- A_6^-$	$L_2^+ L_2^+$	$L_2^- L_2^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	$L_2^+ L_2^+$	$L_2^- L_2^-$	$H_4 H_5$	$H_6 H_6$
dim	2	2	2	2
$S'_2 S'_2$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$L_2^+ L_2^+$	$L_2^- L_2^-$	$H_4 H_5$	$H_6 H_6$	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4K_5	K_6K_6	H_4H_5	H_6H_6
	dim	2	2	2	2
P_4P_5	2	1	0	-1	0
P_6P_6	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4K_5	K_6K_6	H_4H_5	H_6H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 148

Space group $R\bar{3}$, No. 148, with SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$

A matrix

corep		$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	$F_2^+F_2^+$	$F_2^-F_2^-$
	dim	2	2	2	2	2	2
$\Sigma_2\Sigma_2$	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$\Gamma_4^+\Gamma_5^+$	$\Gamma_4^-\Gamma_5^-$	$\Gamma_6^+\Gamma_6^+$	$\Gamma_6^-\Gamma_6^-$	$F_2^+F_2^+$	$F_2^-F_2^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y : $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep	$Z_4^+ Z_5^+$	$Z_4^- Z_5^-$	$Z_6^+ Z_6^+$	$Z_6^- Z_6^-$	$L_2^+ L_2^+$	$L_2^- L_2^-$
dim	2	2	2	2	2	2
$Y_2 Y_2$	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$Z_4^+ Z_5^+$	$Z_4^- Z_5^-$	$Z_6^+ Z_6^+$	$Z_6^- Z_6^-$	$L_2^+ L_2^+$	$L_2^- L_2^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

SG 149

Space group $P312$, No. 149, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep	Γ_4	$\Gamma_5 \Gamma_6$	A_4	$A_5 A_6$
dim	2	2	2	2
Δ_4	1	1 0	-1	0
Δ_5	1	1 0	-1	0
Δ_6	1	0 2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5 \Gamma_6$	A_4	$A_5 A_6$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
	dim	2	2	1	1	1
T_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	Bipartite graph
	2	2	1	1	1	
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
	dim	2	2	2
Σ_3	1	1	1	-1
Σ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_4
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_3M_4	K_4	K_5	K_6
dim		2	1	1	1
T'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_3M_4	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4	A_5A_6	H_4	H_5	H_6
dim		2	2	1	1	1
S_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_4	A_5A_6	L_3L_4
	dim	2	2	2
R_3	1	1	1	-1
R_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_3L_4	H_4	H_5	H_6
	dim	2	1	1	1
S'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_3L_4	H_4	H_5	H_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

SG 150

Space group $P321$, No. 150, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6
	dim	2	2	2	2
Δ_4	1	1	0	-1	0
Δ_5	1	1	0	-1	0
Δ_6	1	0	2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
	dim	2	2	2	1	1
T_3	1	1	1	-1	0	-1
T_4	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
dim	2	2	2
Σ_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_3M_4	L_3L_4
dim	2	2
U_2	1	2
		-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4	A_5A_6	H_4	H_5	H_6
	dim	2	2	2	1	1
S_3	1	1	1	-1	0	-1
S_4	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_4	A_5A_6	L_3L_4
	dim	2	2	2
R_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4	H_5	H_6
dim		2	2	1	1
S'_3	1	1	-1	-1	0
S'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
dim		2	1	1	2	1	1
P_4	1	1	0	0	-1	0	0
P_5	1	1	0	0	-1	0	0
P_6	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/2}^1$

SG 151

Space group $P3_112$, No. 151, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6
dim		2	2	2	2
Δ_4	1	1	0	0	-2
Δ_5	1	1	0	-1	0
Δ_6	1	0	2	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6	
	2	2	2	2	Bipartite graph
v^1	2	1	2	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
dim		2	2	1	1	1
T_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
dim		2	2	2
Σ_3	1	1	1	-1
Σ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

		Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
		2	2	2	Bipartite graph
v^1	0	1		1	$H_{2/2}^1$
v^2	1	0		1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_4
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		M_3M_4	L_3L_4	
		2	2	Bipartite graph
v^1	1	1		$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_4	K_5	K_6
dim		2	1	1	1
T'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_3M_4	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_4	A_5A_6	H_4	H_5	H_6
dim		2	2	1	1	1
S_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix



corep		A_4	A_5A_6	L_3L_4
	dim	2	2	2
R_3	1	1	1	-1
R_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4	H_5	H_6
	dim	2	1	1	1
S'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_3L_4	H_4	H_5	H_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

SG 152

Space group $P3_121$, No. 152, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6
	dim	2	2	2	2
Δ_4	1	1	0	0	-2
Δ_5	1	1	0	-1	0
Δ_6	1	0	2	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6	
	2	2	2	2	Bipartite graph
v^1	2	1	2	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
	dim	2	2	2	1	1
T_3	1	1	1	-1	0	-1
T_4	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
	dim	2	2	2
Σ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_4
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_4	K_5	K_6
dim		2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_4	A_5A_6	H_4	H_5	H_6
	dim	2	2	2	1	1
S_3	1	1	1	-1	-1	0
S_4	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_4	A_5A_6	L_3L_4
	dim	2	2	2
R_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4	H_5	H_6
	dim	2	2	1	1
S'_3	1	1	-1	0	-1
S'_4	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
	dim	2	1	1	2	1	1
P_4	1	1	0	0	0	-1	-1
P_5	1	1	0	0	-1	0	0
P_6	1	0	1	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	1	0	1	1	0	1	$H_{4/3}^2$
v^2	1	0	1	1	1	0	$H_{4/3}^2$
v^3	1	1	0	1	0	1	$H_{4/3}^2$
v^4	1	1	0	1	1	0	$H_{4/3}^2$

SG 153

Space group $P3_212$, No. 153, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6
	dim	2	2	2	2
Δ_4	1	1	0	-1	0
Δ_5	1	1	0	0	-2
Δ_6	1	0	2	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6	
	2	2	2	2	Bipartite graph
v^1	2	1	2	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
dim	2	2	1	1	1
T_2	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
dim	2	2	2
Σ_3	1	1	1
Σ_4	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_4
dim		2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_4	K_5	K_6
dim		2	1	1	1
T'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_3M_4	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_4	A_5A_6	H_4	H_5	H_6
dim	2	2	1	1	1
S_2	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4	A_5A_6	L_3L_4
dim	2	2	2
R_3	1	1	-1
R_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4	H_5	H_6
	dim	2	1	1	1
S'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_3L_4	H_4	H_5	H_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

SG 154

Space group $P3_221$, No. 154, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6
	dim	2	2	2	2
Δ_4	1	1	0	-1	0
Δ_5	1	1	0	0	-2
Δ_6	1	0	2	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6	
	2	2	2	2	Bipartite graph
v^1	2	1	2	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
	dim	2	2	2	1	1
T_3	1	1	1	-1	0	-1
T_4	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
	dim	2	2	2
Σ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_3M_4	L_3L_4
	dim	2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_3M_4	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4	A_5A_6	H_4	H_5	H_6
	dim	2	2	2	1	1
S_3	1	1	1	-1	0	-1
S_4	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_4	A_5A_6	L_3L_4
	dim	2	2	2
R_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4	H_5	H_6
dim		2	2	1	1
S'_3	1	1	-1	-1	0
S'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
dim		2	1	1	2	1	1
P_4	1	1	0	0	-1	0	0
P_5	1	1	0	0	0	-1	-1
P_6	1	0	1	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	1	0	1	1	0	1	$H_{4/3}^2$
v^2	1	0	1	1	1	0	$H_{4/3}^2$
v^3	1	1	0	1	0	1	$H_{4/3}^2$
v^4	1	1	0	1	1	0	$H_{4/3}^2$

SG 155

Space group $R32$, No. 155, with SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	F_3F_4
	dim	2	2	2
Σ_3	1	1	1	-1
Σ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	F_3F_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y : $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep		Z_4	Z_5Z_6	L_3L_4
	dim	2	2	2
Y_3	1	1	1	-1
Y_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_4	Z_5Z_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

SG 156

Space group $P3m1$, No. 156, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6
dim		2	2	2	2
Δ_4	2	1	0	-1	0
Δ_5	1	0	1	0	-1
Δ_6	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
dim		2	2	1	1	1
T_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
dim	2	2	2
Σ_3	1	1	1
Σ_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_3M_4	L_3L_4
dim	2	2
U_3	1	1
U_4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_3M_4	K_4	K_5	K_6
dim	2	1	1	1
T'_2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_3M_4	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_4	A_5A_6	H_4	H_5	H_6
dim		2	2	1	1	1
S_2	1	2	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix



corep		A_4	A_5A_6	L_3L_4
	dim	2	2	2
R_3	1	1	1	-1
R_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4	H_5	H_6
	dim	2	1	1	1
S'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_3L_4	H_4	H_5	H_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

SG 157

Space group $P31m$, No. 157, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix



corep		Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6
	dim	2	2	2	2
Δ_4	2	1	0	-1	0
Δ_5	1	0	1	0	-1
Δ_6	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	A_4	A_5A_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
	dim	2	2	2	1	1
T_3	1	1	1	-1	-1	0
T_4	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
	dim	2	2	2
Σ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_4
dim		2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_4	K_5	K_6
dim		2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_4	A_5A_6	H_4	H_5	H_6
	dim	2	2	2	1	1
S_3	1	1	1	-1	-1	0
S_4	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_4	A_5A_6	H_4	H_5	H_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_4	A_5A_6	L_3L_4
	dim	2	2	2
R_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4	A_5A_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4	H_5	H_6
	dim	2	2	1	1
S'_3	1	1	-1	-1	0
S'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
	dim	2	1	1	2	1	1
P_4	2	1	0	0	-1	0	0
P_5	1	0	1	0	0	-1	0
P_6	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

SG 158

Space group $P3c1$, No. 158, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_4	$\Gamma_5\Gamma_6$	A_4A_4	A_5A_5	A_6A_6
	dim	2	2	2	2	4
Δ_4	2	1	0	0	0	-2
Δ_5	1	0	1	-2	0	0
Δ_6	1	0	1	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	A_4A_4	A_5A_5	A_6A_6	
	2	2	2	2	4	Bipartite graph
v^1	0	2	1	1	0	$H_{4/4}^2$
v^2	2	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
dim	2	2	1	1	1
T_2	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	1	1	1	Bipartite graph
v^1	0	1	0	0	2	$H_{3/2}^1$
v^2	0	1	0	1	1	$H_{3/2}^1$
v^3	0	1	0	2	0	$H_{3/2}^1$
v^4	0	1	1	0	1	$H_{3/2}^1$
v^5	0	1	1	1	0	$H_{3/2}^1$
v^6	0	1	2	0	0	$H_{3/2}^1$
v^7	1	0	0	0	2	$H_{3/2}^1$
v^8	1	0	0	1	1	$H_{3/2}^1$
v^9	1	0	0	2	0	$H_{3/2}^1$
v^{10}	1	0	1	0	1	$H_{3/2}^1$
v^{11}	1	0	1	1	0	$H_{3/2}^1$
v^{12}	1	0	2	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
dim	2	2	2
Σ_3	1	1	1
Σ_4	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_3M_4	L_3L_3	L_4L_4
	dim	2	2	2
U_3	1	1	-2	0
U_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_3	L_4L_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_3M_4	K_4	K_5	K_6
	dim	2	1	1	1
T'_2	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_3M_4	K_4	K_5	K_6	
	2	1	1	1	Bipartite graph
v^1	1	0	0	2	$H_{3/2}^1$
v^2	1	0	1	1	$H_{3/2}^1$
v^3	1	0	2	0	$H_{3/2}^1$
v^4	1	1	0	1	$H_{3/2}^1$
v^5	1	1	1	0	$H_{3/2}^1$
v^6	1	2	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_4A_4	A_5A_5	A_6A_6	H_4H_4	H_5H_5	H_6H_6
dim	2	2	4	2	2	2
S_2S_2	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4A_4	A_5A_5	A_6A_6	H_4H_4	H_5H_5	H_6H_6	Bipartite graph
	2	2	4	2	2	2	
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4A_4	A_5A_5	A_6A_6	L_3L_3	L_4L_4
dim	2	2	4	2	2
R_3	1	2	0	2	-2
R_4	1	0	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_4A_4	A_5A_5	A_6A_6	L_3L_3	L_4L_4	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	0	1	0	0	1	$H_{2/2}^1$
v^3	1	0	0	1	0	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of *S'*: $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_3L_3	L_4L_4	H_4H_4	H_5H_5	H_6H_6	
dim	2	2	2	2	2	
$S'_2S'_2$	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_3L_3	L_4L_4	H_4H_4	H_5H_5	H_6H_6	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_4	K_5	K_6	H_4H_4	H_5H_5	H_6H_6	
dim	1	1	1	2	2	2	
P_4	1	1	0	0	-2	0	0
P_5	1	0	1	0	0	-2	0
P_6	1	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5	K_6	H_4H_4	H_5H_5	H_6H_6	
	1	1	1	2	2	2	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	2	0	0	1	0	$H_{3/2}^1$
v^3	2	0	0	1	0	0	$H_{3/2}^1$

SG 159

Space group *P31c*, No. 159, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	A_4A_4	A_5A_5	A_6A_6
dim	2	2	2	2	4
Δ_4	2	1 0	0	0	-2
Δ_5	1	0 1	-2	0	0
Δ_6	1	0 1	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	A_4A_4	A_5A_5	A_6A_6	
	2	2	2	2	4	Bipartite graph
v^1	0	2	1	1	0	$H_{4/4}^2$
v^2	2	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6
dim	2	2	2	1	1
T_3	1	1 1	-1	-1	0
T_4	1	1 1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_4	$\Gamma_5\Gamma_6$	K_4	K_5	K_6	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix



corep		Γ_4	$\Gamma_5\Gamma_6$	M_3M_4
	dim	2	2	2
Σ_2	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	M_3M_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_3	L_4L_4
	dim	2	2	2
U_3	1	1	-2	0
U_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_3	L_4L_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4A_4	A_5A_5	A_6A_6	H_4	H_5	H_6
	dim	2	2	4	1	1	2
S_3	1	2	0	2	-1	0	-1
S_4	1	0	2	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_4A_4	A_5A_5	A_6A_6	H_4	H_5	H_6	
	2	2	4	1	1	2	Bipartite graph
v^1	0	0	1	0	0	2	$H_{3/4}^1$
v^2	0	0	1	1	1	1	$H_{4/4}^1$
v^3	0	0	1	2	2	0	$H_{5/4}^1$
v^4	0	1	0	0	2	0	$H_{3/2}^1$
v^5	1	0	0	2	0	0	$H_{3/2}^1$
v^6	1	1	0	0	0	2	$H_{4/4}^2$
v^7	1	1	0	1	1	1	$H_{5/4}^3$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_4A_4	A_5A_5	A_6A_6	L_3L_3	L_4L_4
	dim	2	2	4	2	2
R_2R_2	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_4A_4	A_5A_5	A_6A_6	L_3L_3	L_4L_4	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_3	L_4L_4	H_4	H_5	H_6
dim		2	2	1	1	2
S'_3	1	2	0	-1	0	-1
S'_4	1	0	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_3L_3	L_4L_4	H_4	H_5	H_6	
	2	2	1	1	2	Bipartite graph
v^1	0	1	0	2	0	$H_{3/2}^1$
v^2	1	0	2	0	0	$H_{3/2}^1$
v^3	1	1	0	0	2	$H_{4/4}^2$
v^4	1	1	1	1	1	$H_{5/4}^3$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
dim		2	1	1	1	1	2
P_4	2	1	0	0	0	0	-1
P_5	1	0	1	0	-1	0	0
P_6	1	0	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	1	1	2	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	1	0	1	0	0	$H_{2/1}^1$
v^3	1	0	0	0	0	1	$H_{2/1}^1$

SG 160

Space group $R3m$, No. 160, with SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	F_3F_4
dim	2	2	2
Σ_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	F_3F_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y : $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$
A matrix

corep	Z_4	Z_5Z_6	L_3L_4
dim	2	2	2
Y_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_4	Z_5Z_6	L_3L_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

SG 161

Space group $R3c$, No. 161, with SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep	Γ_4	$\Gamma_5\Gamma_6$	F_3F_4
dim	2	2	2
Σ_2	1	2	2
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_4	$\Gamma_5\Gamma_6$	F_3F_4	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y: $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$
A matrix

corep	Z_4Z_4	Z_5Z_5	Z_6Z_6	L_3L_3	L_4L_4
dim	2	2	4	2	2
Y_2Y_2	2	1	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Z_4Z_4	Z_5Z_5	Z_6Z_6	L_3L_3	L_4L_4	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

SG 162

Space group $P\bar{3}1m$, No. 162, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	
dim	2	2	2	2	2	2	2	2	
Δ_4	2	1	1	0	0	-1	-1	0	0
$\Delta_5\Delta_6$	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6
dim	2	2	2	2	2	2
T_3T_4	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$
dim	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$M_3^+M_4^+$	$M_3^-M_4^-$	$L_3^+L_4^+$	$L_3^-L_4^-$
dim	2	2	2	2
U_3U_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_3^+M_4^+$	$M_3^-M_4^-$	$L_3^+L_4^+$	$L_3^-L_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	$M_3^+M_4^+$	$M_3^-M_4^-$	K_4	K_5K_6
dim	2	2	2	2
$T_3'T_4'$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	K_4	$K_5 K_6$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	H_4	$H_5 H_6$
dim	2	2	2	2	2	2
$S_3 S_4$	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	H_4	$H_5 H_6$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$
dim	2	2	2	2	2	2
$R_3 R_4$	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	$L_3^+ L_4^+$	$L_3^- L_4^-$	H_4	$H_5 H_6$
dim	2	2	2	2
$S'_3 S'_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$L_3^+ L_4^+$	$L_3^- L_4^-$	H_4	$H_5 H_6$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_4	$K_5 K_6$	H_4	$H_5 H_6$
dim	2	2	2	2
P_4	2	1	0	-1 0
$P_5 P_6$	2	0	1	0 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4	K_5K_6	H_4	H_5H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 163

Space group $P\bar{3}1c$, No. 163, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4A_4	A_5A_6
	dim	2	2	2	2	4	4
Δ_4	2	1	1	0	0	0	-2
$\Delta_5\Delta_6$	2	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4A_4	A_5A_6	
	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	2	1	0	$H_{3/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	2	0	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$
v^6	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6
	dim	2	2	2	2	2	2
T_3T_4	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$
dim	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_3^+M_4^+$	$M_3^-M_4^-$	L_2L_2
dim	2	2	4
U_3U_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	$L_2 L_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	$M_3^+ M_4^+$	$M_3^- M_4^-$	K_4	$K_5 K_6$
dim	2	2	2	2
$T_3' T_4'$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	K_4	$K_5 K_6$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	$A_4 A_4$	$A_5 A_6$	$H_4 H_5$	H_6
dim	4	4	2	2
$S_3 S_4$	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$A_4 A_4$	$A_5 A_6$	$H_4 H_5$	H_6	
	4	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	2	$H_{3/2}^1$
v^5	1	0	1	1	$H_{3/2}^1$
v^6	1	0	2	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4A_4	A_5A_6	L_2L_2
dim	4	4	4
R_3R_3	2	1	1
R_4R_4	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_4	A_5A_6	L_2L_2	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_2L_2	H_4H_5	H_6
dim	4	2	2
$S'_3S'_4$	2	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_2L_2	H_4H_5	H_6	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_4	K_5K_6	H_4H_5	H_6
dim	2	2	2	2
P_4	2	1	0	0
P_5P_6	2	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4	K_5K_6	H_4H_5	H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	1	0	$H_{2/1}^1$
v^2	1	0	0	1	$H_{2/1}^1$

SG 164

Space group $P\bar{3}m1$, No. 164, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$
dim		2	2	2	2	2	2	2	2
Δ_4	2	1	1	0	0	-1	-1	0	0
$\Delta_5\Delta_6$	2	0	0	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4^+	A_4^-	$A_5^+A_6^+$	$A_5^-A_6^-$	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6
dim		2	2	2	2	2	2
T_3T_4	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$
dim	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_3^+M_4^+$	$M_3^-M_4^-$	$L_3^+L_4^+$	$L_3^-L_4^-$
dim	2	2	2	2
U_3U_4	2	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	$M_3^+ M_4^+$	$M_3^- M_4^-$	K_4	$K_5 K_6$
dim	2	2	2	2
$T_3' T_4'$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	K_4	$K_5 K_6$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	H_4	$H_5 H_6$
dim	2	2	2	2	2	2
$S_3 S_4$	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	H_4	$H_5 H_6$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$

v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$
dim	2	2	2	2	2	2
$R_3 R_4$	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4^+	A_4^-	$A_5^+ A_6^+$	$A_5^- A_6^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	$L_3^+ L_4^+$	$L_3^- L_4^-$	H_4	$H_5 H_6$
dim	2	2	2	2
$S'_3 S'_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$L_3^+ L_4^+$	$L_3^- L_4^-$	H_4	$H_5 H_6$	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5K_6	H_4	H_5H_6
	dim	2	2	2	2
P_4P_5	2	1	0	-1	0
P_6P_6	2	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4	K_5K_6	H_4	H_5H_6	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$

SG 165

Space group $P\bar{3}c1$, No. 165, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4A_4	A_5A_6
	dim	2	2	2	2	4	4
Δ_4	2	1	1	0	0	0	-2
$\Delta_5\Delta_6$	2	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	A_4A_4	A_5A_6	Bipartite graph
	2	2	2	2	4	4	
v^1	0	0	0	2	1	0	$H_{3/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	0	2	0	1	0	$H_{3/2}^1$
v^4	0	2	0	0	0	1	$H_{3/2}^1$
v^5	1	1	0	0	0	1	$H_{3/2}^1$
v^6	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6
dim	2	2	2	2	2	2
T_3T_4	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	K_4	K_5K_6	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$
dim	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$M_3^+ M_4^+$	$M_3^- M_4^-$	$L_2 L_2$
dim	2	2	4
$U_3 U_4$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	$L_2 L_2$	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	$M_3^+ M_4^+$	$M_3^- M_4^-$	K_4	$K_5 K_6$
dim	2	2	2	2
$T'_3 T'_4$	2	1	1	-1
			-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	K_4	$K_5 K_6$	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	$A_4 A_4$	$A_5 A_6$	$H_4 H_4$	$H_5 H_5$	$H_6 H_6$
dim	4	4	4	2	2
$S_3 S_3$	2	1	1	-1	-1
$S_4 S_4$	2	1	1	-1	0
				0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_4A_4	A_5A_6	H_4H_4	H_5H_5	H_6H_6	
	4	4	4	2	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4A_4	A_5A_6	L_2L_2
dim	4	4	4
R_3R_4	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_4	A_5A_6	L_2L_2	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_2L_2	H_4H_4	H_5H_5	H_6H_6
dim	4	4	2	2
$S'_3S'_3$	2	1	-1	0
$S'_4S'_4$	2	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_2L_2	H_4H_4	H_5H_5	H_6H_6	
	4	4	2	2	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_4	K_5K_6	H_4H_4	H_5H_5	H_6H_6
dim	2	2	4	2	2
P_4P_5	2	1 0	-2	0	0
P_6P_6	2	0 1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5K_6	H_4H_4	H_5H_5	H_6H_6	
	2	2	4	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	2	0	1	0	0	$H_{3/2}^1$

SG 166

Space group $R\bar{3}m$, No. 166, with SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$F_3^+F_4^+$	$F_3^-F_4^-$
dim	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1 1	1 1	1 1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+\Gamma_6^+$	$\Gamma_5^-\Gamma_6^-$	$F_3^+F_4^+$	$F_3^-F_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y: $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$
A matrix

corep	Z_4^+	Z_4^-	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$
dim	2	2	2	2	2	2
$Y_3 Y_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Z_4^+	Z_4^-	$Z_5^+ Z_6^+$	$Z_5^- Z_6^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

SG 167

Space group $R\bar{3}c$, No. 167, with SOC

Σ

High symmetry line $\Sigma(0, u, -u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), aF(0, \frac{1}{2}, -\frac{1}{2})$
A matrix

corep	Γ_4^+	Γ_4^-	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$F_3^+ F_4^+$	$F_3^- F_4^-$
dim	2	2	2	2	2	2
$\Sigma_3 \Sigma_4$	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_4^+	Γ_4^-	$\Gamma_5^+ \Gamma_6^+$	$\Gamma_5^- \Gamma_6^-$	$F_3^+ F_4^+$	$F_3^- F_4^-$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$

v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Y

High symmetry line $Y(u, \frac{1}{2}, -u)$, high symmetry points at both ends of Y: $Z(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}), L(0, \frac{1}{2}, 0)$

A matrix

corep		Z_4Z_4	Z_5Z_6	L_2L_2
	dim	4	4	4
Y_3Y_3	2	1	1	-1
Y_4Y_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Z_4Z_4	Z_5Z_6	L_2L_2	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

SG 168

Space group $P6$, No. 168, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$
	dim	2	2	2	2	2	2
Δ_7	1	1	0	0	-1	0	0
Δ_8	1	1	0	0	-1	0	0
Δ_9	1	0	1	0	0	-1	0
Δ_{10}	1	0	1	0	0	-1	0
Δ_{11}	1	0	0	1	0	0	-1
Δ_{12}	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	1	0	0	1	0	0	$H_{2/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6
	dim	2	2	2	2	1
T_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6	Bipartite graph
	2	2	2	2	1	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4
	dim	2	2	2	2
Σ_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix



corep		M_3M_4	L_3L_4
dim		2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_3M_4	K_4K_5	K_6
dim		2	2	1
T'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4K_5	K_6	
	2	2	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_4H_5	H_6
dim		2	2	2	2	1
S_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_4H_5	H_6	
	2	2	2	2	1	Bipartite graph

v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4
dim		2	2	2	2
R_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4H_5	H_6
dim		2	2	1
S'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4H_5	H_6	
	2	2	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4K_5	K_6	H_4H_5	H_6
dim		2	1	2	1
P_4	1	1	0	-1	0
P_5	1	1	0	-1	0
P_6	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4K_5	K_6	H_4H_5	H_6	
	2	1	2	1	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 169

Space group $P6_1$, No. 169, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$
dim		2	2	2	2	2	2	2
Δ_7	1	1	0	0	0	0	-1	0
Δ_8	1	1	0	0	0	0	0	-2
Δ_9	1	0	1	0	0	0	-1	0
Δ_{10}	1	0	1	0	0	-1	0	0
Δ_{11}	1	0	0	1	-2	0	0	0
Δ_{12}	1	0	0	1	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	
	2	2	2	2	2	2	2	Bipartite graph
v^1	2	2	2	1	2	2	1	$H_{12/12}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6
	dim	2	2	2	2	1
T_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6	Bipartite graph
	2	2	2	2	1	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4
	dim	2	2	2	2
Σ_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix



v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	L_3L_3	L_4L_4
dim	2	2	2	2	2	2
R_2R_2	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	L_3L_3	L_4L_4	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_3L_3	L_4L_4	H_4H_5	H_6H_6
dim	2	2	2	2
$S'_2S'_2$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_3L_3	L_4L_4	H_4H_5	H_6H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4K_5	K_6	H_4H_5	H_6H_6
	dim	2	1	2	2
P_4	1	1	0	0	-2
P_5	1	1	0	-1	0
P_6	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	K_4K_5	K_6	H_4H_5	H_6H_6	
	2	1	2	2	Bipartite graph
v^1	2	2	2	1	$H_{7/6}^3$

SG 170

Space group $P6_5$, No. 170, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$
	dim	2	2	2	2	2	2	2
Δ_7	1	1	0	0	-2	0	0	0
Δ_8	1	1	0	0	0	-1	0	0
Δ_9	1	0	1	0	0	0	-1	0
Δ_{10}	1	0	1	0	0	-1	0	0
Δ_{11}	1	0	0	1	0	0	-1	0
Δ_{12}	1	0	0	1	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	
	2	2	2	2	2	2	2	Bipartite graph
v^1	2	2	2	1	2	2	1	$H_{12/12}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6
	dim	2	2	2	2	1
T_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6	
	2	2	2	2	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4
	dim	2	2	2	2
Σ_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_3M_4	L_3L_3	L_4L_4
	dim	2	2	2
U_3	1	1	-2	0
U_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_3	L_4L_4	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_3M_4	K_4K_5	K_6
	dim	2	2	1
T'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4K_5	K_6	
	2	2	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	H_4H_5	H_6H_6
	dim	2	2	2	2	2	2
S_2S_2	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	H_4H_5	H_6H_6	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	L_3L_3	L_4L_4
dim	2	2	2	2	2	2
R_2R_2	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	L_3L_3	L_4L_4	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_3L_3	L_4L_4	H_4H_5	H_6H_6
dim	2	2	2	2
$S'_2S'_2$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_3L_3	L_4L_4	H_4H_5	H_6H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4K_5	K_6	H_4H_5	H_6H_6
	dim	2	1	2	2
P_4	1	1	0	-1	0
P_5	1	1	0	0	-2
P_6	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	K_4K_5	K_6	H_4H_5	H_6H_6	
	2	1	2	2	Bipartite graph
v^1	2	2	2	1	$H_{7/6}^3$

SG 171

Space group $P6_2$, No. 171, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$
	dim	2	2	2	2	2	2
Δ_7	1	1	0	0	0	-1	0
Δ_8	1	1	0	0	-1	0	0
Δ_9	1	0	1	0	-1	0	0
Δ_{10}	1	0	1	0	0	0	-1
Δ_{11}	1	0	0	1	0	0	-1
Δ_{12}	1	0	0	1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	
	2	2	2	2	2	2	Bipartite graph
v^1	1	1	1	1	1	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6
	dim	2	2	2	2	1
T_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6	
	2	2	2	2	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4
	dim	2	2	2	2
Σ_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_3M_4	L_3L_4
dim		2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_3M_4	K_4K_5	K_6
dim		2	2	1
T'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4K_5	K_6	
	2	2	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_4H_5	H_6
dim		2	2	2	2	1
S_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_4H_5	H_6	Bipartite graph
	2	2	2	2	1	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4
dim		2	2	2	2
R_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4H_5	H_6
dim		2	2	1
S'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4H_5	H_6	Bipartite graph
	2	2	1	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4K_5	K_6	H_4H_5	H_6
dim		2	1	2	1
P_4	1	1	0	-1	0
P_5	1	1	0	0	-1
P_6	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		K_4K_5	K_6	H_4H_5	H_6	
		2	1	2	1	Bipartite graph
v^1		1	1	1	1	$H_{4/3}^2$

SG 172

Space group $P6_4$, No. 172, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$
dim		2	2	2	2	2	2
Δ_7	1	1	0	0	0	0	-1
Δ_8	1	1	0	0	0	-1	0
Δ_9	1	0	1	0	-1	0	0
Δ_{10}	1	0	1	0	0	0	-1
Δ_{11}	1	0	0	1	0	-1	0
Δ_{12}	1	0	0	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 1.

		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	
		2	2	2	2	2	2	Bipartite graph
v^1		1	1	1	1	1	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6
dim		2	2	2	2	1
T_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6	Bipartite graph
	2	2	2	2	1	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4
dim		2	2	2	2
Σ_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_4
dim		2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_4K_5	K_6
dim		2	2	1
T'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4K_5	K_6	
	2	2	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_4H_5	H_6
dim		2	2	2	2	1
S_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_4H_5	H_6	
	2	2	2	2	1	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4
dim		2	2	2	2
R_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_3L_4	H_4H_5	H_6
dim		2	2	1
S'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4H_5	H_6	
	2	2	1	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4K_5	K_6	H_4H_5	H_6
dim		2	1	2	1
P_4	1	1	0	0	-1
P_5	1	1	0	-1	0
P_6	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	K_4K_5	K_6	H_4H_5	H_6	
	2	1	2	1	Bipartite graph
v^1	1	1	1	1	$H_{4/3}^2$

SG 173

Space group $P6_3$, No. 173, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$
dim		2	2	2	2	2	2	2
Δ_7	1	1	0	0	0	-1	0	0
Δ_8	1	1	0	0	0	0	-1	0
Δ_9	1	0	1	0	0	0	0	-2
Δ_{10}	1	0	1	0	-2	0	0	0
Δ_{11}	1	0	0	1	0	-1	0	0
Δ_{12}	1	0	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	
	2	2	2	2	2	2	2	Bipartite graph
v^1	0	2	0	1	0	0	1	$H_{4/4}^2$
v^2	1	0	1	0	1	1	0	$H_{4/4}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6
dim		2	2	2	2	1
T_2	1	2	2	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_4K_5	K_6	Bipartite graph
	2	2	2	2	1	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	0	$H_{2/2}^1$
v^3	0	1	0	0	2	$H_{3/2}^1$
v^4	0	1	0	1	0	$H_{2/2}^1$
v^5	1	0	0	0	2	$H_{3/2}^1$
v^6	1	0	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4
dim		2	2	2	2
Σ_2	1	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_3M_4	L_3L_3	L_4L_4
dim		2	2	2
U_3	1	1	-2	0
U_4	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_3	L_4L_4	Bipartite graph
	2	2	2	
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_3M_4	K_4K_5	K_6
	dim	2	2	1
T'_2	1	2	-2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_3M_4	K_4K_5	K_6	Bipartite graph
	2	2	1	
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	H_4H_5	H_6H_6
	dim	2	2	2	2	2	2
S_2S_2	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	H_4H_5	H_6H_6	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix



corep	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	L_3L_3	L_4L_4
dim	2	2	2	2	2	2
R_2R_2	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_7A_7	A_8A_{12}	A_9A_{11}	$A_{10}A_{10}$	L_3L_3	L_4L_4	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_3L_3	L_4L_4	H_4H_5	H_6H_6
dim	2	2	2	2
$S'_2S'_2$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_3L_3	L_4L_4	H_4H_5	H_6H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix



corep		K_4K_5	K_6	H_4H_5	H_6H_6
	dim	2	1	2	2
P_4	1	1	0	-1	0
P_5	1	1	0	-1	0
P_6	1	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4K_5	K_6	H_4H_5	H_6H_6	
	2	1	2	2	Bipartite graph
v^1	0	2	0	1	$H_{3/2}^1$
v^2	1	0	1	0	$H_{2/2}^1$

SG 174

Space group $P\bar{6}$, No. 174, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$
	dim	2	2	2	2	2	2
$\Delta_4\Delta_5$	2	1	0	1	-1	0	-1
Δ_6	1	0	2	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	$H_{2/2}^1$
v^4	1	0	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}
dim		2	2	2	1	1	1	1	1	1
T_3	1	1	1	1	-1	0	-1	0	-1	0
T_4	1	1	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 27.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	Bipartite graph
	2	2	2	1	1	1	1	1	1	
v^1	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{14}	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{16}	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{17}	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{18}	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{20}	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{21}	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{23}	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{25}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0)$, $M(0, \frac{1}{2}, 0)$

A matrix



corep		$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$\Gamma_7\Gamma_8$	$\Gamma_9\Gamma_{10}$	$\Gamma_{11}\Gamma_{12}$	M_3M_4	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_3M_4	L_3L_4
	dim	2	2
U_2	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_3M_4	L_3L_4	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_3M_4	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}
	dim	2	1	1	1	1	1	1
T'_3	1	1	-1	0	-1	0	-1	0
T'_4	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	M_3M_4	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	
	2	1	1	1	1	1	1	Bipartite graph
v^1	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	1	1	1	0	0	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}
dim		2	2	2	1	1	1	1	1	1
S_3	1	1	1	1	-1	0	-1	0	-1	0
S_4	1	1	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 27.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}	
	2	2	2	1	1	1	1	1	1	Bipartite graph
v^1	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{14}	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{16}	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$

v^{17}	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{18}	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{20}	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{21}	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{23}	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{25}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4
	dim	2	2	2	2
R_3	1	1	1	1	-1
R_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7A_8	A_9A_{10}	$A_{11}A_{12}$	L_3L_4	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_3L_4	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}
	dim	2	1	1	1	1	1	1
S'_3	1	1	-1	0	-1	0	-1	0
S'_4	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	L_3L_4	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}	
	2	1	1	1	1	1	1	Bipartite graph
v^1	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	1	1	1	0	0	0	0	$H_{3/2}^1$

SG 175

Space group $P6/m$, No. 175, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	$A_7^+A_8^+$	$A_7^-A_8^-$	$A_9^+A_{10}^+$	$A_9^-A_{10}^-$	$A_{11}^+A_{12}^+$	$A_{11}^-A_{12}^-$
dim	2	2	2	2	2	2	2	2	2	2	2	2
$\Delta_7\Delta_8$	2	1	1	0	0	0	0	-1	-1	0	0	0
$\Delta_9\Delta_{10}$	2	0	0	1	1	0	0	0	0	-1	-1	0
$\Delta_{11}\Delta_{12}$	2	0	0	0	0	1	1	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	$A_7^+A_8^+$	$A_7^-A_8^-$	$A_9^+A_{10}^+$	$A_9^-A_{10}^-$	$A_{11}^+A_{12}^+$	$A_{11}^-A_{12}^-$	Bipartite graph
	2	2	2	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	K_7K_8	K_9K_{10}	$K_{11}K_{12}$
dim	2	2	2	2	2	2	2	2	2
T_3T_4	2	1	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	K_7K_8	K_9K_{10}	$K_{11}K_{12}$	Bipartite graph
	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	$M_3^+M_4^+$	$M_3^-M_4^-$
dim	2	2	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	$M_3^+M_4^+$	$M_3^-M_4^-$	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_3^+M_4^+$	$M_3^-M_4^-$	$L_3^+L_4^+$	$L_3^-L_4^-$
dim	2	2	2	2
U_3U_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$M_3^+M_4^+$	$M_3^-M_4^-$	$L_3^+L_4^+$	$L_3^-L_4^-$	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	$M_3^+M_4^+$	$M_3^-M_4^-$	K_7K_8	K_9K_{10}	$K_{11}K_{12}$
dim	2	2	2	2	2
$T'_3T'_4$	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	$K_7 K_8$	$K_9 K_{10}$	$K_{11} K_{12}$	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	$A_7^+ A_8^+$	$A_7^- A_8^-$	$A_9^+ A_{10}^+$	$A_9^- A_{10}^-$	$A_{11}^+ A_{12}^+$	$A_{11}^- A_{12}^-$	$H_7 H_8$	$H_9 H_{10}$	$H_{11} H_{12}$
dim	2	2	2	2	2	2	2	2	2
$S_3 S_4$	2	1	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	$A_7^+ A_8^+$	$A_7^- A_8^-$	$A_9^+ A_{10}^+$	$A_9^- A_{10}^-$	$A_{11}^+ A_{12}^+$	$A_{11}^- A_{12}^-$	$H_7 H_8$	$H_9 H_{10}$	$H_{11} H_{12}$	
	2	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	$A_7^+ A_8^+$	$A_7^- A_8^-$	$A_9^+ A_{10}^+$	$A_9^- A_{10}^-$	$A_{11}^+ A_{12}^+$	$A_{11}^- A_{12}^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$
dim	2	2	2	2	2	2	2	2
$R_3 R_4$	2	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$A_7^+ A_8^+$	$A_7^- A_8^-$	$A_9^+ A_{10}^+$	$A_9^- A_{10}^-$	$A_{11}^+ A_{12}^+$	$A_{11}^- A_{12}^-$	$L_3^+ L_4^+$	$L_3^- L_4^-$	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	$L_3^+ L_4^+$	$L_3^- L_4^-$	$H_7 H_8$	$H_9 H_{10}$	$H_{11} H_{12}$
dim	2	2	2	2	2
$S'_3 S'_4$	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$L_3^+ L_4^+$	$L_3^- L_4^-$	$H_7 H_8$	$H_9 H_{10}$	$H_{11} H_{12}$	Bipartite graph
	2	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_7K_8	K_9K_{10}	$K_{11}K_{12}$	H_7H_8	H_9H_{10}	$H_{11}H_{12}$	
dim	2	2	2	2	2	2	
P_4P_5	2	1	0	1	-1	0	-1
P_6P_6	2	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_7K_8	K_9K_{10}	$K_{11}K_{12}$	H_7H_8	H_9H_{10}	$H_{11}H_{12}$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

SG 176

Space group $P6_3/m$, No. 176, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	A_4A_6	A_5A_5	
dim	2	2	2	2	2	2	4	4	
$\Delta_7\Delta_8$	2	1	1	0	0	0	0	-1	0
$\Delta_9\Delta_{10}$	2	0	0	1	1	0	0	0	-2
$\Delta_{11}\Delta_{12}$	2	0	0	0	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	A_4A_6	A_5A_5	Bipartite graph
	2	2	2	2	2	2	4	4	
v^1	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^3	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^4	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	1	0	$H_{3/2}^1$
v^6	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	0	0	1	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	K_7K_8	K_9K_{10}	$K_{11}K_{12}$
dim	2	2	2	2	2	2	2	2	2
T_3T_4	2	1	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	K_7K_8	K_9K_{10}	$K_{11}K_{12}$	Bipartite graph
	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	$\Gamma_7^+\Gamma_8^+$	$\Gamma_7^-\Gamma_8^-$	$\Gamma_9^+\Gamma_{10}^+$	$\Gamma_9^-\Gamma_{10}^-$	$\Gamma_{11}^+\Gamma_{12}^+$	$\Gamma_{11}^-\Gamma_{12}^-$	$M_3^+M_4^+$	$M_3^-M_4^-$
dim	2	2	2	2	2	2	2	2
$\Sigma_3\Sigma_4$	2	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	$\Gamma_7^+ \Gamma_8^+$	$\Gamma_7^- \Gamma_8^-$	$\Gamma_9^+ \Gamma_{10}^+$	$\Gamma_9^- \Gamma_{10}^-$	$\Gamma_{11}^+ \Gamma_{12}^+$	$\Gamma_{11}^- \Gamma_{12}^-$	$M_3^+ M_4^+$	$M_3^- M_4^-$	
	2	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$M_3^+ M_4^+$	$M_3^- M_4^-$	$L_2 L_2$
dim	2	2	4
$U_3 U_4$	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	$L_2 L_2$	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	$M_3^+ M_4^+$	$M_3^- M_4^-$	$K_7 K_8$	$K_9 K_{10}$	$K_{11} K_{12}$
dim	2	2	2	2	2
$T'_3 T'_4$	2	1	1	-1	-1
					-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	$M_3^+ M_4^+$	$M_3^- M_4^-$	$K_7 K_8$	$K_9 K_{10}$	$K_{11} K_{12}$	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	$A_4 A_6$	$A_5 A_5$	$H_7 H_{11}$	$H_8 H_{12}$	$H_9 H_9$	$H_{10} H_{10}$	
dim	4	4	2	2	2	2	
$S_3 S_3$	2	1	1	-1	0	-1	0
$S_4 S_4$	2	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$A_4 A_6$	$A_5 A_5$	$H_7 H_{11}$	$H_8 H_{12}$	$H_9 H_9$	$H_{10} H_{10}$	
	4	4	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	1	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	$A_4 A_6$	$A_5 A_5$	$L_2 L_2$	
dim	4	4	4	
$R_3 R_3$	2	1	1	-1
$R_4 R_4$	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_6	A_5A_5	L_2L_2	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_2L_2	H_7H_{11}	H_8H_{12}	H_9H_9	$H_{10}H_{10}$
dim		4	2	2	2	2
$S'_3S'_3$	2	1	-1	0	-1	0
$S'_4S'_4$	2	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_2L_2	H_7H_{11}	H_8H_{12}	H_9H_9	$H_{10}H_{10}$	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_7K_8	K_9K_{10}	$K_{11}K_{12}$	H_7H_{11}	H_8H_{12}	H_9H_9	$H_{10}H_{10}$
dim		2	2	2	2	2	2	2
P_4P_5	2	1	0	1	-1	-1	0	0
P_6P_6	2	0	1	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	K_7K_8	K_9K_{10}	$K_{11}K_{12}$	H_7H_{11}	H_8H_{12}	H_9H_9	$H_{10}H_{10}$	
	2	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	1	0	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	1	0	0	1	0	0	0	$H_{2/1}^1$

SG 177

Space group $P622$, No. 177, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_7	A_8	A_9
dim		2	2	2	2	2	2
Δ_7	1	0	0	1	0	0	-1
Δ_8	1	0	0	1	0	0	-1
Δ_9	1	1	0	0	-1	0	0
Δ_{10}	1	1	0	0	-1	0	0
Δ_{11}	1	0	1	0	0	-1	0
Δ_{12}	1	0	1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	A_7	A_8	A_9	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	1	0	0	1	0	0	$H_{2/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
dim		2	2	2	2	1	1
T_3	1	1	1	1	-1	0	-1
T_4	1	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
dim		2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	L_5
dim		2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_7	A_8	A_9	H_4	H_5	H_6
	dim	2	2	2	2	1	1
S_3	1	1	1	1	-1	0	-1
S_4	1	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7	A_8	A_9	H_4	H_5	H_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_7	A_8	A_9	L_5
	dim	2	2	2	2
R_3	1	1	1	1	-1
R_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7	A_8	A_9	L_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_5	H_4	H_5	H_6
	dim	2	2	1	1
S'_3	1	1	-1	-1	0
S'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_5	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
	dim	2	1	1	2	1	1
P_4	1	1	0	0	-1	0	0
P_5	1	1	0	0	-1	0	0
P_6	1	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/2}^1$

SG 178

Space group $P6_122$, No. 178, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_4A_5	A_6A_7	A_8	A_9
	dim	2	2	2	2	2	2	2
Δ_7	1	0	0	1	0	0	0	-1
Δ_8	1	0	0	1	0	-2	0	0
Δ_9	1	1	0	0	0	0	0	-1
Δ_{10}	1	1	0	0	0	0	-1	0
Δ_{11}	1	0	1	0	-2	0	0	0
Δ_{12}	1	0	1	0	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_7	Γ_8	Γ_9	A_4A_5	A_6A_7	A_8	A_9	
	2	2	2	2	2	2	2	Bipartite graph
v^1	2	2	2	1	1	2	2	$H_{12/12}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	0	-1
T_4	1	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_2L_3	L_4L_5
	dim	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_2L_3	L_4L_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_2L_3	L_4L_5	H_4	H_5H_6
dim	2	2	2	2
$S'_3S'_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_2L_3	L_4L_5	H_4	H_5H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_4	K_5	K_6	H_4	H_5H_6
dim	2	1	1	2	2
P_4	1	1	0	0	0 -2
P_5	1	1	0	0	-1 0
P_6	1	0	1	1	-1 0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5	K_6	H_4	H_5H_6	
	2	1	1	2	2	Bipartite graph
v^1	2	0	2	2	1	$H_{7/6}^3$
v^2	2	1	1	2	1	$H_{7/6}^3$
v^3	2	2	0	2	1	$H_{7/6}^3$

SG 179

Space group $P6_522$, No. 179, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_4A_5	A_6A_7	A_8	A_9
	dim	2	2	2	2	2	2	2
Δ_7	1	0	0	1	-2	0	0	0
Δ_8	1	0	0	1	0	0	-1	0
Δ_9	1	1	0	0	0	0	0	-1
Δ_{10}	1	1	0	0	0	0	-1	0
Δ_{11}	1	0	1	0	0	0	0	-1
Δ_{12}	1	0	1	0	0	-2	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_7	Γ_8	Γ_9	A_4A_5	A_6A_7	A_8	A_9	
	2	2	2	2	2	2	2	Bipartite graph
v^1	2	2	2	1	1	2	2	$H_{12/12}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	0	-1
T_4	1	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_2L_3	L_4L_5
	dim	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_2L_3	L_4L_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4A_5	A_6A_7	A_8	A_9	H_4	H_5H_6
	dim	2	2	2	2	2	2
S_3S_4	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4A_5	A_6A_7	A_8	A_9	H_4	H_5H_6	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4A_5	A_6A_7	A_8	A_9	L_2L_3	L_4L_5
dim	2	2	2	2	2	2
R_3R_4	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4A_5	A_6A_7	A_8	A_9	L_2L_3	L_4L_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_2L_3	L_4L_5	H_4	H_5H_6
dim	2	2	2	2
$S'_3S'_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_2L_3	L_4L_5	H_4	H_5H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5H_6
	dim	2	1	1	2	2
P_4	1	1	0	0	-1	0
P_5	1	1	0	0	0	-2
P_6	1	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5	K_6	H_4	H_5H_6	
	2	1	1	2	2	Bipartite graph
v^1	2	0	2	2	1	$H_{7/6}^3$
v^2	2	1	1	2	1	$H_{7/6}^3$
v^3	2	2	0	2	1	$H_{7/6}^3$

SG 180

Space group $P6_222$, No. 180, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_7	A_8	A_9
	dim	2	2	2	2	2	2
Δ_7	1	0	0	1	-1	0	0
Δ_8	1	0	0	1	0	0	-1
Δ_9	1	1	0	0	0	0	-1
Δ_{10}	1	1	0	0	0	-1	0
Δ_{11}	1	0	1	0	0	-1	0
Δ_{12}	1	0	1	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_7	Γ_8	Γ_9	A_7	A_8	A_9	
	2	2	2	2	2	2	Bipartite graph
v^1	1	1	1	1	1	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	0	-1
T_4	1	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	L_5
	dim	2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_7	A_8	A_9	H_4	H_5	H_6
	dim	2	2	2	2	1	1
S_3	1	1	1	1	-1	-1	0
S_4	1	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7	A_8	A_9	H_4	H_5	H_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7	A_8	A_9	L_5
	dim	2	2	2	2
R_3	1	1	1	1	-1
R_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7	A_8	A_9	L_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_5	H_4	H_5	H_6
	dim	2	2	1	1
S'_3	1	1	-1	-1	0
S'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_5	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
	dim	2	1	1	2	1	1
P_4	1	1	0	0	-1	0	0
P_5	1	1	0	0	0	-1	-1
P_6	1	0	1	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	1	0	1	1	0	1	$H_{4/3}^2$
v^2	1	0	1	1	1	0	$H_{4/3}^2$
v^3	1	1	0	1	0	1	$H_{4/3}^2$
v^4	1	1	0	1	1	0	$H_{4/3}^2$

SG 181

Space group $P6_422$, No. 181, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_7	Γ_8	Γ_9	A_7	A_8	A_9
	dim	2	2	2	2	2	2
Δ_7	1	0	0	1	0	-1	0
Δ_8	1	0	0	1	-1	0	0
Δ_9	1	1	0	0	0	0	-1
Δ_{10}	1	1	0	0	0	-1	0
Δ_{11}	1	0	1	0	-1	0	0
Δ_{12}	1	0	1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	Γ_7	Γ_8	Γ_9	A_7	A_8	A_9	
	2	2	2	2	2	2	Bipartite graph
v^1	1	1	1	1	1	1	$H_{6/6}^5$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	0	-1
T_4	1	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_5
	dim	2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7	A_8	A_9	H_4	H_5	H_6
	dim	2	2	2	2	1	1
S_3	1	1	1	1	-1	-1	0
S_4	1	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7	A_8	A_9	H_4	H_5	H_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7	A_8	A_9	L_5
	dim	2	2	2	2
R_3	1	1	1	1	-1
R_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7	A_8	A_9	L_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_5	H_4	H_5	H_6
	dim	2	2	1	1
S'_3	1	1	-1	-1	0
S'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_5	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
	dim	2	1	1	2	1	1
P_4	1	1	0	0	0	-1	-1
P_5	1	1	0	0	-1	0	0
P_6	1	0	1	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	1	0	1	1	0	1	$H_{4/3}^2$
v^2	1	0	1	1	1	0	$H_{4/3}^2$
v^3	1	1	0	1	0	1	$H_{4/3}^2$
v^4	1	1	0	1	1	0	$H_{4/3}^2$

SG 182

Space group $P6_322$, No. 182, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_4A_5	A_6A_7	A_8	A_9
	dim	2	2	2	2	2	2	2
Δ_7	1	0	0	1	0	0	-1	0
Δ_8	1	0	0	1	0	0	0	-1
Δ_9	1	1	0	0	0	-2	0	0
Δ_{10}	1	1	0	0	-2	0	0	0
Δ_{11}	1	0	1	0	0	0	-1	0
Δ_{12}	1	0	1	0	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_7	Γ_8	Γ_9	A_4A_5	A_6A_7	A_8	A_9	
	2	2	2	2	2	2	2	Bipartite graph
v^1	0	1	1	0	0	1	1	$H_{4/4}^2$
v^2	2	0	0	1	1	0	0	$H_{4/4}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	0	-1
T_4	1	1	1	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_2L_3	L_4L_5
	dim	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_2L_3	L_4L_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4A_5	A_6A_7	A_8	A_9	H_4	H_5H_6
	dim	2	2	2	2	2	2
S_3S_4	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4A_5	A_6A_7	A_8	A_9	H_4	H_5H_6	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4A_5	A_6A_7	A_8	A_9	L_2L_3	L_4L_5
dim	2	2	2	2	2	2
R_3R_4	2	1	1	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	A_4A_5	A_6A_7	A_8	A_9	L_2L_3	L_4L_5	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_2L_3	L_4L_5	H_4	H_5H_6
dim	2	2	2	2
$S'_3S'_4$	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_2L_3	L_4L_5	H_4	H_5H_6	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5H_6
	dim	2	1	1	2	2
P_4	1	1	0	0	-1	0
P_5	1	1	0	0	-1	0
P_6	1	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_4	K_5	K_6	H_4	H_5H_6	
	2	1	1	2	2	Bipartite graph
v^1	0	0	2	0	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	$H_{2/2}^1$

SG 183

Space group $P6mm$, No. 183, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_7	A_8	A_9
	dim	2	2	2	2	2	2
Δ_7	2	1	0	0	-1	0	0
Δ_8	2	0	1	0	0	-1	0
Δ_9	2	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	A_7	A_8	A_9	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
dim		2	2	2	2	1	1
T_3	1	1	1	1	-1	-1	0
T_4	1	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
dim		2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_5
dim		2	2
U_5	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7	A_8	A_9	H_4	H_5	H_6
	dim	2	2	2	2	1	1
S_3	1	1	1	1	-1	-1	0
S_4	1	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7	A_8	A_9	H_4	H_5	H_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7	A_8	A_9	L_5
	dim	2	2	2	2
R_3	1	1	1	1	-1
R_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7	A_8	A_9	L_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_5	H_4	H_5	H_6
	dim	2	2	1	1
S'_3	1	1	-1	-1	0
S'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_5	H_4	H_5	H_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_4	K_5	K_6	H_4	H_5	H_6
	dim	2	1	1	2	1	1
P_4	2	1	0	0	-1	0	0
P_5	1	0	1	0	0	-1	0
P_6	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5	K_6	H_4	H_5	H_6	
	2	1	1	2	1	1	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	0	$H_{2/1}^1$

SG 184

Space group $P6cc$, No. 184, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_7A_7	A_8A_8	A_9A_9
	dim	2	2	2	4	4	4
Δ_7	2	1	0	0	0	0	-2
Δ_8	2	0	1	0	-2	0	0
Δ_9	2	0	0	1	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	A_7A_7	A_8A_8	A_9A_9	
	2	2	2	4	4	4	Bipartite graph
v^1	0	0	2	0	1	0	$H_{3/2}^1$
v^2	0	2	0	1	0	0	$H_{3/2}^1$
v^3	2	0	0	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	-1	0
T_4	1	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	L_5L_5
	dim	2	4
U_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5L_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_4	K_5	K_6
dim		2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7A_7	A_8A_8	A_9A_9	H_4H_5	H_6H_6
dim		4	4	4	2	4
S_3S_4	2	2	2	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7A_7	A_8A_8	A_9A_9	H_4H_5	H_6H_6	
	4	4	4	2	4	Bipartite graph
v^1	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	1	2	0	$H_{3/2}^1$
v^3	0	1	0	0	1	$H_{2/2}^1$
v^4	0	1	0	2	0	$H_{3/2}^1$
v^5	1	0	0	0	1	$H_{2/2}^1$
v^6	1	0	0	2	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix



corep	A_7A_7	A_8A_8	A_9A_9	L_5L_5
dim	4	4	4	4
R_3R_4	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7A_7	A_8A_8	A_9A_9	L_5L_5	
	4	4	4	4	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_5L_5	H_4H_5	H_6H_6
dim	4	2	4
$S'_3S'_4$	2	2	-1 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_5L_5	H_4H_5	H_6H_6	
	4	2	4	Bipartite graph
v^1	1	0	1	$H_{2/2}^1$
v^2	1	2	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	K_4	K_5	K_6	H_4H_5	H_6H_6
dim	2	1	1	2	4
P_4	2	1	0	0	-2
P_5	1	0	1	0	0
P_6	1	0	0	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4	K_5	K_6	H_4H_5	H_6H_6	
	2	1	1	2	4	Bipartite graph
v^1	0	1	1	1	0	$H_{3/2}^1$
v^2	2	0	0	0	1	$H_{3/2}^1$

SG 185

Space group $P6_3cm$, No. 185, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_7A_7	A_8A_9
	dim	2	2	2	4	4
Δ_7	2	1	0	0	-2	0
Δ_8	2	0	1	0	0	-1
Δ_9	2	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_7	Γ_8	Γ_9	A_7A_7	A_8A_9	
	2	2	2	4	4	Bipartite graph
v^1	0	1	1	0	1	$H_{3/2}^1$
v^2	2	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	-1	0
T_4	1	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph

v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	L_5L_5
	dim	2	4
U_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5L_5	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7A_7	A_8A_9	H_4H_4	H_5H_5	H_6H_6
	dim	4	4	4	2	2
S_3S_3	2	1	1	-1	-1	0
S_4S_4	2	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_7A_7	A_8A_9	H_4H_4	H_5H_5	H_6H_6	
	4	4	4	2	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R : $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7A_7	A_8A_9	L_5L_5
	dim	4	4	4
R_3R_4	2	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_7A_7	A_8A_9	L_5L_5	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_5L_5	H_4H_4	H_5H_5	H_6H_6
dim		4	4	2	2
$S'_3S'_3$	2	1	-1	-1	0
$S'_4S'_4$	2	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_5L_5	H_4H_4	H_5H_5	H_6H_6	
	4	4	2	2	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		K_4	K_5	K_6	H_4H_4	H_5H_5	H_6H_6
dim		2	1	1	4	2	2
P_4	2	1	0	0	-2	0	0
P_5	1	0	1	0	0	-2	0
P_6	1	0	0	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	K_4	K_5	K_6	H_4H_4	H_5H_5	H_6H_6	
	2	1	1	4	2	2	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	2	0	0	1	0	$H_{3/2}^1$
v^3	2	0	0	1	0	0	$H_{3/2}^1$

SG 186

Space group $P6_3mc$, No. 186, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_7A_7	A_8A_9
	dim	2	2	2	4	4
Δ_7	2	1	0	0	-2	0
Δ_8	2	0	1	0	0	-1
Δ_9	2	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_7	Γ_8	Γ_9	A_7A_7	A_8A_9	
	2	2	2	4	4	Bipartite graph
v^1	0	1	1	0	1	$H_{3/2}^1$
v^2	2	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	K_4	K_5	K_6
	dim	2	2	2	2	1	1
T_3	1	1	1	1	-1	-1	0
T_4	1	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_7	Γ_8	Γ_9	K_4	K_5	K_6	
	2	2	2	2	1	1	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/2}^1$
v^2	0	0	1	1	0	0	$H_{2/2}^1$
v^3	0	1	0	0	1	1	$H_{3/2}^1$
v^4	0	1	0	1	0	0	$H_{2/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	0	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	$L_5 L_5$
	dim	2	4
U_5	2	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	$L_5 L_5$	
	2	4	Bipartite graph
v^1	2	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_5	K_4	K_5	K_6
	dim	2	2	1	1
T'_3	1	1	-1	-1	0
T'_4	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	K_4	K_5	K_6	
	2	2	1	1	Bipartite graph
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_7A_7	A_8A_9	H_4H_5	H_6
dim	4	4	2	2
S_3S_4	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	A_7A_7	A_8A_9	H_4H_5	H_6	
	4	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	2	$H_{3/2}^1$
v^5	1	0	1	1	$H_{3/2}^1$
v^6	1	0	2	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_7A_7	A_8A_9	L_5L_5
dim	4	4	4
R_3R_3	2	1	1
R_4R_4	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_7A_7	A_8A_9	L_5L_5	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_5L_5	H_4H_5	H_6
dim	4	2	2
$S'_3S'_4$	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_5L_5	H_4H_5	H_6	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	1	2	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_4	K_5	K_6	H_4H_5	H_6
dim	2	1	1	2	2
P_4	2	1	0	0	-1
P_5	1	0	1	0	-1
P_6	1	0	0	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	K_4	K_5	K_6	H_4H_5	H_6	
	2	1	1	2	2	Bipartite graph
v^1	0	1	1	1	0	$H_{3/2}^1$
v^2	1	0	0	0	1	$H_{2/1}^1$

SG 187

Space group $P\bar{6}m2$, No. 187, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_7	Γ_8	Γ_9	A_7	A_8	A_9
	dim	2	2	2	2	2	2
Δ_4	2	0	1	1	0	-1	-1
Δ_5	1	1	0	0	-1	0	0
Δ_6	1	1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_7	Γ_8	Γ_9	A_7	A_8	A_9	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}
	dim	2	2	2	1	1	1	1	1	1
T_3	1	1	1	1	-1	0	-1	0	-1	0
T_4	1	1	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 27.

	Γ_7	Γ_8	Γ_9	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	Bipartite graph
	2	2	2	1	1	1	1	1	1	
v^1	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	1	0	0	$H_{3/2}^1$

v^{14}	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{16}	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{17}	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{18}	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{20}	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{21}	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{23}	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{25}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
dim	2	2	2	2	2
Σ_5	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_5
dim	2	2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}
	dim	2	1	1	1	1	1	1
T'_3	1	1	-1	0	-1	0	-1	0
T'_4	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	M_5	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	
	2	1	1	1	1	1	1	Bipartite graph
v^1	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	1	1	1	0	0	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_7	A_8	A_9	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}
	dim	2	2	2	1	1	1	1	1	1
S_3	1	1	1	1	-1	0	-1	0	-1	0
S_4	1	1	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 27.

	A_7	A_8	A_9	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}	Bipartite graph
	2	2	2	1	1	1	1	1	1	
v^1	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{14}	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{16}	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{17}	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{18}	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{20}	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{21}	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{23}	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{25}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_7	A_8	A_9	L_5
dim	2	2	2	2
R_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7	A_8	A_9	L_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_5	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}
	dim	2	1	1	1	1	1	1
S'_3	1	1	-1	0	-1	0	-1	0
S'_4	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	L_5	H_7	H_8	H_9	H_{10}	H_{11}	H_{12}	
	2	1	1	1	1	1	1	Bipartite graph
v^1	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	1	1	1	0	0	0	0	$H_{3/2}^1$

SG 188

Space group $P\bar{6}c2$, No. 188, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep	Γ_7	Γ_8	Γ_9	A_4A_6	A_5A_7	A_8A_9
	dim	2	2	2	2	4

Δ_4	2	0	1	1	0	0	-2
Δ_5	1	1	0	0	-2	0	0
Δ_6	1	1	0	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_7	Γ_8	Γ_9	A_4A_6	A_5A_7	A_8A_9	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	2	0	0	0	1	$H_{3/2}^1$
v^4	2	0	0	1	1	0	$H_{4/4}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}
	dim	2	2	2	1	1	1	1	1	1
T_3	1	1	1	1	-1	0	-1	0	-1	0
T_4	1	1	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 27.

	Γ_7	Γ_8	Γ_9	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	
	2	2	2	1	1	1	1	1	1	Bipartite graph
v^1	0	0	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	0	0	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	0	0	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	0	0	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	0	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	0	0	1	1	1	0	0	0	0	$H_{3/2}^1$
v^{10}	0	1	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{11}	0	1	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{12}	0	1	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{13}	0	1	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{14}	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{16}	0	1	0	1	0	0	0	0	1	$H_{3/2}^1$

v^{17}	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{18}	0	1	0	1	1	0	0	0	0	$H_{3/2}^1$
v^{19}	1	0	0	0	0	0	0	1	1	$H_{3/2}^1$
v^{20}	1	0	0	0	0	0	1	1	0	$H_{3/2}^1$
v^{21}	1	0	0	0	0	1	0	0	1	$H_{3/2}^1$
v^{22}	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^{23}	1	0	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{24}	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{25}	1	0	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{27}	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
dim	2	2	2	2	2
Σ_5	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/1}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	L_2L_3	L_4L_5
dim	2	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_2L_3	L_4L_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T' : $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		M_5	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}
	dim	2	1	1	1	1	1	1
T'_3	1	1	-1	0	-1	0	-1	0
T'_4	1	1	0	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	M_5	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	Bipartite graph
	2	1	1	1	1	1	1	
v^1	1	0	0	0	0	1	1	$H_{3/2}^1$
v^2	1	0	0	0	1	1	0	$H_{3/2}^1$
v^3	1	0	0	1	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	1	0	1	0	0	1	0	$H_{3/2}^1$
v^6	1	0	1	1	0	0	0	$H_{3/2}^1$
v^7	1	1	0	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	0	$H_{3/2}^1$
v^9	1	1	1	0	0	0	0	$H_{3/2}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S : $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		A_4A_6	A_5A_7	A_8A_9	H_7H_{12}	H_8H_{11}	H_9H_{10}
	dim	2	2	4	2	2	2
S_3S_4	2	1	1	2	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_4A_6	A_5A_7	A_8A_9	H_7H_{12}	H_8H_{11}	H_9H_{10}	Bipartite graph
	2	2	4	2	2	2	
v^1	0	0	1	0	0	2	$H_{3/2}^1$
v^2	0	0	1	0	1	1	$H_{3/2}^1$
v^3	0	0	1	0	2	0	$H_{3/2}^1$
v^4	0	0	1	1	0	1	$H_{3/2}^1$
v^5	0	0	1	1	1	0	$H_{3/2}^1$
v^6	0	0	1	2	0	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$

v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_4A_6	A_5A_7	A_8A_9	L_2L_3	L_4L_5
	dim	2	2	4	2	2
R_2	1	1	0	1	0	-1
R_3	1	0	1	1	-1	0
R_4	1	1	0	1	0	-1
R_5	1	0	1	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_4A_6	A_5A_7	A_8A_9	L_2L_3	L_4L_5	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	0	1	0	1	0	$H_{2/2}^1$
v^3	1	0	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_2L_3	L_4L_5	H_7H_{12}	H_8H_{11}	H_9H_{10}
	dim	2	2	2	2	2
$S'_3S'_4$	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_2L_3	L_4L_5	H_7H_{12}	H_8H_{11}	H_9H_{10}	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	H_7H_{12}	H_8H_{11}	H_9H_{10}
dim	1	1	1	1	1	1	2	2	2
P_4	1	0	0	0	0	1	-2	0	0
P_5	1	0	1	0	0	1	0	-2	0
P_6	1	0	0	1	1	0	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	K_7	K_8	K_9	K_{10}	K_{11}	K_{12}	H_7H_{12}	H_8H_{11}	H_9H_{10}	Bipartite graph
	1	1	1	1	1	1	2	2	2	
v^1	0	0	0	0	0	2	1	0	0	$H_{3/2}^1$
v^2	0	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^3	0	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^4	0	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^5	0	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^7	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^8	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^9	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$

SG 189

Space group $P\bar{6}2m$, No. 189, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7	Γ_8	Γ_9	A_7	A_8	A_9
dim		2	2	2	2	2	2
Δ_4	2	0	1	1	0	-1	-1
Δ_5	1	1	0	0	-1	0	0
Δ_6	1	1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_7	Γ_8	Γ_9	A_7	A_8	A_9	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_7	Γ_8	Γ_9	K_7	K_8	K_9
dim	2	2	2	2	2	2
T_5	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_7	Γ_8	Γ_9	K_7	K_8	K_9	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep	Γ_7	Γ_8	Γ_9	M_5
dim	2	2	2	2
Σ_3	1	1	1	1
Σ_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_5
	dim	2	2
U_3	1	1	-1
U_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_5	K_7	K_8	K_9
	dim	2	2	2	2
T'_5	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_5	K_7	K_8	K_9	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_7	A_8	A_9	H_7	H_8	H_9
dim	2	2	2	2	2	2
S_5	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_7	A_8	A_9	H_7	H_8	H_9	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_7	A_8	A_9	L_5
dim	2	2	2	2
R_3	1	1	1	-1
R_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7	A_8	A_9	L_5	Bipartite graph
	2	2	2	2	
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_5	H_7	H_8	H_9
	dim	2	2	2	2
S'_5	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	L_5	H_7	H_8	H_9	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_7	K_8	K_9	H_7	H_8	H_9
	dim	2	2	2	2	2	2
P_4	2	0	1	1	0	-1	-1
P_5	1	1	0	0	-1	0	0
P_6	1	1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_7	K_8	K_9	H_7	H_8	H_9	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/2}^1$

SG 190

Space group $P\bar{6}2c$, No. 190, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix



corep		Γ_7	Γ_8	Γ_9	A_4A_6	A_5A_7	A_8A_9
	dim	2	2	2	2	2	4
Δ_4	2	0	1	1	0	0	-2
Δ_5	1	1	0	0	-2	0	0
Δ_6	1	1	0	0	0	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_7	Γ_8	Γ_9	A_4A_6	A_5A_7	A_8A_9	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	2	0	0	0	1	$H_{3/2}^1$
v^4	2	0	0	1	1	0	$H_{4/4}^2$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	K_7	K_8	K_9
	dim	2	2	2	2	2	2
T_5	2	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_7	Γ_8	Γ_9	K_7	K_8	K_9	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ: $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_7	Γ_8	Γ_9	M_5
	dim	2	2	2	2
Σ_3	1	1	1	1	-1
Σ_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_7	Γ_8	Γ_9	M_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	L_2L_3	L_4L_5
	dim	2	2	2
U_3	1	1	0	-2
U_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	M_5	L_2L_3	L_4L_5	
	2	2	2	Bipartite graph
v^1	2	1	1	$H_{4/4}^2$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_5	K_7	K_8	K_9
	dim	2	2	2	2
T'_5	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_5	K_7	K_8	K_9	
	2	2	2	2	Bipartite graph
v^1	1	0	0	1	$H_{2/1}^1$
v^2	1	0	1	0	$H_{2/1}^1$
v^3	1	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_4A_6	A_5A_7	A_8A_9	H_4	H_5	H_6	H_7	H_8	H_9
dim		2	2	4	1	1	1	1	2	2
S_2	1	1	0	1	-1	0	0	0	-1	0
S_3	1	0	1	1	0	0	0	-1	0	-1
S_4	1	1	0	1	0	0	-1	0	0	-1
S_5	1	0	1	1	0	-1	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	A_4A_6	A_5A_7	A_8A_9	H_4	H_5	H_6	H_7	H_8	H_9	
	2	2	4	1	1	1	1	2	2	Bipartite graph
v^1	0	0	1	0	0	0	0	1	1	$H_{3/4}^1$
v^2	0	0	1	0	0	1	1	1	0	$H_{4/4}^1$
v^3	0	0	1	1	1	0	0	0	1	$H_{4/4}^1$
v^4	0	0	1	1	1	1	1	0	0	$H_{5/4}^1$
v^5	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^6	1	0	0	1	0	1	0	0	0	$H_{3/2}^1$
v^7	1	1	0	0	0	0	0	1	1	$H_{4/4}^2$
v^8	1	1	0	0	0	1	1	1	0	$H_{5/4}^3$
v^9	1	1	0	1	1	0	0	0	1	$H_{5/4}^3$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_4A_6	A_5A_7	A_8A_9	L_2L_3	L_4L_5
dim		2	2	4	2	2
R_3R_4	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	A_4A_6	A_5A_7	A_8A_9	L_2L_3	L_4L_5	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S' : $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_2L_3	L_4L_5	H_4	H_5	H_6	H_7	H_8	H_9
	dim	2	2	1	1	1	1	2	2
S'_2	1	1	0	0	-1	0	0	-1	0
S'_3	1	0	1	0	0	-1	0	0	-1
S'_4	1	1	0	0	0	0	-1	0	-1
S'_5	1	0	1	-1	0	0	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	L_2L_3	L_4L_5	H_4	H_5	H_6	H_7	H_8	H_9	Bipartite graph
	2	2	1	1	1	1	2	2	
v^1	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^2	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^3	1	1	0	0	0	0	1	1	$H_{4/4}^2$
v^4	1	1	0	0	1	1	1	0	$H_{5/4}^3$
v^5	1	1	1	1	0	0	0	1	$H_{5/4}^3$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of P : $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_7	K_8	K_9	H_4	H_5	H_6	H_7	H_8	H_9
	dim	2	2	2	1	1	1	1	2	2
P_4	2	0	1	1	0	0	0	0	-1	-1
P_5	1	1	0	0	-1	0	-1	0	0	0
P_6	1	1	0	0	0	-1	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	K_7	K_8	K_9	H_4	H_5	H_6	H_7	H_8	H_9	Bipartite graph
	2	2	2	1	1	1	1	2	2	
v^1	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	0	1	1	0	0	$H_{3/2}^1$
v^6	1	0	0	0	1	1	0	0	0	$H_{3/2}^1$
v^7	1	0	0	1	0	0	1	0	0	$H_{3/2}^1$
v^8	1	0	0	1	1	0	0	0	0	$H_{3/2}^1$

SG 191

Space group $P6/mmm$, No. 191, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$

A matrix

corep		Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_7^+	A_7^-	A_8^+	A_8^-	A_9^+	A_9^-
dim		2	2	2	2	2	2	2	2	2	2	2	2
Δ_7	2	1	1	0	0	0	0	-1	-1	0	0	0	0
Δ_8	2	0	0	1	1	0	0	0	0	-1	-1	0	0
Δ_9	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_7^+	A_7^-	A_8^+	A_8^-	A_9^+	A_9^-	Bipartite graph
	2	2	2	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

v^{11}	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	0	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9
dim	2	2	2	2	2	2	2	2	2
T_5	2	1	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9	Bipartite graph
v^1	2	2	2	2	2	2	2	2	2	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-
dim	2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-	Bipartite graph
	2	2	2	2	2	2	2		
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_5^+	M_5^-	L_5^+	L_5^-
dim	2	2	2	2
U_5	2	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_5^+	M_5^-	L_5^+	L_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep		M_5^+	M_5^-	K_7	K_8	K_9
	dim	2	2	2	2	2
T'_5	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_5^+	M_5^-	K_7	K_8	K_9	Bipartite graph
	2	2	2	2	2	
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		A_7^+	A_7^-	A_8^+	A_8^-	A_9^+	A_9^-	H_7	H_8	H_9
	dim	2	2	2	2	2	2	2	2	2
S_5	2	1	1	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	A_7^+	A_7^-	A_8^+	A_8^-	A_9^+	A_9^-	H_7	H_8	H_9	Bipartite graph
	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$

v^{16}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		A_7^+	A_7^-	A_8^+	A_8^-	A_9^+	A_9^-	L_5^+	L_5^-
dim		2	2	2	2	2	2	2	2
R_5	2	1	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	A_7^+	A_7^-	A_8^+	A_8^-	A_9^+	A_9^-	L_5^+	L_5^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		L_5^+	L_5^-	H_7	H_8	H_9
dim		2	2	2	2	2
S'_5	2	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_5^+	L_5^-	H_7	H_8	H_9	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_7	K_8	K_9	H_7	H_8	H_9
	dim	2	2	2	2	2	2
P_4	2	0	1	1	0	-1	-1
P_5P_6	2	1	0	0	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_7	K_8	K_9	H_7	H_8	H_9	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

SG 192

Space group $P6/mcc$, No. 192, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_7A_8	A_9A_{11}	$A_{10}A_{12}$
	dim	2	2	2	2	2	2	4	4	4
Δ_7	2	1	1	0	0	0	0	-2	0	0
Δ_8	2	0	0	1	1	0	0	0	-2	0
Δ_9	2	0	0	0	0	1	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_7A_8	A_9A_{11}	$A_{10}A_{12}$	Bipartite graph
	2	2	2	2	2	2	4	4	4	
v^1	0	0	0	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^6	0	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	2	0	0	0	0	1	0	0	$H_{3/2}^1$
v^8	1	1	0	0	0	0	1	0	0	$H_{3/2}^1$
v^9	2	0	0	0	0	0	1	0	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9
dim	2	2	2	2	2	2	2	2	2
T_5	2	1	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9	Bipartite graph
	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$

v^{16}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-
dim		2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5^+	M_5^-	L_3L_4
dim		2	2	4
U_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_5^+	M_5^-	L_3L_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_5^+	M_5^-	K_7	K_8	K_9
dim	2	2	2	2	2
T'_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_5^+	M_5^-	K_7	K_8	K_9	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_7A_8	A_9A_{11}	$A_{10}A_{12}$	H_4H_7	H_5H_6	H_8H_9
dim	4	4	4	2	2	4
S_2S_3	2	1	1	1	-1	0
S_4S_5	2	1	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

A_7A_8	A_9A_{11}	$A_{10}A_{12}$	H_4H_7	H_5H_6	H_8H_9	
4	4	4	2	2	4	Bipartite graph

v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	1	1	0	$H_{3/2}^1$
v^3	0	1	0	0	0	1	$H_{2/2}^1$
v^4	0	1	0	1	1	0	$H_{3/2}^1$
v^5	1	0	0	0	0	1	$H_{2/2}^1$
v^6	1	0	0	1	1	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		A_7A_8	A_9A_{11}	$A_{10}A_{12}$	L_3L_4
	dim	4	4	4	4
R_2R_3	2	1	1	1	-1
R_4R_5	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	A_7A_8	A_9A_{11}	$A_{10}A_{12}$	L_3L_4	
	4	4	4	4	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep		L_3L_4	H_4H_7	H_5H_6	H_8H_9
	dim	4	2	2	4
$S'_2S'_3$	2	1	0	-1	-1
$S'_4S'_5$	2	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_4H_7	H_5H_6	H_8H_9	
	4	2	2	4	Bipartite graph
v^1	1	0	0	1	$H_{2/2}^1$
v^2	1	1	1	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_7	K_8	K_9	H_4H_7	H_5H_6	H_8H_9
	dim	2	2	2	2	2	4
P_4	2	0	1	1	0	0	-2
P_5P_6	2	1	0	0	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	K_7	K_8	K_9	H_4H_7	H_5H_6	H_8H_9	
	2	2	2	2	2	4	Bipartite graph
v^1	0	0	2	0	0	1	$H_{3/2}^1$
v^2	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	2	0	0	0	1	$H_{3/2}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

SG 193

Space group $P6_3/mcm$, No. 193, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_4A_5	A_6
	dim	2	2	2	2	2	2	4	4
Δ_7	2	1	1	0	0	0	0	-2	0
Δ_8	2	0	0	1	1	0	0	0	-1
Δ_9	2	0	0	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_4A_5	A_6	
	2	2	2	2	2	2	4	4	Bipartite graph
v^1	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^6	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	2	0	0	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T : $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9
dim	2	2	2	2	2	2	2	2	2
T_5	2	1	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9	Bipartite graph
v^1	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-
dim	2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U: $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_5^+	M_5^-	L_3L_4
dim	2	2	4
U_5	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_5^+	M_5^-	L_3L_4	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$

A matrix

corep	M_5^+	M_5^-	K_7	K_8	K_9
dim	2	2	2	2	2
T'_5	2	1	1	-1	-1
				-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_5^+	M_5^-	K_7	K_8	K_9	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_4A_5	A_6	H_7H_7	H_8H_9
dim	4	4	4	4
S_5S_5	4	1	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	A_4A_5	A_6	H_7H_7	H_8H_9	
	4	4	4	4	Bipartite graph
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	A_4A_5	A_6	L_3L_4
dim	4	4	4
R_2R_3	2	1	1 -1
R_4R_5	2	1	1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_5	A_6	L_3L_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of *S'*: $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	L_3L_4	H_7H_7	H_8H_9
dim	4	4	4
$S'_5S'_5$	4	1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_3L_4	H_7H_7	H_8H_9	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	K_7	K_8	K_9	H_7H_7	H_8H_9
dim	2	2	2	4	4
P_4	2	0	1	1	0 -2
P_5P_6	2	1	0	0	-2 0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	K_7	K_8	K_9	H_7H_7	H_8H_9	Bipartite graph
v^1	0	0	2	0	1	$H_{3/2}^1$
v^2	0	1	1	0	1	$H_{3/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	2	0	0	1	0	$H_{3/2}^1$

SG 194

Space group $P6_3/mmc$, No. 194, with SOC

Δ

High symmetry line $\Delta(0, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), A(0, 0, \frac{1}{2})$
A matrix

corep		Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_4A_5	A_6
	dim	2	2	2	2	2	2	4	4
Δ_7	2	1	1	0	0	0	0	-2	0
Δ_8	2	0	0	1	1	0	0	0	-1
Δ_9	2	0	0	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	A_4A_5	A_6	Bipartite graph
	2	2	2	2	2	2	4	4	
v^1	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^3	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^4	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^5	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^6	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^7	2	0	0	0	0	0	1	0	$H_{3/2}^1$

T

High symmetry line $T(-u, 2u, 0)$, high symmetry points at both ends of T: $\Gamma(0, 0, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep		Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9
	dim	2	2	2	2	2	2	2	2	2
T_5	2	1	1	1	1	1	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 18.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	K_7	K_8	K_9	Bipartite graph
	2	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$

v^{13}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(0, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-
dim		2	2	2	2	2	2	2	2
Σ_5	2	1	1	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	Γ_9^+	Γ_9^-	M_5^+	M_5^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

U

High symmetry line $U(0, \frac{1}{2}, u)$, high symmetry points at both ends of U : $M(0, \frac{1}{2}, 0), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5^+	M_5^-	L_3L_4
dim		2	2	4
U_5	2	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_5^+	M_5^-	L_3L_4	
	2	2	4	Bipartite graph
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T'

High symmetry line $T'(-2u, u + \frac{1}{2}, 0)$, high symmetry points at both ends of T': $M(0, \frac{1}{2}, 0), K(-\frac{1}{3}, \frac{2}{3}, 0)$
A matrix

corep	M_5^+	M_5^-	K_7	K_8	K_9
dim	2	2	2	2	2
T'_5	2	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_5^+	M_5^-	K_7	K_8	K_9	
	2	2	2	2	2	Bipartite graph
v^1	0	1	0	0	1	$H_{2/1}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	0	1	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$
v^5	1	0	0	1	0	$H_{2/1}^1$
v^6	1	0	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(-u, 2u, \frac{1}{2})$, high symmetry points at both ends of S: $A(0, 0, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep	A_4A_5	A_6	H_4H_5	H_6H_7	H_8	H_9
dim	4	4	2	2	2	2
S_2S_3	2	1	1	-1	0	-1
S_4S_5	2	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

A_4A_5	A_6	H_4H_5	H_6H_7	H_8	H_9	
4	4	2	2	2	2	Bipartite graph

v^1	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	1	0	0	$H_{3/2}^1$

R

High symmetry line $R(0, u, \frac{1}{2})$, high symmetry points at both ends of R: $A(0, 0, \frac{1}{2}), L(0, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	A_4A_5	A_6	L_3L_4	
dim	4	4	4	
R_5R_5	4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	A_4A_5	A_6	L_3L_4	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

S'

High symmetry line $S'(-2u, u + \frac{1}{2}, \frac{1}{2})$, high symmetry points at both ends of S': $L(0, \frac{1}{2}, \frac{1}{2}), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$

A matrix

corep	L_3L_4	H_4H_5	H_6H_7	H_8	H_9	
dim	4	2	2	2	2	
$S'_2S'_3$	2	1	-1	0	-1	0
$S'_4S'_5$	2	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_3L_4	H_4H_5	H_6H_7	H_8	H_9	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	1	1	$H_{3/2}^1$
v^2	1	0	1	1	0	$H_{3/2}^1$
v^3	1	1	0	0	1	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$

P

High symmetry line $P(-\frac{1}{3}, \frac{2}{3}, u)$, high symmetry points at both ends of *P*: $K(-\frac{1}{3}, \frac{2}{3}, 0), H(-\frac{1}{3}, \frac{2}{3}, \frac{1}{2})$
A matrix

corep		K_7	K_8	K_9	H_4H_5	H_6H_7	H_8	H_9
	dim	2	2	2	2	2	2	2
P_4	2	0	1	1	0	0	-1	-1
P_5P_6	2	1	0	0	-1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	K_7	K_8	K_9	H_4H_5	H_6H_7	H_8	H_9	Bipartite graph
	2	2	2	2	2	2	2	
v^1	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	0	0	0	1	$H_{2/1}^1$
v^4	0	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	1	0	0	1	0	0	0	$H_{2/1}^1$

SG 195

Space group *P23*, No. 195, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	X_5
	dim	2	4	2
Δ_3	1	1	2	-1
Δ_4	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	X_5	Bipartite graph
	2	4	2	
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	Γ_5	$\Gamma_6\Gamma_7$	M_5
dim	2	4	2
Σ_2	1	2	4
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	M_5	Bipartite graph
	2	4	2	
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_5	$\Gamma_6\Gamma_7$	R_5	R_6R_7
dim	2	4	2	4
Λ_4	1	1	1	-1
Λ_5	1	1	1	-1
Λ_6	1	0	2	0
				-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	R_5	R_6R_7	Bipartite graph
	2	4	2	4	
v^1	0	1	0	1	$H_{2/4}^1$
v^2	1	0	1	0	$H_{2/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_5	R_5	R_6R_7
dim	2	2	4
S_2	1	2	-2
			-4

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	R_5	R_6R_7	
	2	2	4	Bipartite graph
v^1	1	1	0	$H_{2/2}^1$
v^2	2	0	1	$H_{3/4}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_5	M_5
dim		2	2
Z_3	1	1	-1
Z_4	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	M_5	
	2	2	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_5	R_5	R_6R_7
dim		2	2	4
T_3	1	1	-1	-2
T_4	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	R_5	R_6R_7	
	2	2	4	Bipartite graph
v^1	1	1	0	$H_{2/2}^1$
v^2	2	0	1	$H_{3/4}^1$

SG 196

Space group $F23$, No. 196, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	X_5
dim		2	4	2
Δ_3	1	1	2	-1
Δ_4	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	X_5	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	L_4L_5	L_6L_6
dim		2	4	2	2
Λ_4	1	1	1	-1	0
Λ_5	1	1	1	-1	0
Λ_6	1	0	2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	L_4L_5	L_6L_6	
	2	4	2	2	Bipartite graph
v^1	0	1	1	1	$H_{3/4}^1$
v^2	1	0	1	0	$H_{2/2}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		X_5	W_3	W_4
dim		2	1	1
Z_3	1	1	-1	0
Z_4	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	W_3	W_4	
	2	1	1	Bipartite graph
v^1	1	1	1	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q : $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		L_4L_5	L_6L_6	W_3	W_4
dim		2	2	1	1
Q_2	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	L_4L_5	L_6L_6	W_3	W_4	
	2	2	1	1	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	2	$H_{3/2}^1$
v^5	1	0	1	1	$H_{3/2}^1$
v^6	1	0	2	0	$H_{3/2}^1$

SG 197

Space group $I23$, No. 197, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	N_3N_4
dim		2	4	2
Σ_2	1	2	4	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	N_3N_4	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	H_5	H_6H_7
dim		2	4	2	4
Δ_3	1	1	2	-1	-2
Δ_4	1	1	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5	$\Gamma_6\Gamma_7$	H_5	H_6H_7	Bipartite graph
	2	4	2	4	
v^1	0	1	0	1	$H_{2/4}^1$
v^2	0	1	2	0	$H_{3/4}^1$
v^3	1	0	1	0	$H_{2/2}^1$
v^4	2	0	0	1	$H_{3/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	P_5	P_6	P_7
dim		2	4	2	2	2
Λ_4	1	1	1	-1	-1	0
Λ_5	1	1	1	-1	0	-1
Λ_6	1	0	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	P_5	P_6	P_7	Bipartite graph
	2	4	2	2	2	
v^1	0	1	0	1	1	$H_{3/4}^1$
v^2	1	0	1	0	0	$H_{2/2}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$
A matrix

corep		H_5	H_6H_7	N_3N_4
	dim	2	4	2
G_2	1	2	4	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	H_5	H_6H_7	N_3N_4	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_5	H_6H_7	P_5	P_6	P_7
	dim	2	4	2	2	2
F_4	1	1	1	-1	0	-1
F_5	1	1	1	-1	-1	0
F_6	1	0	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	H_5	H_6H_7	P_5	P_6	P_7	
	2	4	2	2	2	Bipartite graph
v^1	0	1	0	1	1	$H_{3/4}^1$
v^2	1	0	1	0	0	$H_{2/2}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_5	P_6	P_7	N_3N_4
	dim	2	2	2	2
D_3	1	1	1	1	-1
D_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_5	P_6	P_7	N_3N_4	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

SG 198

Space group $P2_13$, No. 198, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	X_2X_3	X_4X_5
	dim	2	4	2	2
Δ_3	1	1	2	0	-2
Δ_4	1	1	2	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	X_2X_3	X_4X_5	
	2	4	2	2	Bipartite graph
v^1	0	1	1	1	$H_{3/4}^1$
v^2	2	0	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	M_5M_5
	dim	2	4	4
Σ_2	1	2	4	-4

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	M_5M_5	
	2	4	4	Bipartite graph
v^1	0	1	1	$H_{2/4}^1$
v^2	2	0	1	$H_{3/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	R_4R_4	R_5R_6	R_7R_7
	dim	2	4	2	2	6
Λ_4	1	1	1	0	-1	-2
Λ_5	1	1	1	0	-1	-2
Λ_6	1	0	2	-2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5	$\Gamma_6\Gamma_7$	R_4R_4	R_5R_6	R_7R_7	
	2	4	2	2	6	Bipartite graph
v^1	0	1	1	1	0	$H_{3/4}^1$
v^2	0	2	1	0	1	$H_{4/8}^1$
v^3	1	0	0	1	0	$H_{2/2}^1$
v^4	1	1	0	0	1	$H_{3/6}^2$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_2X_3	X_4X_5	R_4R_4	R_5R_6	R_7R_7
	dim	2	2	2	2	6
S_2S_2	2	1	1	-1	-1	-3

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_2X_3	X_4X_5	R_4R_4	R_5R_6	R_7R_7	
	2	2	2	2	6	Bipartite graph
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	0	1	1	0	0	$H_{2/1}^1$
v^3	0	3	0	0	1	$H_{4/3}^1$
v^4	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	1	0	0	$H_{2/1}^1$
v^6	1	2	0	0	1	$H_{4/3}^1$
v^7	2	1	0	0	1	$H_{4/3}^1$
v^8	3	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_2X_3	X_4X_5	M_5M_5
dim	2	2	4
Z_3Z_4	2	1	1
			-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_2X_3	X_4X_5	M_5M_5	Bipartite graph
	2	2	4	
v^1	0	2	1	$H_{3/2}^1$
v^2	1	1	1	$H_{3/2}^1$
v^3	2	0	1	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_5M_5	R_4R_4	R_5R_6	R_7R_7
dim	4	2	2	6
T_3T_3	2	1	0	0
T_4T_4	2	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5M_5	R_4R_4	R_5R_6	R_7R_7	Bipartite graph
	4	2	2	6	
v^1	2	0	1	1	$H_{4/4}^4$
v^2	2	1	0	1	$H_{4/4}^4$

SG 199

Space group $I2_13$, No. 199, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix



corep		Γ_5	$\Gamma_6\Gamma_7$	N_3N_4
	dim	2	4	2
Σ_2	1	2	4	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_5	$\Gamma_6\Gamma_7$	N_3N_4	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	H_5	H_6H_7
	dim	2	4	2	4
Δ_3	1	1	2	-1	-2
Δ_4	1	1	2	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5	$\Gamma_6\Gamma_7$	H_5	H_6H_7	
	2	4	2	4	Bipartite graph
v^1	0	1	0	1	$H_{2/4}^1$
v^2	0	1	2	0	$H_{3/4}^1$
v^3	1	0	1	0	$H_{2/2}^1$
v^4	2	0	0	1	$H_{3/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_5	$\Gamma_6\Gamma_7$	P_4	P_5	P_6	P_7
	dim	2	4	1	1	1	3
Λ_4	1	1	1	0	0	-1	-1
Λ_5	1	1	1	0	-1	0	-1
Λ_6	1	0	2	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5	$\Gamma_6\Gamma_7$	P_4	P_5	P_6	P_7	
	2	4	1	1	1	3	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/4}^2$
v^2	0	1	2	1	1	0	$H_{5/4}^1$
v^3	1	0	0	1	1	0	$H_{3/2}^1$
v^4	1	1	0	0	0	2	$H_{4/6}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep		H_5	H_6H_7	N_3N_4
	dim	2	4	2
G_2	1	2	4	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	H_5	H_6H_7	N_3N_4	
	2	4	2	Bipartite graph
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		H_5	H_6H_7	P_4	P_5	P_6	P_7
	dim	2	4	1	1	1	3
F_4	1	0	2	-1	0	0	-1
F_5	1	1	1	0	0	-1	-1
F_6	1	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	H_5	H_6H_7	P_4	P_5	P_6	P_7	
	2	4	1	1	1	3	Bipartite graph
v^1	0	1	1	0	0	1	$H_{3/4}^2$
v^2	0	1	2	1	1	0	$H_{5/4}^1$
v^3	1	0	0	1	1	0	$H_{3/2}^1$
v^4	1	1	0	0	0	2	$H_{4/6}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_4	P_5	P_6	P_7	N_3N_4
	dim	1	1	1	3	2
D_3	1	0	0	0	2	-1
D_4	1	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_4	P_5	P_6	P_7	N_3N_4	
	1	1	1	3	2	Bipartite graph
v^1	0	0	1	1	2	$H_{4/4}^4$
v^2	0	1	0	1	2	$H_{4/4}^4$
v^3	1	0	0	1	2	$H_{4/4}^4$

SG 200

Space group $Pm\bar{3}$, No. 200, with SOC

 Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	X_5^+	X_5^-
	dim	2	2	4	4	2	2
Δ_5	2	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	X_5^+	X_5^-	
	2	2	4	4	2	2	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	M_5^+	M_5^-	
dim	2	2	4	4	2	2	
$\Sigma_3\Sigma_4$	2	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	M_5^+	M_5^-	Bipartite graph
	2	2	4	4	2	2	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$
dim	2	2	4	4	2	2	4	4
$\Lambda_4\Lambda_5$	2	1	1	1	-1	-1	-1	-1
$\Lambda_6\Lambda_6$	2	0	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$	Bipartite graph
	2	2	4	4	2	2	4	4	
v^1	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/2}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/2}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_5^+	X_5^-	R_5^+	R_5^-	$R_6^+ R_7^+$	$R_6^- R_7^-$
dim	2	2	2	2	4	4
$S_3 S_4$	2	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	X_5^+	X_5^-	R_5^+	R_5^-	$R_6^+ R_7^+$	$R_6^- R_7^-$	Bipartite graph
	2	2	2	2	4	4	
v^1	0	1	0	1	0	0	$H_{2/1}^1$
v^2	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$
v^6	1	0	1	0	0	0	$H_{2/1}^1$
v^7	1	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_5^+	X_5^-	M_5^+	M_5^-
dim	2	2	2	2
Z_5	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_5^+	X_5^-	M_5^+	M_5^-	Bipartite graph
	2	2	2	2	
v^1	0	1	0	1	$H_{2/1}^1$
v^2	0	1	1	0	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$
v^4	1	0	1	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5^+	M_5^-	R_5^+	R_5^-	$R_6^+ R_7^+$	$R_6^- R_7^-$
dim		2	2	2	2	4	4
T_5	2	1	1	-1	-1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	M_5^+	M_5^-	R_5^+	R_5^-	$R_6^+ R_7^+$	$R_6^- R_7^-$	Bipartite graph
	2	2	2	2	4	4	
v^1	0	1	0	1	0	0	$H_{2/1}^1$
v^2	0	1	1	0	0	0	$H_{2/1}^1$
v^3	0	2	0	0	0	1	$H_{3/2}^1$
v^4	0	2	0	0	1	0	$H_{3/2}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$
v^6	1	0	1	0	0	0	$H_{2/1}^1$
v^7	1	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	0	$H_{3/2}^1$

SG 201

Space group $Pn\bar{3}$, No. 201, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	$\Gamma_6^+ \Gamma_7^+$	$\Gamma_6^- \Gamma_7^-$	$X_3 X_4$
dim		2	2	4	4	4
Δ_5	2	1	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_5^+	Γ_5^-	$\Gamma_6^+ \Gamma_7^+$	$\Gamma_6^- \Gamma_7^-$	$X_3 X_4$	Bipartite graph
	2	2	4	4	4	
v^1	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	0	1	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$
v^5	2	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	M_3M_4	
dim	2	2	4	4	4	
$\Sigma_3\Sigma_4$	2	1	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	M_3M_4	Bipartite graph
	2	2	4	4	4	
v^1	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	0	1	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$
v^5	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$
dim	2	2	4	4	2	2	4	4
$\Lambda_4\Lambda_5$	2	1	1	1	-1	-1	-1	-1
$\Lambda_6\Lambda_6$	2	0	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$	Bipartite graph
	2	2	4	4	2	2	4	4	
v^1	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/2}^1$
v^3	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^4	0	0	1	0	0	0	1	0	$H_{2/2}^1$
v^5	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	1	0	0	0	1	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$
dim	4	2	2	4	4
S_3S_4	2	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	X_3X_4	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$	Bipartite graph
	4	2	2	4	4	
v^1	1	0	0	0	1	$H_{2/2}^1$
v^2	1	0	0	1	0	$H_{2/2}^1$
v^3	1	0	2	0	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$
v^5	1	2	0	0	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	M_3M_4
dim	4	4
Z_2Z_5	2	1
Z_3Z_4	2	1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	M_3M_4	Bipartite graph
	4	4	
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_3M_4	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$
dim	4	2	2	4	4
T_5	2	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	M_3M_4	R_5^+	R_5^-	$R_6^+R_7^+$	$R_6^-R_7^-$	
	4	2	2	4	4	Bipartite graph
v^1	1	0	0	0	1	$H_{2/2}^1$
v^2	1	0	0	1	0	$H_{2/2}^1$
v^3	1	0	2	0	0	$H_{3/2}^1$
v^4	1	1	1	0	0	$H_{3/2}^1$
v^5	1	2	0	0	0	$H_{3/2}^1$

SG 202

Space group $Fm\bar{3}$, No. 202, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	X_5^+	X_5^-
dim	2	2	4	4	2	2
Δ_5	2	1	1	2	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	X_5^+	X_5^-	
	2	2	4	4	2	2	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	$L_4^+L_4^+$	$L_4^-L_4^-$	$L_5^+L_6^+$	$L_5^-L_6^-$
dim	2	2	4	4	2	2	2	2
$\Lambda_4\Lambda_5$	2	1	1	1	0	0	-1	-1
$\Lambda_6\Lambda_6$	2	0	0	1	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	$L_4^+L_4^+$	$L_4^-L_4^-$	$L_5^+L_6^+$	$L_5^-L_6^-$	Bipartite graph
v^1	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z: $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_5^+	X_5^-	W_5
dim	2	2	2
Z_5	2	1	1
			-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5^+	X_5^-	W_5	Bipartite graph
	2	2	2	
v^1	0	1	1	$H_{2/1}^1$
v^2	1	0	1	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	$L_4^+ L_4^+$	$L_4^- L_4^-$	$L_5^+ L_6^+$	$L_5^- L_6^-$	W_5
dim	2	2	2	2	2
$Q_2 Q_2$	2	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	$L_4^+ L_4^+$	$L_4^- L_4^-$	$L_5^+ L_6^+$	$L_5^- L_6^-$	W_5	
	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	1	$H_{2/1}^1$
v^2	0	0	1	0	1	$H_{2/1}^1$
v^3	0	1	0	0	1	$H_{2/1}^1$
v^4	1	0	0	0	1	$H_{2/1}^1$

SG 203

Space group $Fd\bar{3}$, No. 203, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+ \Gamma_7^+$	$\Gamma_6^- \Gamma_7^-$	$X_3 X_4$
dim	2	2	4	4	4
Δ_5	2	1	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_5^+	Γ_5^-	$\Gamma_6^+ \Gamma_7^+$	$\Gamma_6^- \Gamma_7^-$	$X_3 X_4$	
	2	2	4	4	4	Bipartite graph
v^1	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	0	1	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$
v^5	2	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	$L_4^+L_4^+$	$L_4^-L_4^-$	$L_5^+L_6^+$	$L_5^-L_6^-$
dim	2	2	4	4	2	2	2	2
$\Lambda_4\Lambda_5$	2	1	1	1	0	0	-1	-1
$\Lambda_6\Lambda_6$	2	0	0	1	1	-1	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	$L_4^+L_4^+$	$L_4^-L_4^-$	$L_5^+L_6^+$	$L_5^-L_6^-$	Bipartite graph
	2	2	4	4	2	2	2	2	
v^1	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z: $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_3X_4	W_2W_3	W_4W_5
dim	4	2	2
Z_2Z_5	2	1	-1
Z_3Z_4	2	1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_3X_4	W_2W_3	W_4W_5	Bipartite graph
	4	2	2	
v^1	1	1	1	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		$L_4^+ L_4^+$	$L_4^- L_4^-$	$L_5^+ L_6^+$	$L_5^- L_6^-$	$W_2 W_3$	$W_4 W_5$
dim		2	2	2	2	2	2
$Q_2 Q_2$	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	$L_4^+ L_4^+$	$L_4^- L_4^-$	$L_5^+ L_6^+$	$L_5^- L_6^-$	$W_2 W_3$	$W_4 W_5$	Bipartite graph
	2	2	2	2	2	2	
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

SG 204

Space group $Im\bar{3}$, No. 204, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep		Γ_5^+	Γ_5^-	$\Gamma_6^+ \Gamma_7^+$	$\Gamma_6^- \Gamma_7^-$	$N_3^+ N_4^+$	$N_3^- N_4^-$
dim		2	2	4	4	2	2
$\Sigma_3 \Sigma_4$	2	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_5^+	Γ_5^-	$\Gamma_6^+ \Gamma_7^+$	$\Gamma_6^- \Gamma_7^-$	$N_3^+ N_4^+$	$N_3^- N_4^-$	Bipartite graph
	2	2	4	4	2	2	
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	H_5^+	H_5^-	$H_6^+H_7^+$	$H_6^-H_7^-$
dim	2	2	4	4	2	2	4	4
Δ_5	2	1	1	2	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	H_5^+	H_5^-	$H_6^+H_7^+$	$H_6^-H_7^-$	Bipartite graph
	2	2	4	4	2	2	4	4	
v^1	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^6	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^7	0	0	1	0	0	0	1	0	$H_{2/2}^1$
v^8	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^9	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{19}	2	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	P_5	P_6P_7
dim	2	2	4	4	2	4

$\Lambda_4\Lambda_5$	2	1	1	1	1	-1	-1
$\Lambda_6\Lambda_6$	2	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	P_5	P_6P_7	
	2	2	4	4	2	4	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep	H_5^+	H_5^-	$H_6^+H_7^+$	$H_6^-H_7^-$	$N_3^+N_4^+$	$N_3^-N_4^-$	
dim	2	2	4	4	2	2	
G_3G_4	2	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	H_5^+	H_5^-	$H_6^+H_7^+$	$H_6^-H_7^-$	$N_3^+N_4^+$	$N_3^-N_4^-$	
	2	2	4	4	2	2	Bipartite graph
v^1	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	1	0	0	0	1	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix



corep		H_5^+	H_5^-	$H_6^+ H_7^+$	$H_6^- H_7^-$	P_5	$P_6 P_7$
dim		2	2	4	4	2	4
$F_4 F_5$	2	1	1	1	1	-1	-1
$F_6 F_6$	2	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	H_5^+	H_5^-	$H_6^+ H_7^+$	$H_6^- H_7^-$	P_5	$P_6 P_7$	
	2	2	4	4	2	4	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	1	0	0	1	$H_{2/2}^1$
v^3	0	1	0	0	1	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_5	$P_6 P_7$	$N_3^+ N_4^+$	$N_3^- N_4^-$
dim		2	4	2	2
$D_3 D_4$	2	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	P_5	$P_6 P_7$	$N_3^+ N_4^+$	$N_3^- N_4^-$	
	2	4	2	2	Bipartite graph
v^1	0	1	0	2	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	1	2	0	$H_{3/2}^1$
v^4	1	0	0	1	$H_{2/1}^1$
v^5	1	0	1	0	$H_{2/1}^1$

SG 205

Space group $Pa\bar{3}$, No. 205, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix



corep		Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	X_3X_4
dim		2	2	4	4	4
Δ_5	2	1	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	X_3X_4	Bipartite graph
	2	2	4	4	4	
v^1	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	0	1	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$
v^5	2	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	M_3M_3	M_4M_4
dim		2	2	4	4	4	4
$\Sigma_3\Sigma_4$	2	1	1	2	2	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	M_3M_3	M_4M_4	Bipartite graph
	2	2	4	4	4	4	
v^1	0	0	0	1	0	1	$H_{2/2}^1$
v^2	0	0	0	1	1	0	$H_{2/2}^1$
v^3	0	0	1	0	0	1	$H_{2/2}^1$
v^4	0	0	1	0	1	0	$H_{2/2}^1$
v^5	0	2	0	0	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	0	$H_{3/2}^1$
v^7	1	1	0	0	0	1	$H_{3/2}^1$
v^8	1	1	0	0	1	0	$H_{3/2}^1$
v^9	2	0	0	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_6^+$	$R_5^-R_6^-$	$R_7^+R_7^+$	$R_7^-R_7^-$
dim	2	2	4	4	2	2	2	2	6	6
$\Lambda_4\Lambda_5$	2	1	1	1	0	0	-1	-1	-2	-2
$\Lambda_6\Lambda_6$	2	0	0	1	1	-1	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 32.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_6^+$	$R_5^-R_6^-$	$R_7^+R_7^+$	$R_7^-R_7^-$	Bipartite graph
	2	2	4	4	2	2	2	2	6	6	
v^1	0	0	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^2	0	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^3	0	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	2	0	1	0	0	0	1	$H_{4/4}^4$
v^6	0	0	0	2	0	1	0	0	1	0	$H_{4/4}^4$
v^7	0	0	0	2	1	0	0	0	0	1	$H_{4/4}^4$
v^8	0	0	0	2	1	0	0	0	1	0	$H_{4/4}^4$
v^9	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{10}	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{12}	0	0	1	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	1	1	0	1	0	0	0	1	$H_{4/4}^4$
v^{14}	0	0	1	1	0	1	0	0	1	0	$H_{4/4}^4$
v^{15}	0	0	1	1	1	0	0	0	0	1	$H_{4/4}^4$
v^{16}	0	0	1	1	1	0	0	0	1	0	$H_{4/4}^4$
v^{17}	0	0	2	0	0	1	0	0	0	1	$H_{4/4}^4$
v^{18}	0	0	2	0	0	1	0	0	1	0	$H_{4/4}^4$
v^{19}	0	0	2	0	1	0	0	0	0	1	$H_{4/4}^4$
v^{20}	0	0	2	0	1	0	0	0	1	0	$H_{4/4}^4$
v^{21}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{22}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{23}	0	1	0	1	0	0	0	0	0	1	$H_{3/3}^1$
v^{24}	0	1	0	1	0	0	0	0	1	0	$H_{3/3}^1$
v^{25}	0	1	1	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{26}	0	1	1	0	0	0	0	0	1	0	$H_{3/3}^1$
v^{27}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{28}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{29}	1	0	0	1	0	0	0	0	0	1	$H_{3/3}^1$
v^{30}	1	0	0	1	0	0	0	0	1	0	$H_{3/3}^1$
v^{31}	1	0	1	0	0	0	0	0	0	1	$H_{3/3}^1$
v^{32}	1	0	1	0	0	0	0	0	1	0	$H_{3/3}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_4	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_6^+$	$R_5^-R_6^-$	$R_7^+R_7^+$	$R_7^-R_7^-$
dim	4	2	2	2	2	6	6
S_3S_3	2	1	0	-1	0	-1	-2
S_4S_4	2	1	-1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	X_3X_4	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_6^+$	$R_5^-R_6^-$	$R_7^+R_7^+$	$R_7^-R_7^-$	
	4	2	2	2	2	6	6	Bipartite graph
v^1	1	0	0	1	1	0	0	$H_{3/2}^1$
v^2	1	0	1	1	0	0	0	$H_{3/2}^1$
v^3	1	1	0	0	1	0	0	$H_{3/2}^1$
v^4	1	1	1	0	0	0	0	$H_{3/2}^1$
v^5	2	0	0	0	1	0	1	$H_{4/4}^4$
v^6	2	0	0	1	0	1	0	$H_{4/4}^4$
v^7	2	0	1	0	0	0	1	$H_{4/4}^4$
v^8	2	1	0	0	0	1	0	$H_{4/4}^4$
v^9	3	0	0	0	0	1	1	$H_{5/6}^2$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_4	M_3M_3	M_4M_4
dim	4	4	4
Z_5Z_5	4	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_3X_4	M_3M_3	M_4M_4	
	4	4	4	Bipartite graph
v^1	1	0	1	$H_{2/1}^1$
v^2	1	1	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_3M_3	M_4M_4	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_6^+$	$R_5^-R_6^-$	$R_7^+R_7^+$	$R_7^-R_7^-$
dim	4	4	2	2	2	2	6	6
T_2T_2	2	0	1	0	0	0	-1	-1
T_3T_3	2	0	1	-1	0	-1	0	-1
T_4T_4	2	1	0	0	-1	0	-1	0
T_5T_5	2	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	M_3M_3	M_4M_4	$R_4^+R_4^+$	$R_4^-R_4^-$	$R_5^+R_6^+$	$R_5^-R_6^-$	$R_7^+R_7^+$	$R_7^-R_7^-$	Bipartite graph
	4	4	2	2	2	2	6	6	
v^1	1	1	0	0	0	1	0	1	$H_{4/4}^4$
v^2	1	1	0	0	1	0	1	0	$H_{4/4}^4$
v^3	1	1	0	1	0	0	0	1	$H_{4/4}^4$
v^4	1	1	1	0	0	0	1	0	$H_{4/4}^4$

SG 206

Space group $Ia\bar{3}$, No. 206, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	N_2N_2
dim	2	2	4	4	4
$\Sigma_3\Sigma_4$	2	1	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	N_2N_2	Bipartite graph
	2	2	4	4	4	
v^1	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	1	0	1	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	1	0	0	1	$H_{3/2}^1$
v^5	2	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	H_5^+	H_5^-	$H_6^+H_7^+$	$H_6^-H_7^-$
dim	2	2	4	4	2	2	4	4	
Δ_5	2	1	1	2	2	-1	-1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	H_5^+	H_5^-	$H_6^+H_7^+$	$H_6^-H_7^-$	Bipartite graph
	2	2	4	4	2	2	4	4	
v^1	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^4	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^5	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^6	0	0	1	0	0	0	0	1	$H_{2/2}^1$
v^7	0	0	1	0	0	0	1	0	$H_{2/2}^1$
v^8	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^9	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{18}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{19}	2	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		Γ_5^+	Γ_5^-	$\Gamma_6^+\Gamma_7^+$	$\Gamma_6^-\Gamma_7^-$	P_4P_4	P_5P_6	P_7P_7
dim	2	2	4	4	2	2	6	
$\Lambda_4\Lambda_5$	2	1	1	1	1	0	-1	-2
$\Lambda_6\Lambda_6$	2	0	0	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	H_5^+	H_5^-	$H_6^+ H_7^+$	$H_6^- H_7^-$	$P_4 P_4$	$P_5 P_6$	$P_7 P_7$	Bipartite graph
	2	2	4	4	2	2	6	
v^1	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	2	1	0	1	$H_{4/4}^4$
v^3	0	0	1	0	1	1	0	$H_{3/2}^1$
v^4	0	0	1	1	1	0	1	$H_{4/4}^4$
v^5	0	0	2	0	1	0	1	$H_{4/4}^4$
v^6	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	1	0	1	0	0	1	$H_{3/3}^1$
v^8	0	1	1	0	0	0	1	$H_{3/3}^1$
v^9	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	1	0	0	1	$H_{3/3}^1$
v^{11}	1	0	1	0	0	0	1	$H_{3/3}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$

A matrix

corep	$P_4 P_4$	$P_5 P_6$	$P_7 P_7$	$N_2 N_2$
dim	2	2	6	4
$D_3 D_3$	2	0	0	2
$D_4 D_4$	2	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$P_4 P_4$	$P_5 P_6$	$P_7 P_7$	$N_2 N_2$	Bipartite graph
	2	2	6	4	
v^1	0	1	1	2	$H_{4/4}^4$
v^2	1	0	1	2	$H_{4/4}^4$

SG 207

Space group $P432$, No. 207, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	X_6	X_7
	dim	2	2	4	2	2
Δ_5	1	1	0	1	-1	0
Δ_6	1	0	1	1	0	-1
Δ_7	1	0	1	1	0	-1
Δ_8	1	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	X_6	X_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	0	1	0	0	1	$H_{2/2}^1$
v^3	1	0	0	1	0	$H_{2/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	M_6	M_7
	dim	2	2	4	2	2
Σ_3	1	1	1	2	-1	-1
Σ_4	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	M_6	M_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/4}^1$
v^2	0	0	1	1	1	$H_{3/4}^1$
v^3	0	0	1	2	0	$H_{3/4}^1$
v^4	0	1	0	0	1	$H_{2/2}^1$
v^5	0	1	0	1	0	$H_{2/2}^1$
v^6	1	0	0	0	1	$H_{2/2}^1$
v^7	1	0	0	1	0	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Γ_8	R_6	R_7	R_8
	dim	2	2	4	2	2	4
Λ_4	1	1	1	1	-1	-1	-1
Λ_5	1	1	1	1	-1	-1	-1
Λ_6	1	0	0	2	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_6	Γ_7	Γ_8	R_6	R_7	R_8	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/4}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	0	1	0	1	0	0	$H_{2/2}^1$
v^4	1	0	0	0	1	0	$H_{2/2}^1$
v^5	1	0	0	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_6	X_7	R_6	R_7	R_8
	dim	2	2	2	2	4
S_3	1	1	1	-1	-1	-2
S_4	1	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_6	X_7	R_6	R_7	R_8	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	0	$H_{2/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/4}^1$
v^4	1	0	0	1	0	$H_{2/2}^1$
v^5	1	0	1	0	0	$H_{2/2}^1$
v^6	1	1	0	0	1	$H_{3/4}^1$
v^7	2	0	0	0	1	$H_{3/4}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_6	X_7	M_6	M_7
dim		2	2	2	2
Z_3	1	1	1	-1	-1
Z_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_6	X_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	R_6	R_7	R_8
dim		2	2	2	2	4
T_5	1	1	0	-1	0	-1
T_6	1	0	1	0	-1	-1
T_7	1	0	1	0	-1	-1
T_8	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	M_6	M_7	R_6	R_7	R_8	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	0	$H_{2/2}^1$
v^2	1	0	1	0	0	$H_{2/2}^1$
v^3	1	1	0	0	1	$H_{3/4}^1$

SG 208

Space group $P4_232$, No. 208, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	Γ_8	X_6	X_7
	dim	2	2	4	2	2
Δ_5	1	1	0	1	-1	0
Δ_6	1	0	1	1	-1	0
Δ_7	1	0	1	1	0	-1
Δ_8	1	1	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Γ_8	X_6	X_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	1	1	0	1	1	$H_{4/4}^2$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	Γ_8	M_6	M_7
	dim	2	2	4	2	2
Σ_3	1	1	1	2	-1	-1
Σ_4	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	M_6	M_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/4}^1$
v^2	0	0	1	1	1	$H_{3/4}^1$
v^3	0	0	1	2	0	$H_{3/4}^1$
v^4	0	1	0	0	1	$H_{2/2}^1$
v^5	0	1	0	1	0	$H_{2/2}^1$
v^6	1	0	0	0	1	$H_{2/2}^1$
v^7	1	0	0	1	0	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix



corep		Γ_6	Γ_7	Γ_8	R_6	R_7	R_8
	dim	2	2	4	2	2	4
Λ_4	1	1	1	1	-1	-1	-1
Λ_5	1	1	1	1	-1	-1	-1
Λ_6	1	0	0	2	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_6	Γ_7	Γ_8	R_6	R_7	R_8	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/4}^1$
v^2	0	1	0	0	1	0	$H_{2/2}^1$
v^3	0	1	0	1	0	0	$H_{2/2}^1$
v^4	1	0	0	0	1	0	$H_{2/2}^1$
v^5	1	0	0	1	0	0	$H_{2/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_6	X_7	R_6	R_7	R_8
	dim	2	2	2	2	4
S_3	1	1	1	-1	-1	-2
S_4	1	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_6	X_7	R_6	R_7	R_8	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	0	$H_{2/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/4}^1$
v^4	1	0	0	1	0	$H_{2/2}^1$
v^5	1	0	1	0	0	$H_{2/2}^1$
v^6	1	1	0	0	1	$H_{3/4}^1$
v^7	2	0	0	0	1	$H_{3/4}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_6	X_7	M_6	M_7
	dim	2	2	2	2
Z_3	1	1	1	-1	-1
Z_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_6	X_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	R_6	R_7	R_8
	dim	2	2	2	2	4
T_5	1	0	1	0	-1	-1
T_6	1	1	0	0	-1	-1
T_7	1	1	0	-1	0	-1
T_8	1	0	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6	M_7	R_6	R_7	R_8	
	2	2	2	2	4	Bipartite graph
v^1	1	1	0	0	1	$H_{3/4}^1$
v^2	1	1	1	1	0	$H_{4/4}^2$

SG 209

Space group $F432$, No. 209, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	X_6	X_7
	dim	2	2	4	2	2
Δ_5	1	1	0	1	-1	0
Δ_6	1	0	1	1	0	-1
Δ_7	1	0	1	1	0	-1
Δ_8	1	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	X_6	X_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	0	1	0	0	1	$H_{2/2}^1$
v^3	1	0	0	1	0	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Γ_8	L_4	L_5L_6
	dim	2	2	4	2	2
Λ_4	1	1	1	1	-1	0
Λ_5	1	1	1	1	-1	0
Λ_6	1	0	0	2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	L_4	L_5L_6	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	0	1	0	1	0	$H_{2/2}^1$
v^3	1	0	0	1	0	$H_{2/2}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		X_6	X_7	W_5
	dim	2	2	2
Z_3	1	1	1	-1
Z_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_6	X_7	W_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q : $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		L_4	L_5L_6	W_5
	dim	2	2	2
Q_3	1	1	1	-1
Q_4	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_4	L_5L_6	W_5	
	2	2	2	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

SG 210

Space group $F4_132$, No. 210, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	X_6	X_7
	dim	2	2	4	2	2
Δ_5	1	1	0	1	0	-1
Δ_6	1	0	1	1	0	-1
Δ_7	1	0	1	1	-1	0
Δ_8	1	1	0	1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	Γ_6	Γ_7	Γ_8	X_6	X_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	1	1	0	1	1	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Γ_8	L_4	L_5L_6
	dim	2	2	4	2	2
Λ_4	1	1	1	1	-1	0
Λ_5	1	1	1	1	-1	0
Λ_6	1	0	0	2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	L_4	L_5L_6	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	0	1	0	1	0	$H_{2/2}^1$
v^3	1	0	0	1	0	$H_{2/2}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		X_6	X_7	W_2W_4	W_3	W_5
	dim	2	2	2	1	1
Z_3	1	1	1	-2	0	0
Z_4	1	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 9.

	X_6	X_7	W_2W_4	W_3	W_5	
	2	2	2	1	1	Bipartite graph
v^1	0	2	1	0	2	$H_{5/4}^3$
v^2	0	2	1	1	1	$H_{5/4}^3$
v^3	0	2	1	2	0	$H_{5/4}^3$
v^4	1	1	1	0	2	$H_{5/4}^3$
v^5	1	1	1	1	1	$H_{5/4}^3$

v^6	1	1	1	2	0	$H_{5/4}^3$
v^7	2	0	1	0	2	$H_{5/4}^3$
v^8	2	0	1	1	1	$H_{5/4}^3$
v^9	2	0	1	2	0	$H_{5/4}^3$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q : $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		L_4	L_5L_6	W_2W_4	W_3	W_5
dim		2	2	2	1	1
Q_3	1	1	1	-1	-1	0
Q_4	1	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_4	L_5L_6	W_2W_4	W_3	W_5	
	2	2	2	1	1	Bipartite graph
v^1	0	1	0	1	1	$H_{3/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	1	$H_{3/2}^1$
v^4	1	0	1	0	0	$H_{2/2}^1$

SG 211

Space group $I432$, No. 211, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Γ_8	N_5
dim		2	2	4	2
Σ_3	1	1	1	2	-1
Σ_4	1	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	N_5	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/4}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	H_6	H_7	H_8
	dim	2	2	4	2	2	4
Δ_5	1	1	0	1	-1	0	-1
Δ_6	1	0	1	1	0	-1	-1
Δ_7	1	0	1	1	0	-1	-1
Δ_8	1	1	0	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_6	Γ_7	Γ_8	H_6	H_7	H_8	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/4}^1$
v^2	0	0	1	1	1	0	$H_{3/4}^1$
v^3	0	1	0	0	1	0	$H_{2/2}^1$
v^4	1	0	0	1	0	0	$H_{2/2}^1$
v^5	1	1	0	0	0	1	$H_{3/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_6	Γ_7	Γ_8	P_5	P_6	P_7
	dim	2	2	4	2	2	2
Λ_4	1	1	1	1	-1	-1	0
Λ_5	1	1	1	1	-1	0	-1
Λ_6	1	0	0	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	P_5	P_6	P_7	
	2	2	4	2	2	2	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/4}^1$
v^2	0	1	0	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	0	0	$H_{2/2}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$
A matrix

corep		H_6	H_7	H_8	N_5
	dim	2	2	4	2
G_3	1	1	1	2	-1
G_4	1	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	H_6	H_7	H_8	N_5	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/4}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		H_6	H_7	H_8	P_5	P_6	P_7
	dim	2	2	4	2	2	2
F_4	1	1	1	1	-1	0	-1
F_5	1	1	1	1	-1	-1	0
F_6	1	0	0	2	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	H_6	H_7	H_8	P_5	P_6	P_7	
	2	2	4	2	2	2	Bipartite graph
v^1	0	0	1	0	1	1	$H_{3/4}^1$
v^2	0	1	0	1	0	0	$H_{2/2}^1$
v^3	1	0	0	1	0	0	$H_{2/2}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$

A matrix

corep		P_5	P_6	P_7	N_5
	dim	2	2	2	2
D_3	1	1	1	1	-1
D_4	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_5	P_6	P_7	N_5	
	2	2	2	2	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

SG 212

Space group $P4_332$, No. 212, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	X_3X_5	X_4X_6	X_7
dim		2	2	4	2	2	2
Δ_5	1	1	0	1	0	0	-1
Δ_6	1	0	1	1	0	-2	0
Δ_7	1	0	1	1	0	0	-1
Δ_8	1	1	0	1	-2	0	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	X_3X_5	X_4X_6	X_7	
	2	2	4	2	2	2	Bipartite graph
v^1	0	0	2	1	1	2	$H_{6/8}^1$
v^2	1	1	1	1	1	2	$H_{7/8}^{11}, H_{7/8}^9$
v^3	2	2	0	1	1	2	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	M_6M_7
dim		2	2	4	4
Σ_3	1	1	1	2	-2
Σ_4	1	1	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Γ_8	M_6M_7	
	2	2	4	4	Bipartite graph
v^1	0	0	1	1	$H_{2/4}^1$
v^2	0	2	0	1	$H_{3/4}^1$
v^3	1	1	0	1	$H_{3/4}^1$
v^4	2	0	0	1	$H_{3/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_6	Γ_7	Γ_8	R_4R_5	R_6	R_7R_8
dim	2	2	4	2	2	6
Λ_4	1	1	1	1	0	-1 -2
Λ_5	1	1	1	1	0	-1 -2
Λ_6	1	0	0	2	-2	0 -2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6	Γ_7	Γ_8	R_4R_5	R_6	R_7R_8	
	2	2	4	2	2	6	Bipartite graph
v^1	0	0	1	1	1	0	$H_{3/4}^1$
v^2	0	0	2	1	0	1	$H_{4/8}^1$
v^3	0	1	0	0	1	0	$H_{2/2}^1$
v^4	0	1	1	0	0	1	$H_{3/6}^2$
v^5	1	0	0	0	1	0	$H_{2/2}^1$
v^6	1	0	1	0	0	1	$H_{3/6}^2$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_3X_5	X_4X_6	X_7	R_4R_5	R_6	R_7R_8
dim	2	2	2	2	2	6
S_3S_4	2	1	1	1	-1	-1 -3

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	X_3X_5	X_4X_6	X_7	R_4R_5	R_6	R_7R_8	
	2	2	2	2	2	6	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	3	0	0	1	$H_{4/3}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	1	2	0	0	1	$H_{4/3}^1$
v^7	0	2	1	0	0	1	$H_{4/3}^1$
v^8	0	3	0	0	0	1	$H_{4/3}^1$
v^9	1	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	2	0	0	1	$H_{4/3}^1$
v^{12}	1	1	1	0	0	1	$H_{4/3}^1$
v^{13}	1	2	0	0	0	1	$H_{4/3}^1$
v^{14}	2	0	1	0	0	1	$H_{4/3}^1$
v^{15}	2	1	0	0	0	1	$H_{4/3}^1$
v^{16}	3	0	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_5	X_4X_6	X_7	M_6M_7
dim	2	2	2	4
Z_3Z_4	2	1	1	1
				-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	X_3X_5	X_4X_6	X_7	M_6M_7	
	2	2	2	4	Bipartite graph
v^1	0	0	2	1	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	2	0	1	$H_{3/2}^1$
v^4	1	0	1	1	$H_{3/2}^1$
v^5	1	1	0	1	$H_{3/2}^1$
v^6	2	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6M_7	R_4R_5	R_6	R_7R_8
	dim	4	2	2	6
T_5T_7	2	1	-1	-1	-1
T_6T_8	2	1	0	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6M_7	R_4R_5	R_6	R_7R_8	
	4	2	2	6	Bipartite graph
v^1	2	0	1	1	$H_{4/4}^4$
v^2	2	1	0	1	$H_{4/4}^4$

SG 213

Space group $P4_132$, No. 213, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	X_3X_5	X_4X_6	X_7
	dim	2	2	4	2	2	2
Δ_5	1	1	0	1	0	-2	0
Δ_6	1	0	1	1	0	0	-1
Δ_7	1	0	1	1	-2	0	0
Δ_8	1	1	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	X_3X_5	X_4X_6	X_7	
	2	2	4	2	2	2	Bipartite graph
v^1	0	0	2	1	1	2	$H_{6/8}^1$
v^2	1	1	1	1	1	2	$H_{7/8}^{11}, H_{7/8}^9$
v^3	2	2	0	1	1	2	$H_{8/8}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	M_6M_7
	dim	2	2	4	4
Σ_3	1	1	1	2	-2
Σ_4	1	1	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Γ_8	M_6M_7	
	2	2	4	4	Bipartite graph
v^1	0	0	1	1	$H_{2/4}^1$
v^2	0	2	0	1	$H_{3/4}^1$
v^3	1	1	0	1	$H_{3/4}^1$
v^4	2	0	0	1	$H_{3/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	R_4R_5	R_6	R_7R_8
	dim	2	2	4	2	2	6
Λ_4	1	1	1	1	0	-1	-2
Λ_5	1	1	1	1	0	-1	-2
Λ_6	1	0	0	2	-2	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6	Γ_7	Γ_8	R_4R_5	R_6	R_7R_8	
	2	2	4	2	2	6	Bipartite graph
v^1	0	0	1	1	1	0	$H_{3/4}^1$
v^2	0	0	2	1	0	1	$H_{4/8}^1$
v^3	0	1	0	0	1	0	$H_{2/2}^1$
v^4	0	1	1	0	0	1	$H_{3/6}^2$
v^5	1	0	0	0	1	0	$H_{2/2}^1$
v^6	1	0	1	0	0	1	$H_{3/6}^2$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	X_3X_5	X_4X_6	X_7	R_4R_5	R_6	R_7R_8	
dim	2	2	2	2	2	6	
S_3S_4	2	1	1	1	-1	-1	-3

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	X_3X_5	X_4X_6	X_7	R_4R_5	R_6	R_7R_8	
	2	2	2	2	2	6	Bipartite graph
v^1	0	0	1	0	1	0	$H_{2/1}^1$
v^2	0	0	1	1	0	0	$H_{2/1}^1$
v^3	0	0	3	0	0	1	$H_{4/3}^1$
v^4	0	1	0	0	1	0	$H_{2/1}^1$
v^5	0	1	0	1	0	0	$H_{2/1}^1$
v^6	0	1	2	0	0	1	$H_{4/3}^1$
v^7	0	2	1	0	0	1	$H_{4/3}^1$
v^8	0	3	0	0	0	1	$H_{4/3}^1$
v^9	1	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	1	0	0	$H_{2/1}^1$
v^{11}	1	0	2	0	0	1	$H_{4/3}^1$
v^{12}	1	1	1	0	0	1	$H_{4/3}^1$
v^{13}	1	2	0	0	0	1	$H_{4/3}^1$
v^{14}	2	0	1	0	0	1	$H_{4/3}^1$
v^{15}	2	1	0	0	0	1	$H_{4/3}^1$
v^{16}	3	0	0	0	0	1	$H_{4/3}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	X_3X_5	X_4X_6	X_7	M_6M_7	
dim	2	2	2	4	
Z_3Z_4	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	X_3X_5	X_4X_6	X_7	M_6M_7	
	2	2	2	4	Bipartite graph
v^1	0	0	2	1	$H_{3/2}^1$
v^2	0	1	1	1	$H_{3/2}^1$
v^3	0	2	0	1	$H_{3/2}^1$
v^4	1	0	1	1	$H_{3/2}^1$
v^5	1	1	0	1	$H_{3/2}^1$
v^6	2	0	0	1	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		M_6M_7	R_4R_5	R_6	R_7R_8
	dim	4	2	2	6
T_5T_7	2	1	0	0	-2
T_6T_8	2	1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_6M_7	R_4R_5	R_6	R_7R_8	
	4	2	2	6	Bipartite graph
v^1	2	0	1	1	$H_{4/4}^4$
v^2	2	1	0	1	$H_{4/4}^4$

SG 214

Space group $I4_132$, No. 214, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Γ_8	N_5
	dim	2	2	4	2
Σ_3	1	1	1	2	-1
Σ_4	1	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	N_5	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/4}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	H_6	H_7	H_8
	dim	2	2	4	2	2	4
Δ_5	1	1	0	1	-1	0	-1
Δ_6	1	0	1	1	-1	0	-1
Δ_7	1	0	1	1	0	-1	-1
Δ_8	1	1	0	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Γ_8	H_6	H_7	H_8	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/4}^1$
v^2	0	0	1	1	1	0	$H_{3/4}^1$
v^3	1	1	0	0	0	1	$H_{3/4}^1$
v^4	1	1	0	1	1	0	$H_{4/4}^2$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_6	Γ_7	Γ_8	P_4	P_5	P_6	P_7
	dim	2	2	4	1	1	1	3
Λ_4	1	1	1	1	0	0	-1	-1
Λ_5	1	1	1	1	0	-1	0	-1
Λ_6	1	0	0	2	-1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6	Γ_7	Γ_8	P_4	P_5	P_6	P_7	
	2	2	4	1	1	1	3	Bipartite graph
v^1	0	0	1	1	0	0	1	$H_{3/4}^2$
v^2	0	0	1	2	1	1	0	$H_{5/4}^1$
v^3	0	1	0	0	1	1	0	$H_{3/2}^1$
v^4	0	1	1	0	0	0	2	$H_{4/6}^1$
v^5	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	1	0	1	0	0	0	2	$H_{4/6}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$
A matrix

corep		H_6	H_7	H_8	N_5
	dim	2	2	4	2
G_3	1	1	1	2	-1
G_4	1	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	H_6	H_7	H_8	N_5	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/4}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_6	H_7	H_8	P_4	P_5	P_6	P_7
	dim	2	2	4	1	1	1	3
F_4	1	0	0	2	-1	0	0	-1
F_5	1	1	1	1	0	0	-1	-1
F_6	1	1	1	1	0	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	H_6	H_7	H_8	P_4	P_5	P_6	P_7	
	2	2	4	1	1	1	3	Bipartite graph
v^1	0	0	1	1	0	0	1	$H_{3/4}^2$
v^2	0	0	1	2	1	1	0	$H_{5/4}^1$
v^3	0	1	0	0	1	1	0	$H_{3/2}^1$
v^4	0	1	1	0	0	0	2	$H_{4/6}^1$
v^5	1	0	0	0	1	1	0	$H_{3/2}^1$
v^6	1	0	1	0	0	0	2	$H_{4/6}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_4	P_5	P_6	P_7	N_5
	dim	1	1	1	3	2
D_3	1	0	0	0	2	-1
D_4	1	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_4	P_5	P_6	P_7	N_5	
	1	1	1	3	2	Bipartite graph
v^1	0	0	1	1	2	$H_{4/4}^4$
v^2	0	1	0	1	2	$H_{4/4}^4$
v^3	1	0	0	1	2	$H_{4/4}^4$

SG 215

Space group $P\bar{4}3m$, No. 215, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	X_6	X_7
	dim	2	2	4	2	2
Δ_5	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	X_6	X_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6	Γ_7	Γ_8	M_6	M_7
	dim	2	2	4	2	2
Σ_3	1	1	1	2	-1	-1
Σ_4	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	M_6	M_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/4}^1$
v^2	0	0	1	1	1	$H_{3/4}^1$
v^3	0	0	1	2	0	$H_{3/4}^1$
v^4	0	1	0	0	1	$H_{2/2}^1$
v^5	0	1	0	1	0	$H_{2/2}^1$
v^6	1	0	0	0	1	$H_{2/2}^1$
v^7	1	0	0	1	0	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Γ_8	R_6	R_7	R_8
	dim	2	2	4	2	2	4
Λ_4	2	1	1	1	-1	-1	-1
Λ_5	1	0	0	1	0	0	-1
Λ_6	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_6	Γ_7	Γ_8	R_6	R_7	R_8	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/3}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_6	X_7	R_6	R_7	R_8
	dim	2	2	2	2	4
S_3	1	1	1	-1	-1	-2
S_4	1	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	X_6	X_7	R_6	R_7	R_8	
	2	2	2	2	4	Bipartite graph
v^1	0	1	0	1	0	$H_{2/2}^1$
v^2	0	1	1	0	0	$H_{2/2}^1$
v^3	0	2	0	0	1	$H_{3/4}^1$
v^4	1	0	0	1	0	$H_{2/2}^1$
v^5	1	0	1	0	0	$H_{2/2}^1$
v^6	1	1	0	0	1	$H_{3/4}^1$
v^7	2	0	0	0	1	$H_{3/4}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_6	X_7	M_6	M_7
	dim	2	2	2	2
Z_3	1	1	1	-1	-1
Z_4	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	X_6	X_7	M_6	M_7	
	2	2	2	2	Bipartite graph
v^1	0	1	0	1	$H_{2/2}^1$
v^2	0	1	1	0	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$
v^4	1	0	1	0	$H_{2/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	R_6	R_7	R_8
	dim	2	2	2	2	4
T_5	2	1	1	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	M_6	M_7	R_6	R_7	R_8	Bipartite graph
	2	2	2	2	4	
v^1	0	1	0	1	0	$H_{2/1}^1$
v^2	0	1	1	0	0	$H_{2/1}^1$
v^3	0	2	0	0	1	$H_{3/2}^1$
v^4	1	0	0	1	0	$H_{2/1}^1$
v^5	1	0	1	0	0	$H_{2/1}^1$
v^6	1	1	0	0	1	$H_{3/2}^1$
v^7	2	0	0	0	1	$H_{3/2}^1$

SG 216

Space group $F\bar{4}3m$, No. 216, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_6	Γ_7	Γ_8	X_6	X_7
	dim	2	2	4	2	2
Δ_5	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	X_6	X_7	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	L_4	L_5L_6
	dim	2	2	4	2	2
Λ_4	2	1	1	1	-1	0
Λ_5	1	0	0	1	0	-1
Λ_6	1	0	0	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	L_4	L_5L_6	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/3}^1$
v^2	0	1	0	1	0	$H_{2/1}^1$
v^3	1	0	0	1	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		X_6	X_7	W_5	W_6	W_7	W_8
	dim	2	2	1	1	1	1
Z_3	1	1	1	0	-1	0	-1
Z_4	1	1	1	-1	0	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_6	X_7	W_5	W_6	W_7	W_8	
	2	2	1	1	1	1	Bipartite graph
v^1	0	1	0	0	1	1	$H_{3/2}^1$
v^2	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	0	1	1	$H_{3/2}^1$
v^6	1	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	1	0	0	1	$H_{3/2}^1$
v^8	1	0	1	1	0	0	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep		L_4	L_5L_6	W_5	W_6	W_7	W_8
dim		2	2	1	1	1	1
Q_2	1	2	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	L_4	L_5L_6	W_5	W_6	W_7	W_8	Bipartite graph
	2	2	1	1	1	1	
v^1	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	1	0	1	0	1	$H_{3/2}^1$
v^5	0	1	0	1	1	0	$H_{3/2}^1$
v^6	0	1	0	2	0	0	$H_{3/2}^1$
v^7	0	1	1	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	1	0	$H_{3/2}^1$
v^9	0	1	1	1	0	0	$H_{3/2}^1$
v^{10}	0	1	2	0	0	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	2	$H_{3/2}^1$
v^{12}	1	0	0	0	1	1	$H_{3/2}^1$
v^{13}	1	0	0	0	2	0	$H_{3/2}^1$
v^{14}	1	0	0	1	0	1	$H_{3/2}^1$
v^{15}	1	0	0	1	1	0	$H_{3/2}^1$
v^{16}	1	0	0	2	0	0	$H_{3/2}^1$
v^{17}	1	0	1	0	0	1	$H_{3/2}^1$
v^{18}	1	0	1	0	1	0	$H_{3/2}^1$
v^{19}	1	0	1	1	0	0	$H_{3/2}^1$
v^{20}	1	0	2	0	0	0	$H_{3/2}^1$

SG 217

Space group $I\bar{4}3m$, No. 217, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	N_5
	dim	2	2	4	2
Σ_3	1	1	1	2	-1
Σ_4	1	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	Γ_6	Γ_7	Γ_8	N_5	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/4}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	H_6	H_7	H_8
	dim	2	2	4	2	2	4
Δ_5	2	1	1	2	-1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 11.

	Γ_6	Γ_7	Γ_8	H_6	H_7	H_8	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	1	0	2	0	$H_{3/2}^1$
v^3	0	0	1	1	1	0	$H_{3/2}^1$
v^4	0	0	1	2	0	0	$H_{3/2}^1$
v^5	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	2	0	0	0	1	$H_{3/2}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$
v^9	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	1	0	0	0	1	$H_{3/2}^1$
v^{11}	2	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_6	Γ_7	Γ_8	P_6	P_7	P_8
	dim	2	2	4	2	2	4
Λ_4	2	1	1	1	-1	-1	-1
Λ_5	1	0	0	1	0	0	-1
Λ_6	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	Γ_6	Γ_7	Γ_8	P_6	P_7	P_8	Bipartite graph
	2	2	4	2	2	4	
v^1	0	0	1	0	0	1	$H_{2/3}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep		H_6	H_7	H_8	N_5
	dim	2	2	4	2
G_3	1	1	1	2	-1
G_4	1	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	H_6	H_7	H_8	N_5	Bipartite graph
	2	2	4	2	
v^1	0	0	1	2	$H_{3/4}^1$
v^2	0	1	0	1	$H_{2/2}^1$
v^3	1	0	0	1	$H_{2/2}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		H_6	H_7	H_8	P_6	P_7	P_8
	dim	2	2	4	2	2	4

F_4	2	1	1	1	-1	-1	-1
F_5	1	0	0	1	0	0	-1
F_6	1	0	0	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	H_6	H_7	H_8	P_6	P_7	P_8	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	1	0	0	1	$H_{2/3}^1$
v^2	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	1	0	1	0	0	$H_{2/1}^1$
v^4	1	0	0	0	1	0	$H_{2/1}^1$
v^5	1	0	0	1	0	0	$H_{2/1}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_6	P_7	P_8	N_5
	dim	2	2	4	2
D_5	2	1	1	2	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_6	P_7	P_8	N_5	
	2	2	4	2	Bipartite graph
v^1	0	0	1	2	$H_{3/2}^1$
v^2	0	1	0	1	$H_{2/1}^1$
v^3	1	0	0	1	$H_{2/1}^1$

SG 218

Space group $P\bar{4}3n$, No. 218, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	Γ_8	$X_6 X_7$
	dim	2	2	4	4
Δ_5	2	1	1	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Γ_8	$X_6 X_7$	
	2	2	4	4	Bipartite graph
v^1	0	0	1	1	$H_{2/2}^1$
v^2	0	2	0	1	$H_{3/2}^1$
v^3	1	1	0	1	$H_{3/2}^1$
v^4	2	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6	Γ_7	Γ_8	M_6	M_7
	dim	2	2	4	2	2
Σ_3	1	1	1	2	-1	-1
Σ_4	1	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	M_6	M_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/4}^1$
v^2	0	0	1	1	1	$H_{3/4}^1$
v^3	0	0	1	2	0	$H_{3/4}^1$
v^4	0	1	0	0	1	$H_{2/2}^1$
v^5	0	1	0	1	0	$H_{2/2}^1$
v^6	1	0	0	0	1	$H_{2/2}^1$
v^7	1	0	0	1	0	$H_{2/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	R_6R_7	R_8R_8
	dim	2	2	4	4	8
Λ_4	2	1	1	1	-2	-2
Λ_5	1	0	0	1	0	-2
Λ_6	1	0	0	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Γ_8	R_6R_7	R_8R_8	
	2	2	4	4	8	Bipartite graph
v^1	0	0	2	0	1	$H_{3/6}^1$
v^2	0	2	0	1	0	$H_{3/2}^1$
v^3	1	1	0	1	0	$H_{3/2}^1$
v^4	2	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		X_6X_7	R_6R_7	R_8R_8
	dim	4	4	8
S_3S_4	2	2	-2	-4

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_6X_7	R_6R_7	R_8R_8	
	4	4	8	Bipartite graph
v^1	1	1	0	$H_{2/2}^1$
v^2	2	0	1	$H_{3/4}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		X_6X_7	M_6	M_7
	dim	4	2	2
Z_3	1	2	-1	-1
Z_4	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	$X_6 X_7$	M_6	M_7	
	4	2	2	Bipartite graph
v^1	1	0	2	$H_{3/4}^1$
v^2	1	1	1	$H_{3/4}^1$
v^3	1	2	0	$H_{3/4}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T : $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6	M_7	$R_6 R_7$	$R_8 R_8$
	dim	2	2	4	8
T_5	2	1	1	-2	-4

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	M_6	M_7	$R_6 R_7$	$R_8 R_8$	
	2	2	4	8	Bipartite graph
v^1	0	2	1	0	$H_{3/2}^1$
v^2	0	4	0	1	$H_{5/4}^1$
v^3	1	1	1	0	$H_{3/2}^1$
v^4	1	3	0	1	$H_{5/4}^1$
v^5	2	0	1	0	$H_{3/2}^1$
v^6	2	2	0	1	$H_{5/4}^1$
v^7	3	1	0	1	$H_{5/4}^1$
v^8	4	0	0	1	$H_{5/4}^1$

SG 219

Space group $F\bar{4}3c$, No. 219, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_6	Γ_7	Γ_8	X_6	X_7
	dim	2	2	4	2	2
Δ_5	2	1	1	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	X_6	X_7	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_6	Γ_7	Γ_8	L_4L_4	L_5L_5	L_6L_6
dim	2	2	4	2	2	4
Λ_4	2	1	1	1	0	0
Λ_5	1	0	0	1	0	-2
Λ_6	1	0	0	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Γ_8	L_4L_4	L_5L_5	L_6L_6	
	2	2	4	2	2	4	Bipartite graph
v^1	0	0	2	1	1	1	$H_{5/6}^4$
v^2	0	2	0	0	0	1	$H_{3/2}^1$
v^3	1	1	0	0	0	1	$H_{3/2}^1$
v^4	2	0	0	0	0	1	$H_{3/2}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_6	X_7	W_5W_7	W_6W_8
dim	2	2	2	2
Z_3	1	1	0	-2
Z_4	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_6	X_7	W_5W_7	W_6W_8	
	2	2	2	2	Bipartite graph
v^1	0	2	1	1	$H_{4/4}^2$
v^2	1	1	1	1	$H_{4/4}^2$
v^3	2	0	1	1	$H_{4/4}^2$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep	L_4L_4	L_5L_5	L_6L_6	W_5W_7	W_6W_8
dim	2	2	4	2	2
Q_2Q_2	2	1	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	L_4L_4	L_5L_5	L_6L_6	W_5W_7	W_6W_8	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

SG 220

Space group $I\bar{4}3d$, No. 220, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ: $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep	Γ_6	Γ_7	Γ_8	N_2N_3	N_4N_5
dim	2	2	4	2	2
Σ_3	1	1	2	0	-2
Σ_4	1	1	2	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	Γ_6	Γ_7	Γ_8	N_2N_3	N_4N_5	
	2	2	4	2	2	Bipartite graph
v^1	0	0	1	1	1	$H_{3/4}^1$
v^2	0	2	0	1	1	$H_{4/4}^2$
v^3	1	1	0	1	1	$H_{4/4}^2$
v^4	2	0	0	1	1	$H_{4/4}^2$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_6	Γ_7	Γ_8	H_6H_7	H_8H_8
dim	2	2	4	4	8
Δ_5	2	1	1	2	-2 -4

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_6	Γ_7	Γ_8	H_6H_7	H_8H_8	
	2	2	4	4	8	Bipartite graph
v^1	0	0	1	1	0	$H_{2/2}^1$
v^2	0	0	2	0	1	$H_{3/4}^1$
v^3	0	2	0	1	0	$H_{3/2}^1$
v^4	0	2	1	0	1	$H_{4/4}^1$
v^5	0	4	0	0	1	$H_{5/4}^1$
v^6	1	1	0	1	0	$H_{3/2}^1$
v^7	1	1	1	0	1	$H_{4/4}^1$
v^8	1	3	0	0	1	$H_{5/4}^1$
v^9	2	0	0	1	0	$H_{3/2}^1$
v^{10}	2	0	1	0	1	$H_{4/4}^1$
v^{11}	2	2	0	0	1	$H_{5/4}^1$
v^{12}	3	1	0	0	1	$H_{5/4}^1$
v^{13}	4	0	0	0	1	$H_{5/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	Γ_6	Γ_7	Γ_8	P_4	P_5	P_6	P_7	P_8
dim	2	2	4	1	1	2	3	3

Λ_4	2	1	1	1	0	0	-1	-1	-1
Λ_5	1	0	0	1	0	-1	0	-1	0
Λ_6	1	0	0	1	-1	0	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	Γ_6	Γ_7	Γ_8	P_4	P_5	P_6	P_7	P_8	
	2	2	4	1	1	2	3	3	Bipartite graph
v^1	0	0	1	0	1	0	0	1	$H_{3/3}^1$
v^2	0	0	1	1	0	0	1	0	$H_{3/3}^1$
v^3	0	0	1	1	1	1	0	0	$H_{4/3}^1$
v^4	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^5	0	1	1	0	0	0	1	1	$H_{4/4}^4$
v^6	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^7	1	0	1	0	0	0	1	1	$H_{4/4}^4$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep	H_6H_7	H_8H_8	N_2N_3	N_4N_5
dim	4	8	2	2
G_3G_4	2	4	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	H_6H_7	H_8H_8	N_2N_3	N_4N_5	
	4	8	2	2	Bipartite graph
v^1	0	1	0	4	$H_{5/4}^1$
v^2	0	1	1	3	$H_{5/4}^1$
v^3	0	1	2	2	$H_{5/4}^1$
v^4	0	1	3	1	$H_{5/4}^1$
v^5	0	1	4	0	$H_{5/4}^1$
v^6	1	0	0	2	$H_{3/2}^1$
v^7	1	0	1	1	$H_{3/2}^1$
v^8	1	0	2	0	$H_{3/2}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		H_6H_7	H_8H_8	P_4	P_5	P_6	P_7	P_8
	dim	4	8	1	1	2	3	3
F_4	1	0	2	0	-1	0	-1	0
F_5	1	0	2	-1	0	0	0	-1
F_6	2	2	2	0	0	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	H_6H_7	H_8H_8	P_4	P_5	P_6	P_7	P_8	Bipartite graph
	4	8	1	1	2	3	3	
v^1	0	1	0	2	0	0	2	$H_{5/6}^1$
v^2	0	1	1	1	0	1	1	$H_{5/6}^1$
v^3	0	1	1	2	1	0	1	$H_{6/6}^1$
v^4	0	1	2	0	0	2	0	$H_{5/6}^1$
v^5	0	1	2	1	1	1	0	$H_{6/6}^1$
v^6	0	1	2	2	2	0	0	$H_{7/6}^1$
v^7	1	0	0	0	2	0	0	$H_{3/2}^1$
v^8	1	1	0	0	0	2	2	$H_{6/8}^2$
v^9	1	1	0	1	1	1	2	$H_{7/8}^2, H_{7/8}^1$
v^{10}	1	1	1	0	1	2	1	$H_{7/8}^2, H_{7/8}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D : $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$

A matrix

corep		P_4	P_5	P_6	P_7	P_8	N_2N_3	N_4N_5
	dim	1	1	2	3	3	2	2
D_2	1	0	1	1	1	0	0	-1
D_3	1	1	0	1	0	1	-1	0
D_4	1	0	0	0	1	1	-1	0
D_5	1	0	0	0	1	1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	P_4	P_5	P_6	P_7	P_8	N_2N_3	N_4N_5	Bipartite graph
	1	1	2	3	3	2	2	
v^1	0	0	1	1	1	2	2	$H_{7/8}^{10}, H_{7/8}^3$
v^2	0	1	0	0	1	1	1	$H_{4/4}^4$
v^3	1	0	0	1	0	1	1	$H_{4/4}^4$

SG 221

Space group $Pm\bar{3}m$, No. 221, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_6^+	X_6^-	X_7^+	X_7^-
	dim	2	2	2	2	4	4	2	2	2	2
Δ_6	2	1	1	0	0	1	1	-1	-1	0	0
Δ_7	2	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_6^+	X_6^-	X_7^+	X_7^-	Bipartite graph
	2	2	2	2	4	4	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_6^+	M_6^-	M_7^+	M_7^-
	dim	2	2	2	2	4	4	2	2	2	2
Σ_5	2	1	1	1	1	2	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 36.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_6^+	M_6^-	M_7^+	M_7^-	Bipartite graph
	2	2	2	2	4	4	2	2	2	2	
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^{13}	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^{14}	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^{15}	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^{16}	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^{18}	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^{19}	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^{20}	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{23}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{24}	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{26}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{31}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{32}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{33}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{34}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{35}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{36}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0)$, $R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	
dim	2	2	2	2	4	4	2	2	2	2	4	4	
Λ_4	2	1	1	1	1	1	-1	-1	-1	-1	-1	-1	
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	Bipartite graph
	2	2	2	2	4	4	2	2	2	2	4	4	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	X_6^+	X_6^-	X_7^+	X_7^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	
dim	2	2	2	2	2	2	2	2	4	4	
S_5	2	1	1	1	1	-1	-1	-1	-1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 36.

	X_6^+	X_6^-	X_7^+	X_7^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	Bipartite graph
	2	2	2	2	2	2	2	2	4	4	
v^1	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	0	0	$H_{2/1}^1$
v^5	0	0	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^6	0	0	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^7	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{12}	0	0	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	0	0	2	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	0	0	2	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{23}	0	2	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{24}	0	2	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{25}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{26}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{27}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{28}	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{29}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{30}	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{31}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{32}	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{33}	1	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{34}	1	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{35}	2	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{36}	2	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z : $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	X_6^+	X_6^-	X_7^+	X_7^-	M_6^+	M_6^-	M_7^+	M_7^-	
dim	2	2	2	2	2	2	2	2	
Z_5	2	1	1	1	1	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	X_6^+	X_6^-	X_7^+	X_7^-	M_6^+	M_6^-	M_7^+	M_7^-	Bipartite graph
	2	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	0	1	0	0	$H_{2/1}^1$
v^4	0	0	0	1	1	0	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^8	0	0	1	0	1	0	0	0	$H_{2/1}^1$
v^9	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	1	0	0	0	$H_{2/1}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep	M_6^+	M_6^-	M_7^+	M_7^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	
dim	2	2	2	2	2	2	2	2	4	4	
T_6	2	1	1	0	0	-1	-1	0	0	-1	-1
T_7	2	0	0	1	1	0	0	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

M_6^+	M_6^-	M_7^+	M_7^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	Bipartite graph
2	2	2	2	2	2	2	2	4	4	

v^1	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^2	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^3	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^6	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^7	0	1	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^8	0	1	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^9	0	1	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	0	1	1	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{13}	1	0	0	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{14}	1	0	0	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{15}	1	0	1	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{16}	1	0	1	0	0	0	0	0	1	0	$H_{3/2}^1$

SG 222

Space group $Pn\bar{3}n$, No. 222, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0)$, $X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5X_6	X_7X_8
	dim	2	2	2	2	4	4	4	4
Δ_6	2	1	1	0	0	1	1	-2	0
Δ_7	2	0	0	1	1	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5X_6	X_7X_8	
	2	2	2	2	4	4	4	4	Bipartite graph
v^1	0	0	0	0	0	2	1	1	$H_{4/4}^2$
v^2	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	2	0	1	1	$H_{4/4}^2$
v^4	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^7	0	1	0	1	0	1	1	1	$H_{5/4}^3$
v^8	0	1	0	1	1	0	1	1	$H_{5/4}^3$

v^9	0	1	1	0	0	1	1	1	$H_{5/4}^3$
v^{10}	0	1	1	0	1	0	1	1	$H_{5/4}^3$
v^{11}	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{12}	1	0	0	1	0	1	1	1	$H_{5/4}^3$
v^{13}	1	0	0	1	1	0	1	1	$H_{5/4}^3$
v^{14}	1	0	1	0	0	1	1	1	$H_{5/4}^3$
v^{15}	1	0	1	0	1	0	1	1	$H_{5/4}^3$
v^{16}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{17}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_5	
dim	2	2	2	2	4	4	4	
Σ_5	2	1	1	1	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_5	Bipartite graph
	2	2	2	2	4	4	4	
v^1	0	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	1	$H_{2/2}^1$
v^3	0	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	0	1	1	0	0	1	$H_{3/2}^1$
v^5	0	0	2	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	1	$H_{3/2}^1$
v^7	0	1	1	0	0	0	1	$H_{3/2}^1$
v^8	0	2	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	2	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_5	R_6R_7
	dim	2	2	2	2	4	4	4	8
Λ_4	2	1	1	1	1	1	1	-2	-2
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_5	R_6R_7	Bipartite graph
	2	2	2	2	4	4	4	8	
v^1	0	0	0	0	0	2	0	1	$H_{3/4}^1$
v^2	0	0	0	0	1	1	0	1	$H_{3/4}^1$
v^3	0	0	0	0	2	0	0	1	$H_{3/4}^1$
v^4	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^5	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^9	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5X_6	X_7X_8	R_5	R_6R_7
	dim	4	4	4	8
S_2S_5	2	1	1	-1	-2
S_3S_4	2	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 5.

	X_5X_6	X_7X_8	R_5	R_6R_7	Bipartite graph
	4	4	4	8	
v^1	0	1	1	0	$H_{2/2}^1$
v^2	0	2	0	1	$H_{3/4}^1$
v^3	1	0	1	0	$H_{2/2}^1$
v^4	1	1	0	1	$H_{3/4}^1$
v^5	2	0	0	1	$H_{3/4}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep	$X_5 X_6$	$X_7 X_8$	M_5
dim	4	4	4
$Z_2 Z_5$	2	1	1
$Z_3 Z_4$	2	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	$X_5 X_6$	$X_7 X_8$	M_5	
	4	4	4	Bipartite graph
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	M_5	R_5	$R_6 R_7$
dim	4	4	8
T_6	2	1	-1
T_7	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	M_5	R_5	$R_6 R_7$	
	4	4	8	Bipartite graph
v^1	1	1	0	$H_{2/2}^1$
v^2	2	0	1	$H_{3/4}^1$

SG 223

Space group $Pm\bar{3}n$, No. 223, with SOC

 Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5
	dim	2	2	2	2	4	4	4
Δ_6	2	1	1	0	0	1	1	-1
Δ_7	2	0	0	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5	
	2	2	2	2	4	4	4	Bipartite graph
v^1	0	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	1	$H_{2/2}^1$
v^3	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_6^+	M_6^-	M_7^+	M_7^-
	dim	2	2	2	2	4	4	2	2	2	2
Σ_5	2	1	1	1	1	2	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 36.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_6^+	M_6^-	M_7^+	M_7^-	
	2	2	2	2	4	4	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	0	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	0	1	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	0	0	2	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^5	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	0	2	0	0	$H_{3/2}^1$
v^7	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	0	0	1	1	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	0	1	2	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	0	0	0	2	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	0	0	1	1	$H_{3/2}^1$
v^{13}	0	0	0	0	1	0	0	0	2	0	$H_{3/2}^1$
v^{14}	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$

v^{15}	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^{16}	0	0	0	0	1	0	0	2	0	0	$H_{3/2}^1$
v^{17}	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^{18}	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^{19}	0	0	0	0	1	0	1	1	0	0	$H_{3/2}^1$
v^{20}	0	0	0	0	1	0	2	0	0	0	$H_{3/2}^1$
v^{21}	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{22}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{23}	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{24}	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{25}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{26}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{27}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{28}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{29}	0	1	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{30}	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{31}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{32}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{33}	1	0	0	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{34}	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{35}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{36}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_5	R_6R_7
	dim	2	2	2	2	4	4	4	8
Λ_4	2	1	1	1	1	1	1	-2	-2
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	0	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_5	R_6R_7	Bipartite graph
	2	2	2	2	4	4	4	8	
v^1	0	0	0	0	0	2	0	1	$H_{3/4}^1$
v^2	0	0	0	0	1	1	0	1	$H_{3/4}^1$
v^3	0	0	0	0	2	0	0	1	$H_{3/4}^1$
v^4	0	0	0	2	0	0	1	0	$H_{3/2}^1$
v^5	0	0	1	1	0	0	1	0	$H_{3/2}^1$
v^6	0	0	2	0	0	0	1	0	$H_{3/2}^1$

v^7	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^8	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^9	0	2	0	0	0	0	1	0	$H_{3/2}^1$
v^{10}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{11}	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	1	0	$H_{3/2}^1$
v^{13}	2	0	0	0	0	0	1	0	$H_{3/2}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S: $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	R_5	R_6R_7
	dim	4	4	8
S_2S_3	2	1	-1	-2
S_4S_5	2	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	R_5	R_6R_7	
	4	4	8	Bipartite graph
v^1	1	1	0	$H_{2/2}^1$
v^2	2	0	1	$H_{3/4}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	M_6^+	M_6^-	M_7^+	M_7^-
	dim	4	2	2	2	2
Z_5	2	2	-1	-1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	X_5	M_6^+	M_6^-	M_7^+	M_7^-	
	4	2	2	2	2	Bipartite graph
v^1	1	0	0	0	2	$H_{3/2}^1$
v^2	1	0	0	1	1	$H_{3/2}^1$
v^3	1	0	0	2	0	$H_{3/2}^1$
v^4	1	0	1	0	1	$H_{3/2}^1$
v^5	1	0	1	1	0	$H_{3/2}^1$

v^6	1	0	2	0	0	$H_{3/2}^1$
v^7	1	1	0	0	1	$H_{3/2}^1$
v^8	1	1	0	1	0	$H_{3/2}^1$
v^9	1	1	1	0	0	$H_{3/2}^1$
v^{10}	1	2	0	0	0	$H_{3/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_6^+	M_6^-	M_7^+	M_7^-	R_5	R_6R_7
	dim	2	2	2	2	4	8
T_6	2	0	0	1	1	-1	-2
T_7	2	1	1	0	0	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	M_6^+	M_6^-	M_7^+	M_7^-	R_5	R_6R_7	
	2	2	2	2	4	8	Bipartite graph
v^1	0	1	0	1	1	0	$H_{3/2}^1$
v^2	0	1	1	0	1	0	$H_{3/2}^1$
v^3	0	2	0	2	0	1	$H_{5/4}^1$
v^4	0	2	1	1	0	1	$H_{5/4}^1$
v^5	0	2	2	0	0	1	$H_{5/4}^1$
v^6	1	0	0	1	1	0	$H_{3/2}^1$
v^7	1	0	1	0	1	0	$H_{3/2}^1$
v^8	1	1	0	2	0	1	$H_{5/4}^1$
v^9	1	1	1	1	0	1	$H_{5/4}^1$
v^{10}	1	1	2	0	0	1	$H_{5/4}^1$
v^{11}	2	0	0	2	0	1	$H_{5/4}^1$
v^{12}	2	0	1	1	0	1	$H_{5/4}^1$
v^{13}	2	0	2	0	0	1	$H_{5/4}^1$

SG 224

Space group $Pn\bar{3}m$, No. 224, with SOC

Δ

High symmetry line $\Delta(0, u, 0)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(0, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5
	dim	2	2	2	2	4	4	4
Δ_6	2	1	1	0	0	1	1	-1
Δ_7	2	0	0	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5	Bipartite graph
	2	2	2	2	4	4	4	
v^1	0	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	1	$H_{2/2}^1$
v^3	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	1	$H_{3/2}^1$

Σ

High symmetry line $\Sigma(u, u, 0)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_5
	dim	2	2	2	2	4	4	4
Σ_5	2	1	1	1	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	M_5	Bipartite graph
	2	2	2	2	4	4	4	
v^1	0	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	1	$H_{2/2}^1$
v^3	0	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	0	1	1	0	0	1	$H_{3/2}^1$
v^5	0	0	2	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	1	$H_{3/2}^1$
v^7	0	1	1	0	0	0	1	$H_{3/2}^1$
v^8	0	2	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	2	0	0	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-
dim		2	2	2	2	4	4	2	2	2	2	4	4
Λ_4	2	1	1	1	1	1	1	-1	-1	-1	-1	-1	-1
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	0	0	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 20.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	Bipartite graph
	2	2	2	2	4	4	2	2	2	2	4	4	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^4	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^5	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^6	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^7	0	0	0	1	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^8	0	0	0	1	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{16}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{17}	1	0	0	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{18}	1	0	0	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{19}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{20}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$

S

High symmetry line $S(u, \frac{1}{2}, u)$, high symmetry points at both ends of S : $X(0, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		X_5	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-
dim		4	2	2	2	2	4	4
S_5	2	2	-1	-1	-1	-1	-2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	X_5	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	
	4	2	2	2	2	4	4	Bipartite graph
v^1	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	1	0	0	0	2	0	0	$H_{3/2}^1$
v^4	1	0	0	1	1	0	0	$H_{3/2}^1$
v^5	1	0	0	2	0	0	0	$H_{3/2}^1$
v^6	1	0	1	0	1	0	0	$H_{3/2}^1$
v^7	1	0	1	1	0	0	0	$H_{3/2}^1$
v^8	1	0	2	0	0	0	0	$H_{3/2}^1$
v^9	1	1	0	0	1	0	0	$H_{3/2}^1$
v^{10}	1	1	0	1	0	0	0	$H_{3/2}^1$
v^{11}	1	1	1	0	0	0	0	$H_{3/2}^1$
v^{12}	1	2	0	0	0	0	0	$H_{3/2}^1$

Z

High symmetry line $Z(u, \frac{1}{2}, 0)$, high symmetry points at both ends of Z: $X(0, \frac{1}{2}, 0), M(\frac{1}{2}, \frac{1}{2}, 0)$
A matrix

corep		X_5	M_5
	dim	4	4
Z_2Z_5	2	1	-1
Z_3Z_4	2	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^2$ and the corresponding bipartite graph, number of Hilbert basis: 1.

	X_5	M_5	
	4	4	Bipartite graph
v^1	1	1	$H_{2/2}^1$

T

High symmetry line $T(\frac{1}{2}, \frac{1}{2}, u)$, high symmetry points at both ends of T: $M(\frac{1}{2}, \frac{1}{2}, 0), R(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep		M_5	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-
	dim	4	2	2	2	2	4	4
T_6	2	1	0	0	-1	-1	-1	-1
T_7	2	1	-1	-1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	M_5	R_6^+	R_6^-	R_7^+	R_7^-	R_8^+	R_8^-	
	4	2	2	2	2	4	4	Bipartite graph
v^1	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	1	0	1	0	1	0	0	$H_{3/2}^1$
v^4	1	0	1	1	0	0	0	$H_{3/2}^1$
v^5	1	1	0	0	1	0	0	$H_{3/2}^1$
v^6	1	1	0	1	0	0	0	$H_{3/2}^1$

SG 225

Space group $Fm\bar{3}m$, No. 225, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_6^+	X_6^-	X_7^+	X_7^-
	dim	2	2	2	2	4	4	2	2	2	2
Δ_6	2	1	1	0	0	1	1	-1	-1	0	0
Δ_7	2	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_6^+	X_6^-	X_7^+	X_7^-	
	2	2	2	2	4	4	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4^+	L_4^-	L_5^+	L_6^+	L_5^-	L_6^-
dim	2	2	2	2	4	4	2	2	2	2	2	2
Λ_4	2	1	1	1	1	1	-1	-1	0	0		
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	0	0	-1	-1	

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4^+	L_4^-	L_5^+	L_6^+	L_5^-	L_6^-	Bipartite graph
	2	2	2	2	4	4	2	2	2	2			
v^1	0	0	0	0	0	1	0	1	0	1			$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0			$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1			$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0			$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1			$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0			$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1			$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0			$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	1	0	0			$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	1	0	0	0			$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0			$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0			$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0			$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0			$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0			$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0			$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep	X_6^+	X_6^-	X_7^+	X_7^-	W_6	W_7
dim	2	2	2	2	2	2
Z_5	2	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	X_6^+	X_6^-	X_7^+	X_7^-	W_6	W_7	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	L_4^+	L_4^-	$L_5^+ L_6^+$	$L_5^- L_6^-$	W_6	W_7
dim	2	2	2	2	2	2
$Q_3 Q_4$	2	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^6$ and the corresponding bipartite graph, number of Hilbert basis: 8.

	L_4^+	L_4^-	$L_5^+ L_6^+$	$L_5^- L_6^-$	W_6	W_7	
	2	2	2	2	2	2	Bipartite graph
v^1	0	0	0	1	0	1	$H_{2/1}^1$
v^2	0	0	0	1	1	0	$H_{2/1}^1$
v^3	0	0	1	0	0	1	$H_{2/1}^1$
v^4	0	0	1	0	1	0	$H_{2/1}^1$
v^5	0	1	0	0	0	1	$H_{2/1}^1$
v^6	0	1	0	0	1	0	$H_{2/1}^1$
v^7	1	0	0	0	0	1	$H_{2/1}^1$
v^8	1	0	0	0	1	0	$H_{2/1}^1$

SG 226

Space group $Fm\bar{3}c$, No. 226, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_6^+	X_6^-	X_7^+	X_7^-
	dim	2	2	2	2	4	4	2	2	2	2
Δ_6	2	1	1	0	0	1	1	-1	-1	0	0
Δ_7	2	0	0	1	1	1	1	0	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_6^+	X_6^-	X_7^+	X_7^-	
	2	2	2	2	4	4	2	2	2	2	Bipartite graph
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4L_4	L_5L_6
	dim	2	2	2	2	4	4	4	4
Λ_4	2	1	1	1	1	1	1	0	-2
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4L_4	L_5L_6	
	2	2	2	2	4	4	4	4	Bipartite graph
v^1	0	0	0	0	0	2	1	1	$H_{4/4}^2$
v^2	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	2	0	1	1	$H_{4/4}^2$

v^4	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{13}	2	0	0	0	0	0	0	1	$H_{3/2}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z: $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	X_6^+	X_6^-	X_7^+	X_7^-	W_6W_7
dim	2	2	2	2	4
Z_5	2	1	1	1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	X_6^+	X_6^-	X_7^+	X_7^-	W_6W_7	Bipartite graph
	2	2	2	2	4	
v^1	0	0	0	2	1	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	2	0	1	$H_{3/2}^1$
v^4	0	1	0	1	1	$H_{3/2}^1$
v^5	0	1	1	0	1	$H_{3/2}^1$
v^6	0	2	0	0	1	$H_{3/2}^1$
v^7	1	0	0	1	1	$H_{3/2}^1$
v^8	1	0	1	0	1	$H_{3/2}^1$
v^9	1	1	0	0	1	$H_{3/2}^1$
v^{10}	2	0	0	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep	L_4L_4	L_5L_6	W_6W_7
dim	4	4	4

Q_3Q_3	2	1	1	-1
Q_4Q_4	2	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	L_4L_4	L_5L_6	W_6W_7	Bipartite graph
	4	4	4	
v^1	0	1	1	$H_{2/2}^1$
v^2	1	0	1	$H_{2/2}^1$

SG 227

Space group $Fd\bar{3}m$, No. 227, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5
	dim	2	2	2	2	4	4	4
Δ_6	2	1	1	0	0	1	1	-1
Δ_7	2	0	0	1	1	1	1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5	Bipartite graph
	2	2	2	2	4	4	4	
v^1	0	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	1	$H_{2/2}^1$
v^3	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4^+	L_4^-	$L_5^+L_6^+$	$L_5^-L_6^-$
dim	2	2	2	2	4	4	2	2	2	2
Λ_4	2	1	1	1	1	1	-1	-1	0	0
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	0	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^{10}$ and the corresponding bipartite graph, number of Hilbert basis: 16.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4^+	L_4^-	$L_5^+L_6^+$	$L_5^-L_6^-$	Bipartite graph
	2	2	2	2	4	4	2	2	2	2	
v^1	0	0	0	0	0	1	0	1	0	1	$H_{3/2}^1$
v^2	0	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^3	0	0	0	0	0	1	1	0	0	1	$H_{3/2}^1$
v^4	0	0	0	0	0	1	1	0	1	0	$H_{3/2}^1$
v^5	0	0	0	0	1	0	0	1	0	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	0	1	1	0	$H_{3/2}^1$
v^7	0	0	0	0	1	0	1	0	0	1	$H_{3/2}^1$
v^8	0	0	0	0	1	0	1	0	1	0	$H_{3/2}^1$
v^9	0	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^{10}	0	0	0	1	0	0	1	0	0	0	$H_{2/1}^1$
v^{11}	0	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{12}	0	0	1	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z: $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep	X_5	W_3W_4	W_5W_6	W_7	
dim	4	2	2	2	
Z_2Z_4	2	1	-1	-1	0
Z_3Z_5	2	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	X_5	W_3W_4	W_5W_6	W_7	Bipartite graph
	4	2	2	2	
v^1	1	0	1	1	$H_{3/2}^1$
v^2	1	1	0	1	$H_{3/2}^1$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$
A matrix

corep	L_4^+	L_4^-	$L_5^+L_6^+$	$L_5^-L_6^-$	W_3W_4	W_5W_6	W_7
dim	2	2	2	2	2	2	2
Q_3Q_4	2	1	1	1	1	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	L_4^+	L_4^-	$L_5^+L_6^+$	$L_5^-L_6^-$	W_3W_4	W_5W_6	W_7	Bipartite graph
	2	2	2	2	2	2	2	
v^1	0	0	0	1	0	0	1	$H_{2/1}^1$
v^2	0	0	0	1	0	1	0	$H_{2/1}^1$
v^3	0	0	0	1	1	0	0	$H_{2/1}^1$
v^4	0	0	1	0	0	0	1	$H_{2/1}^1$
v^5	0	0	1	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	1	$H_{2/1}^1$
v^8	0	1	0	0	0	1	0	$H_{2/1}^1$
v^9	0	1	0	0	1	0	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{11}	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{12}	1	0	0	0	1	0	0	$H_{2/1}^1$

SG 228

Space group $Fd\bar{3}c$, No. 228, with SOC

Δ

High symmetry line $\Delta(u, 0, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), X(\frac{1}{2}, 0, \frac{1}{2})$
A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5
dim	2	2	2	2	4	4	4
Δ_6	2	1	1	0	0	1	1
Δ_7	2	0	0	1	1	1	1

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 6.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	X_5	
	2	2	2	2	4	4	4	Bipartite graph
v^1	0	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	1	$H_{2/2}^1$
v^3	0	1	0	1	0	0	1	$H_{3/2}^1$
v^4	0	1	1	0	0	0	1	$H_{3/2}^1$
v^5	1	0	0	1	0	0	1	$H_{3/2}^1$
v^6	1	0	1	0	0	0	1	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4L_4	L_5L_6
	dim	2	2	2	2	4	4	4	4
Λ_4	2	1	1	1	1	1	1	0	-2
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	-2	0

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 13.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	L_4L_4	L_5L_6	
	2	2	2	2	4	4	4	4	Bipartite graph
v^1	0	0	0	0	0	2	1	1	$H_{4/4}^2$
v^2	0	0	0	0	1	1	1	1	$H_{4/4}^2$
v^3	0	0	0	0	2	0	1	1	$H_{4/4}^2$
v^4	0	0	0	2	0	0	0	1	$H_{3/2}^1$
v^5	0	0	1	1	0	0	0	1	$H_{3/2}^1$
v^6	0	0	2	0	0	0	0	1	$H_{3/2}^1$
v^7	0	1	0	1	0	0	0	1	$H_{3/2}^1$
v^8	0	1	1	0	0	0	0	1	$H_{3/2}^1$
v^9	0	2	0	0	0	0	0	1	$H_{3/2}^1$
v^{10}	1	0	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	1	0	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	1	1	0	0	0	0	0	1	$H_{3/2}^1$
v^{13}	2	0	0	0	0	0	0	1	$H_{3/2}^1$

Z

High symmetry line $Z(\frac{1}{2}, u, u + \frac{1}{2})$, high symmetry points at both ends of Z : $X(\frac{1}{2}, 0, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		X_5	W_3W_6	W_4W_5	W_7W_7
	dim	4	2	2	4
Z_2Z_4	2	1	0	0	-2
Z_3Z_5	2	1	-1	-1	0

$\ker M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 3.

	X_5	W_3W_6	W_4W_5	W_7W_7	
	4	2	2	4	Bipartite graph
v^1	2	0	2	1	$H_{5/4}^3$
v^2	2	1	1	1	$H_{5/4}^3$
v^3	2	2	0	1	$H_{5/4}^3$

Q

High symmetry line $Q(\frac{1}{2}, \frac{1}{2} - u, u + \frac{1}{2})$, high symmetry points at both ends of Q: $L(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}), W(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})$

A matrix

corep		L_4L_4	L_5L_6	W_3W_6	W_4W_5	W_7W_7
	dim	4	4	2	2	4
Q_3Q_3	2	1	1	0	-1	-1
Q_4Q_4	2	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	L_4L_4	L_5L_6	W_3W_6	W_4W_5	W_7W_7	
	4	4	2	2	4	Bipartite graph
v^1	0	1	0	0	1	$H_{2/2}^1$
v^2	0	1	1	1	0	$H_{3/2}^1$
v^3	1	0	0	0	1	$H_{2/2}^1$
v^4	1	0	1	1	0	$H_{3/2}^1$

SG 229

Space group $Im\bar{3}m$, No. 229, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	N_5^+	N_5^-
dim	2	2	2	2	4	4	2	2
Σ_5	2	1	1	1	1	2	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	N_5^+	N_5^-	Bipartite graph
	2	2	2	2	4	4	2	2	
v^1	0	0	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$

A matrix

corep	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	H_6^+	H_6^-	H_7^+	H_7^-	H_8^+	H_8^-
dim	2	2	2	2	4	4	2	2	2	2	4	4
Δ_6	2	1	1	0	0	1	1	-1	-1	0	0	-1 -1
Δ_7	2	0	0	1	1	1	1	0	0	-1	-1	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^{12}$ and the corresponding bipartite graph, number of Hilbert basis: 28.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	H_6^+	H_6^-	H_7^+	H_7^-	H_8^+	H_8^-	Bipartite graph
	2	2	2	2	4	4	2	2	2	2	4	4	
v^1	0	0	0	0	0	1	0	0	0	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	0	1	0	0	0	0	1	0	$H_{2/2}^1$
v^3	0	0	0	0	0	1	0	1	0	1	0	0	$H_{3/2}^1$
v^4	0	0	0	0	0	1	0	1	1	0	0	0	$H_{3/2}^1$
v^5	0	0	0	0	0	1	1	0	0	1	0	0	$H_{3/2}^1$
v^6	0	0	0	0	0	1	1	0	1	0	0	0	$H_{3/2}^1$

v^7	0	0	0	0	1	0	0	0	0	0	0	1	$H_{2/2}^1$
v^8	0	0	0	0	1	0	0	0	0	0	1	0	$H_{2/2}^1$
v^9	0	0	0	0	1	0	0	1	0	1	0	0	$H_{3/2}^1$
v^{10}	0	0	0	0	1	0	0	1	1	0	0	0	$H_{3/2}^1$
v^{11}	0	0	0	0	1	0	1	0	0	1	0	0	$H_{3/2}^1$
v^{12}	0	0	0	0	1	0	1	0	1	0	0	0	$H_{3/2}^1$
v^{13}	0	0	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{14}	0	0	0	1	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{15}	0	0	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^{16}	0	0	1	0	0	0	0	0	1	0	0	0	$H_{2/1}^1$
v^{17}	0	1	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{18}	0	1	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{19}	0	1	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{20}	0	1	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{21}	0	1	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{22}	0	1	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{23}	1	0	0	0	0	0	0	1	0	0	0	0	$H_{2/1}^1$
v^{24}	1	0	0	0	0	0	1	0	0	0	0	0	$H_{2/1}^1$
v^{25}	1	0	0	1	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{26}	1	0	0	1	0	0	0	0	0	0	1	0	$H_{3/2}^1$
v^{27}	1	0	1	0	0	0	0	0	0	0	0	1	$H_{3/2}^1$
v^{28}	1	0	1	0	0	0	0	0	0	0	1	0	$H_{3/2}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	P_6	P_7	P_8
	dim	2	2	2	2	4	4	2	2	4
Λ_4	2	1	1	1	1	1	1	-1	-1	-1
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	P_6	P_7	P_8	Bipartite graph
	2	2	2	2	4	4	2	2	4	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^3	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$

v^7	0	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	0	0	1	0	0	$H_{2/1}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$
A matrix

corep		H_6^+	H_6^-	H_7^+	H_7^-	H_8^+	H_8^-	N_5^+	N_5^-
dim		2	2	2	2	4	4	2	2
G_5	2	1	1	1	1	2	2	-1	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 14.

	H_6^+	H_6^-	H_7^+	H_7^-	H_8^+	H_8^-	N_5^+	N_5^-	Bipartite graph
	2	2	2	2	4	4	2	2	
v^1	0	0	0	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	0	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	0	0	0	1	2	0	$H_{3/2}^1$
v^4	0	0	0	0	1	0	0	2	$H_{3/2}^1$
v^5	0	0	0	0	1	0	1	1	$H_{3/2}^1$
v^6	0	0	0	0	1	0	2	0	$H_{3/2}^1$
v^7	0	0	0	1	0	0	0	1	$H_{2/1}^1$
v^8	0	0	0	1	0	0	1	0	$H_{2/1}^1$
v^9	0	0	1	0	0	0	0	1	$H_{2/1}^1$
v^{10}	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^{11}	0	1	0	0	0	0	0	1	$H_{2/1}^1$
v^{12}	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	1	0	0	0	0	0	0	1	$H_{2/1}^1$
v^{14}	1	0	0	0	0	0	1	0	$H_{2/1}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F: $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		H_6^+	H_6^-	H_7^+	H_7^-	H_8^+	H_8^-	P_6	P_7	P_8
dim		2	2	2	2	4	4	2	2	4
F_4	2	1	1	1	1	1	1	-1	-1	-1
F_5F_6	2	0	0	0	0	1	1	0	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 10.

	H_6^+	H_6^-	H_7^+	H_7^-	H_8^+	H_8^-	P_6	P_7	P_8	Bipartite graph
	2	2	2	2	4	4	2	2	4	
v^1	0	0	0	0	0	1	0	0	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	0	0	1	$H_{2/2}^1$
v^3	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^4	0	0	0	1	0	0	1	0	0	$H_{2/1}^1$
v^5	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^6	0	0	1	0	0	0	1	0	0	$H_{2/1}^1$
v^7	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^8	0	1	0	0	0	0	1	0	0	$H_{2/1}^1$
v^9	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	1	0	0	0	0	0	1	0	0	$H_{2/1}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D : $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep	P_6	P_7	P_8	N_5^+	N_5^-
dim	2	2	4	2	2
D_5	2	1	1	2	-1 -1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 7.

	P_6	P_7	P_8	N_5^+	N_5^-	Bipartite graph
	2	2	4	2	2	
v^1	0	0	1	0	2	$H_{3/2}^1$
v^2	0	0	1	1	1	$H_{3/2}^1$
v^3	0	0	1	2	0	$H_{3/2}^1$
v^4	0	1	0	0	1	$H_{2/1}^1$
v^5	0	1	0	1	0	$H_{2/1}^1$
v^6	1	0	0	0	1	$H_{2/1}^1$
v^7	1	0	0	1	0	$H_{2/1}^1$

SG 230

Space group $Ia\bar{3}d$, No. 230, with SOC

Σ

High symmetry line $\Sigma(0, 0, u)$, high symmetry points at both ends of Σ : $\Gamma(0, 0, 0), N(0, 0, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	N_3N_4
dim	2	2	2	2	4	4	4	
Σ_5	2	1	1	1	1	2	2	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^7$ and the corresponding bipartite graph, number of Hilbert basis: 12.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	N_3N_4	
	2	2	2	2	4	4	4	Bipartite graph
v^1	0	0	0	0	0	1	1	$H_{2/2}^1$
v^2	0	0	0	0	1	0	1	$H_{2/2}^1$
v^3	0	0	0	2	0	0	1	$H_{3/2}^1$
v^4	0	0	1	1	0	0	1	$H_{3/2}^1$
v^5	0	0	2	0	0	0	1	$H_{3/2}^1$
v^6	0	1	0	1	0	0	1	$H_{3/2}^1$
v^7	0	1	1	0	0	0	1	$H_{3/2}^1$
v^8	0	2	0	0	0	0	1	$H_{3/2}^1$
v^9	1	0	0	1	0	0	1	$H_{3/2}^1$
v^{10}	1	0	1	0	0	0	1	$H_{3/2}^1$
v^{11}	1	1	0	0	0	0	1	$H_{3/2}^1$
v^{12}	2	0	0	0	0	0	1	$H_{3/2}^1$

Δ

High symmetry line $\Delta(u, -u, u)$, high symmetry points at both ends of Δ : $\Gamma(0, 0, 0), H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	H_5	H_6H_7
dim	2	2	2	2	4	4	4	8	
Δ_6	2	1	1	0	0	1	1	-1	-2
Δ_7	2	0	0	1	1	1	1	-1	-2

$\ker M \cap \mathbb{Z}_{\geq 0}^8$ and the corresponding bipartite graph, number of Hilbert basis: 26.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	H_5	H_6H_7	
	2	2	2	2	4	4	4	8	Bipartite graph
v^1	0	0	0	0	0	1	1	0	$H_{2/2}^1$
v^2	0	0	0	0	0	2	0	1	$H_{3/4}^1$
v^3	0	0	0	0	1	0	1	0	$H_{2/2}^1$
v^4	0	0	0	0	1	1	0	1	$H_{3/4}^1$
v^5	0	0	0	0	2	0	0	1	$H_{3/4}^1$
v^6	0	1	0	1	0	0	1	0	$H_{3/2}^1$
v^7	0	1	0	1	0	1	0	1	$H_{4/4}^1$
v^8	0	1	0	1	1	0	0	1	$H_{4/4}^1$

v^9	0	1	1	0	0	0	1	0	$H_{3/2}^1$
v^{10}	0	1	1	0	0	1	0	1	$H_{4/4}^1$
v^{11}	0	1	1	0	1	0	0	1	$H_{4/4}^1$
v^{12}	0	2	0	2	0	0	0	1	$H_{5/4}^1$
v^{13}	0	2	1	1	0	0	0	1	$H_{5/4}^1$
v^{14}	0	2	2	0	0	0	0	1	$H_{5/4}^1$
v^{15}	1	0	0	1	0	0	1	0	$H_{3/2}^1$
v^{16}	1	0	0	1	0	1	0	1	$H_{4/4}^1$
v^{17}	1	0	0	1	1	0	0	1	$H_{4/4}^1$
v^{18}	1	0	1	0	0	0	1	0	$H_{3/2}^1$
v^{19}	1	0	1	0	0	1	0	1	$H_{4/4}^1$
v^{20}	1	0	1	0	1	0	0	1	$H_{4/4}^1$
v^{21}	1	1	0	2	0	0	0	1	$H_{5/4}^1$
v^{22}	1	1	1	1	0	0	0	1	$H_{5/4}^1$
v^{23}	1	1	2	0	0	0	0	1	$H_{5/4}^1$
v^{24}	2	0	0	2	0	0	0	1	$H_{5/4}^1$
v^{25}	2	0	1	1	0	0	0	1	$H_{5/4}^1$
v^{26}	2	0	2	0	0	0	0	1	$H_{5/4}^1$

Λ

High symmetry line $\Lambda(u, u, u)$, high symmetry points at both ends of Λ : $\Gamma(0, 0, 0), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$
A matrix

corep		Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	P_4P_5	P_6	P_7P_8
	dim	2	2	2	2	4	4	2	2	6
Λ_4	2	1	1	1	1	1	1	0	-1	-2
$\Lambda_5\Lambda_6$	2	0	0	0	0	1	1	-1	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^9$ and the corresponding bipartite graph, number of Hilbert basis: 17.

	Γ_6^+	Γ_6^-	Γ_7^+	Γ_7^-	Γ_8^+	Γ_8^-	P_4P_5	P_6	P_7P_8	Bipartite graph
	2	2	2	2	4	4	2	2	6	
v^1	0	0	0	0	0	1	1	1	0	$H_{3/2}^1$
v^2	0	0	0	0	0	2	1	0	1	$H_{4/4}^4$
v^3	0	0	0	0	1	0	1	1	0	$H_{3/2}^1$
v^4	0	0	0	0	1	1	1	0	1	$H_{4/4}^4$
v^5	0	0	0	0	2	0	1	0	1	$H_{4/4}^4$
v^6	0	0	0	1	0	0	0	1	0	$H_{2/1}^1$
v^7	0	0	0	1	0	1	0	0	1	$H_{3/3}^1$
v^8	0	0	0	1	1	0	0	0	1	$H_{3/3}^1$
v^9	0	0	1	0	0	0	0	1	0	$H_{2/1}^1$
v^{10}	0	0	1	0	0	1	0	0	1	$H_{3/3}^1$

v^{11}	0	0	1	0	1	0	0	0	1	$H_{3/3}^1$
v^{12}	0	1	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{13}	0	1	0	0	0	1	0	0	1	$H_{3/3}^1$
v^{14}	0	1	0	0	1	0	0	0	1	$H_{3/3}^1$
v^{15}	1	0	0	0	0	0	0	1	0	$H_{2/1}^1$
v^{16}	1	0	0	0	0	1	0	0	1	$H_{3/3}^1$
v^{17}	1	0	0	0	1	0	0	0	1	$H_{3/3}^1$

G

High symmetry line $G(u, -u, \frac{1}{2})$, high symmetry points at both ends of G : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), N(0, 0, \frac{1}{2})$

A matrix

corep	H_5	H_6H_7	N_3N_4
dim	4	8	4
G_2G_3	2	1	2
G_4G_5	2	1	2

$\ker M \cap \mathbb{Z}_{\geq 0}^3$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	H_5	H_6H_7	N_3N_4	
	4	8	4	Bipartite graph
v^1	0	1	2	$H_{3/4}^1$
v^2	1	0	1	$H_{2/2}^1$

F

High symmetry line $F(u + \frac{1}{4}, \frac{1}{4} - 3u, u + \frac{1}{4})$, high symmetry points at both ends of F : $H(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}), P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$

A matrix

corep	H_5	H_6H_7	P_4P_5	P_6	P_7P_8
dim	4	8	2	2	6
F_4F_5	2	0	2	-1	0
F_6	2	2	2	0	-1

$\ker M \cap \mathbb{Z}_{\geq 0}^5$ and the corresponding bipartite graph, number of Hilbert basis: 4.

	H_5	H_6H_7	P_4P_5	P_6	P_7P_8	
	4	8	2	2	6	Bipartite graph
v^1	0	1	1	0	1	$H_{3/4}^2$
v^2	0	1	2	2	0	$H_{5/4}^1$
v^3	1	0	0	2	0	$H_{3/2}^1$
v^4	1	1	0	0	2	$H_{4/6}^1$

D

High symmetry line $D(u, u, \frac{1}{2} - u)$, high symmetry points at both ends of D: $P(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}), N(0, 0, \frac{1}{2})$
A matrix

corep		P_4P_5	P_6	P_7P_8	N_3N_4
	dim	2	2	6	4
D_2D_3	2	1	1	1	-1
D_4D_5	2	0	0	2	-1

ker $M \cap \mathbb{Z}_{\geq 0}^4$ and the corresponding bipartite graph, number of Hilbert basis: 2.

	P_4P_5	P_6	P_7P_8	N_3N_4	
	2	2	6	4	Bipartite graph
v^1	0	1	1	2	$H_{4/4}^4$
v^2	1	0	1	2	$H_{4/4}^4$

-
- [1] C. Bradley and A. Cracknell, *Mathematical theory of symmetry in solids: representation theory for point groups and space groups* (OUP, Oxford, 2009).
- [2] G.-B. Liu, M. Chu, Z. Zhang, Z.-M. Yu, and Y. Yao, SpaceGroupIrep: A package for irreducible representations of space group, *Computer Physics Communications* **265**, 107993 (2021).
- [3] G.-B. Liu, Z. Zhang, Z.-M. Yu, and Y. Yao, MSGCorep: A package for corepresentations of magnetic space groups, *Computer Physics Communications* **288**, 108722 (2023).
- [4] Bilbao Crystallographic Server, <https://www.cryst.ehu.es/>.
- [5] W. Bruns and B. Ichim, Normaliz: Algorithms for affine monoids and rational cones, *Journal of Algebra* **324**, 1098 (2010).
- [6] Z. Zhang, Z.-M. Yu, G.-B. Liu, Z. Li, S. A. Yang, and Y. Yao, *MagneticKP: A package for quickly constructing kp models of magnetic and non-magnetic crystals* (2022), number: arXiv:2205.05830 arXiv:2205.05830 [cond-mat].
- [7] H. C. Po, A. Vishwanath, and H. Watanabe, Symmetry-based indicators of band topology in the 230 space groups, *Nature Communications* **8**, 931 (2017).
- [8] Z. Zhang, Z.-M. Yu, G.-B. Liu, and Y. Yao, MagneticTB: A package for tight-binding model of magnetic and non-magnetic materials, *Computer Physics Communications* **270**, 108153 (2022).
- [9] I. R. Beattie, P. J. Jones, G. Reid, and M. Webster, The Crystal Structure and Raman Spectrum of $\text{Ge}_5\text{Cl}_{12} \cdot \text{GeCl}_4$ and the Vibrational Spectrum of Ge_2Cl_6 , *Inorganic Chemistry* **37**, 6032 (1998).
- [10] G. Kresse and J. Hafner, Ab initio molecular-dynamics simulation of the liquid-metal–amorphous-semiconductor transition in germanium, *Physical Review B* **49**, 14251 (1994), publisher: American Physical Society.
- [11] G. Kresse and J. Furthmüller, Efficient iterative schemes for ab initio total-energy calculations using a plane-wave basis set, *Physical Review B* **54**, 11169 (1996), publisher: American Physical Society.
- [12] J. P. Perdew, K. Burke, and M. Ernzerhof, Generalized Gradient Approximation Made Simple, *Physical Review Letters* **77**, 3865 (1996), publisher: American Physical Society.
- [13] T. M. Project, Materials Data on Ge_3Cl_8 by Materials Project [10.17188/1204166](https://doi.org/10.17188/1204166) (2017).
- [14] J. Gao, Q. Wu, C. Persson, and Z. Wang, Irvsp: to obtain irreducible representations of electronic states in the VASP, *Computer Physics Communications* **261**, 107760 (2021), arXiv: 2002.04032.